SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |16-01-2026|

THE IMPACT OF POLYPHARMACY ON RENAL FUNCTION IN THE ELDERLY AND EARLY CLINICAL MONITORING STRATEGIES

Panjiyev Asilbek G'ayrat o'g'li
Student of General Medicine, Faculty of Medicine
Termiz university of Economics and Service
Email: panjiyevasilbek66@mail.com

Eshqobilov Ozodbek Abdusodiqovich
Assistant, Department of Fundamental Medical Sciences
Termiz university of Economics and Service
Email: ozotbekbestboy@gmail.com

ABSTRACT

Polypharmacy, defined as the simultaneous use of multiple medications, is a widespread and complex clinical problem in modern medicine, particularly among elderly patients. Age-related physiological changes, especially the decline in renal function, significantly increase the nephrotoxic effects of medications in the setting of polypharmacy. This article analyzes the mechanisms through which polypharmacy affects renal function in the elderly, as well as the associated clinical risks and strategies for early detection and monitoring. The study findings demonstrate that optimization of drug therapy, regular laboratory monitoring, and an individualized clinical approach play a crucial role in preventing renal injury in elderly patients.

Keywords: polypharmacy, elderly age, renal function, nephrotoxicity, clinical monitoring.

INTRODUCTION

In recent decades, a significant increase in the proportion of elderly individuals has been observed worldwide as a result of rising life expectancy. This demographic shift has placed new and complex demands on healthcare systems. Elderly patients frequently present with multiple chronic conditions simultaneously, which often necessitates the long-term use of several medications. As a result, polypharmacy emerges as a common clinical phenomenon, defined as the regular use of multiple drugs at the same time. Although polypharmacy may be clinically justified in certain situations, it substantially increases the risk of adverse drug reactions, drug–drug interactions, and toxic effects on target organs.

The kidneys play a central role in the elimination of medications and their metabolites from the body. Therefore, under conditions of polypharmacy, the kidneys are among the organs most vulnerable to drug-induced injury. In elderly individuals, physiological age-related changes in renal function are commonly observed, including a gradual decline in glomerular filtration rate, reduced renal blood flow, and diminished tubular secretion and reabsorption capacity. Against this background, medications may accumulate in the body, reach toxic concentrations, and lead to the development of subclinical or clinically significant renal damage.


Clinical practice indicates that drug-related renal injury in elderly patients is often detected at a late stage. This delay is attributed, on the one hand, to the paucity of specific clinical symptoms and, on the other hand, to the limited sensitivity of traditional laboratory markers such as serum creatinine in reflecting true renal function in elderly individuals. Consequently, renal damage is frequently identified only after significant functional impairment has already occurred, thereby limiting therapeutic options and worsening prognosis. For these reasons, early assessment and

SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |16-01-2026|

dynamic monitoring of renal function in elderly patients receiving multiple medications are of critical clinical importance.

Literature Review: Polypharmacy is widely recognized as one of the most prevalent and clinically challenging problems in geriatric medicine. In the literature, polypharmacy is commonly defined as the regular use of five or more medications by a single patient. However, modern approaches emphasize that polypharmacy should be evaluated not only based on the number of medications but also considering the appropriateness of drug selection, potential drug–drug interactions, dosage adequacy, and alignment with the patient's functional status.

Elderly patients frequently suffer from multiple chronic diseases such as arterial hypertension,




diabetes mellitus, ischemic heart disease, arthritis, chronic obstructive pulmonary disease, and dyslipidemia, which makes polypharmacy almost inevitable. However, studies consistently report that polypharmacy significantly increases the incidence of adverse drug reactions and drug-related hospitalizations.

A decline in renal function is considered one of the key factors amplifying the risks associated with polypharmacy in elderly individuals. Age-related reductions in glomerular filtration rate, renal blood flow, and tubular function lead to decreased drug clearance. As a result, even standard drug dosages may produce higher systemic exposure in elderly patients, increasing the likelihood of nephrotoxicity. Clinically, this process often develops gradually and without overt symptoms, contributing to delayed diagnosis of renal impairment.

METHODOLOGY

This study employed an analytical and systematic approach to assess the impact of polypharmacy on renal function in elderly patients and to substantiate early clinical monitoring strategies. The study design was clinically oriented and based on the principles of evidence-based

SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |16-01-2026|



medicine. Observational and comparative analysis methods were used, and existing scientific data on the mechanisms of drug-induced renal injury associated with polypharmacy were systematized.

The methodology consisted of several key stages. In the first stage, clinical criteria for defining polypharmacy in elderly patients were established. The number of medications taken regularly, their pharmacological classes, duration of use, and potential drug–drug interactions were evaluated. Polypharmacy was defined as the continuous use of five or more medications, and the impact of drug burden on renal function was analyzed on an individual basis.

In the second stage, parameters for assessing renal function were selected. Key indicators included glomerular filtration rate, serum creatinine, cystatin C levels, and urinary albumin excretion. When estimating glomerular filtration rate, factors such as age, sex, and body mass were taken into account, and dynamic monitoring results were considered particularly important. Minimal changes in laboratory parameters were also regarded as diagnostically significant for identifying subclinical renal injury.

RESULTS AND DISCUSSION

The results of the analysis demonstrate that polypharmacy exerts a significant and multifactorial impact on renal function in elderly patients. Age-related physiological decline in renal reserve, when combined with the simultaneous use of multiple medications, creates favorable conditions for the development of drug-induced kidney injury. The findings indicate that elderly patients receiving five or more medications show a higher frequency of reduced glomerular filtration rate and subtle laboratory signs of renal impairment compared with those receiving fewer drugs. Importantly, these changes often occur in the absence of pronounced clinical symptoms, which explains the frequent delay in diagnosis.

One of the key findings is that nephrotoxic risk increases not only due to the number of medications but also because of inappropriate drug combinations and insufficient dose adjustment according to renal function. Drugs commonly used in elderly patients—such as nonsteroidal anti-inflammatory drugs, certain antihypertensive agents, diuretics, and antimicrobial medications—were found to contribute cumulatively to renal stress. When administered together, these agents may reduce renal perfusion, alter intraglomerular hemodynamics, and promote tubular toxicity. As a result, even

standard therapeutic doses may lead to excessive drug accumulation and progressive renal damage in elderly individuals.

The analysis also highlights the limited diagnostic value of serum creatinine alone in elderly patients. Due to age-related reductions in muscle mass, creatinine levels may remain within normal ranges despite a significant decline in glomerular filtration rate. This finding underscores the importance of using estimated glomerular filtration rate calculations and additional biomarkers, such as cystatin C and urinary albumin excretion, for early detection of renal dysfunction in the context of polypharmacy. Regular monitoring of these parameters allows clinicians to identify subclinical renal injury at an early stage and adjust therapy accordingly.

CONCLUSION

Polypharmacy has become an integral component of modern clinical practice in elderly patients due to the high prevalence of chronic diseases and the need for long-term treatment. However, it represents a significant risk factor for renal dysfunction, as age-related physiological decline reduces renal reserve and alters the pharmacokinetic and pharmacodynamic properties of medications.

The analysis confirms that renal injury associated with polypharmacy often progresses silently and is diagnosed at advanced stages, resulting in poorer clinical outcomes. Multiple interrelated mechanisms contribute to renal impairment under conditions of polypharmacy, including drug–drug interactions, combined use of nephrotoxic agents, reduced renal perfusion, and age-related metabolic changes. Notably, even standard medication doses may exert toxic effects in elderly patients with reduced glomerular filtration rate.

Early assessment of renal function, optimization of drug regimens, and continuous clinical and laboratory monitoring are essential strategies for preventing renal injury and improving prognosis in elderly patients exposed to polypharmacy.

REFERENCES

1. Qodirov A.A. Clinical course of kidney diseases and modern diagnostic approaches. Tashkent: Abu Ali ibn Sino Medical Publishing House, 2019.
2. Karimov B.B., Rahmonov D.T. Clinical features of pharmacotherapy for chronic diseases in the elderly. Uzbekistan Medical Journal, 2020; No. 3.
3. To‘raqulov N.M. Pathophysiological basis and clinical consequences of chronic renal failure. Tashkent: Medical Publishing House, 2018.
4. Ismoilov J.X., Abdullayev S.R. Early signs of drug-induced renal injury. Medicine and Innovations, 2021; No. 2.
5. Mamatqulov O.Sh., Yo‘ldoshev A.N. Renal dysfunction in arterial hypertension. Clinical Medicine, 2019; No. 6.
6. Rahimova M.K. Optimization of drug therapy and the problem of polypharmacy in elderly patients. Uzbekistan Health Journal, 2021; No. 4.
7. Xolmatov F.S., Qosimova M.N. Clinical significance of albuminuria and glomerular filtration rate. Uzbekistan Nephrology Bulletin, 2022; No. 1.
8. Abdurahmonov U.A. Functional relationship between the kidney and the heart (cardiorenal syndrome). Tashkent: Science and Technology, 2020.
9. Saidov A.M., Tursunova L.K. Modern laboratory diagnostic methods in kidney diseases. Tashkent: Innovative Development Publishing House, 2021.
10. Nurmatov K.Sh. Prevention and clinical monitoring of chronic kidney diseases in the elderly. Republican Scientific and Practical Medical Journal, 2022; No. 2.

JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |16-01-2026|

11. Jo‘rayev B.E., Hamroyev S.S. Nephrotoxic effects of medications and their clinical significance. Theory and Practice of Medicine, 2020; No. 5.
12. Usmonov A.I. Age-related changes in renal function in gerontology. Tashkent, 2019.