SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |16-01-2026|

THE ROLE OF GENETIC FACTORS IN THE DEVELOPMENT OF PSORIASIS

Azimova Umida Panji qizi
Student of Termiz university of Economics and Service

Eshqobilov Ozodbek Abdisodiqovich
Lecturer, Faculty of Medicine
Termiz university of Economics and Service
E-mail: ozotbekbestboy@gmail.com

ABSTRACT

This article examines the relationship between psoriasis and genetic factors, as well as the causes and types of the disease. Research indicates that psoriasis is a hereditary condition, and genetic predisposition is one of the key factors in its development. The study also analyzes the combined effects of genetic factors with environmental influences, immune system disorders, and stress. In addition, the article provides information on secondary diseases that may develop as a result of psoriasis, as well as various conditions that contribute to the onset and progression of the disease.

Keywords: psoriasis, genetics, immune system, lymphocytes, heredity, stress, multifactorial, epidermis, keratinization, hyperproliferation, neurogenic, virological, fat, carbohydrate, protein, fungus, antidepressant, allergy, external factors, T-lymphocytes.

INTRODUCTION

Psoriasis is one of the most widespread chronic dermatological diseases. It is not contagious; however, it is closely associated with genetic factors. The disease is characterized by the appearance of slightly elevated, round or oval-shaped, red or pink lesions on the skin surface. These lesions often increase as a result of scratching and gradually enlarge, becoming covered with whitish or silvery scaling plaques. A seasonal pattern is typical for psoriasis, with exacerbations most commonly occurring in autumn and winter, while partial remission is often observed during the summer months.

Unlike some other skin diseases, itching may not always be present in psoriasis. Lesions are most frequently localized on the knees, elbows, and scalp. Psoriasis is considered a multifactorial disease, and its pathogenesis is complex. It is manifested by hyperproliferation of epidermal cells and disturbances in the keratinization process, while inflammatory reactions develop in the dermis. In addition, psoriasis may be accompanied by pathological changes in various organs and systems.

Scientific studies indicate that genetic factors play a significant role in the development of psoriasis. Approximately 40% of cases are considered to be hereditary in nature. Psoriasis belongs to the group of diseases that can be transmitted from generation to generation. According to statistical data, 20–37% of patients with psoriasis have relatives on the paternal side who are affected by the disease. If one of the parents has psoriasis, the probability of transmission to the child is about 25%. In cases where both parents are affected, the risk of psoriasis in the offspring increases to approximately 75%. If psoriasis occurs in a child born to healthy parents, the likelihood of the disease developing in subsequent children is estimated to be around 17%.In addition to genetic predisposition, several other factors contribute to the development of psoriasis. These include virological, neurogenic, infectious, endocrine, allergic factors, as well as metabolic disorders, particularly disturbances in lipid and carbohydrate metabolism. It is important to emphasize that psoriasis is not an infectious disease and cannot be transmitted via airborne, contact, sexual, or other routes.

In addition, the development of psoriasis may be triggered by chronic inflammatory and infectious diseases such as tonsillitis, sinusitis, cholecystitis, and adnexitis, as well as decreased

sebum production of the skin (dryness), use of low-quality cosmetic products, exposure to chemical substances used for cleaning purposes, frequent hand washing with shower gels and antibacterial soaps, excessive alcohol consumption, fungal infections, and bacterial diseases caused by Staphylococcus and Streptococcus, after which psoriasis may also develop. Other contributing factors include mechanical skin injuries, allergic predisposition, drug abuse, obesity, smoking, HIV infection, and the use of certain medications, including antidepressants, anticonvulsants, drugs used for the treatment of cardiovascular and malarial diseases, as well as medications prescribed for mental disorders.

As mentioned above, psoriasis belongs to the group of multifactorial diseases. Multifactorial diseases develop as a result of the interaction between genetic predisposition and environmental factors. In individuals with multifactorial susceptibility, eliminating certain external factors such as meat-based foods, grilled meat products, and alcoholic beverages like wine from the diet may reduce the risk of disease development. The classification of psoriasis as a multifactorial disease is supported by the conclusions of Mordovtsev and several other researchers who studied the disease. In particular, epidemiological studies conducted among the population of the Faroe Islands revealed that 2.84% of the inhabitants were affected by psoriasis, and in 91% of these cases, a family history of psoriasis was identified, indicating a strong genetic component.

When examining other causative factors in greater depth, immune system dysfunction occupies a central role. In this condition, improper functioning of the immune system leads to excessive activation and proliferation of T-lymphocytes, which migrate to the epidermis and initiate inflammatory processes. To date, the exact mechanisms underlying these immune alterations and the reasons for increased T-lymphocyte activation remain unclear. However, it has been established that a combination of genetic predisposition and environmental factors partially contributes to these changes. Hormonal fluctuations also play a role in the onset of psoriasis, most commonly manifesting during puberty.

Neurogenic factor (stress, psychological influences, emotional strain). This factor involves disturbances in the trophic (nutritional-regulatory) function of the central nervous system. According to some researchers, the immune system may respond to emotional and psychological pressure in a manner similar to its response to physical injury. Therefore, neurological and psychological problems may contribute to the onset of psoriasis or aggravate its course. In patients with psoriasis, disorders of the central and autonomic nervous systems—such as neurosis, diencephalitis, and vegetative dystonia—have been reported. Clinically, these conditions may manifest with a marked increase in biologically active substances in the body.

Metabolic Disorders. Disruptions in carbohydrate, protein, and lipid metabolism may lead to functional changes in the formation of the stratum corneum and in the normal process of epidermopoiesis. In individuals with psoriasis, an increase in prostaglandins and polyamines has been observed, whereas vitamin D3 levels tend to decrease.

Endocrine system changes. Psoriasis may also arise as a result of functional alterations in endocrine glands. Clinical manifestations are often noted in pregnant women. The first half of pregnancy, abortion, childbirth, and the end of lactation may trigger psoriasis exacerbations and contribute to a more severe disease course.

Infectious–allergic factor. Psoriasis may be associated with chronic infectious diseases, viruses, microbes, and allergic reactions to products formed during microbial activity. Chronic infections can weaken the immune system and thereby create conditions for easier penetration of potential psoriatic triggers into the body. In addition, the entry of such agents may provoke hypersensitivity reactions, which can also contribute to disease development.

SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |16-01-2026|

Secondary conditions in psoriasis. Patients with psoriasis may develop secondary complications such as arthritis, joint pain, cardiovascular diseases, and others. Psychological problems are also common: due to visible skin damage, patients may experience low self-esteem, depression, stress, and other mental health disturbances.

Common sites of psoriasis involvement

Psoriasis most frequently affects:

Elbows,

Knees,

Trunk,

Palms and soles,

Skin folds,

Scalp,

Fingernails and toenails,

Face,

Genital area.

Stages of the disease

Psoriasis progresses through several stages.

Progressive stage: In this stage, round or oval plaque-like lesions appear. They tend to enlarge and are typically erythematous (red). Scaling resembling dandruff may be present in the central areas. Some patients experience itching during this period. In addition, several characteristic clinical phenomena may be observed.

Stationary stage. The stationary stage usually develops 1–4 weeks after the initial manifestations of the disease. During this stage, no new lesions appear. Existing plaques become larger and acquire a pale pink coloration. Scaling is pronounced, while itching is minimal or almost absent.

Regressive stage. In the regressive stage, scaling gradually decreases. Lesions begin to resolve from the central part, forming ring-shaped patterns. Since psoriasis is a chronic disease, it often follows a cyclical course. Typically, the disease progresses through two or three stages with mild symptoms or apparent regression of lesions, followed by subsequent exacerbation. In some patients, psoriasis worsens during summer and spring, whereas in others it exacerbates in autumn and winter. If remission does not occur in any season, the condition is referred to as mixed-type psoriasis.

In certain cases, the appearance of psoriatic lesions depends on skin color. In individuals with lighter skin, lesions usually appear pink, red, or brown, while in individuals with darker skin tones they may appear purple or dark brown. In mild cases, psoriasis affects less than 5% of the body surface area; in moderate cases, 5–10%; and in severe cases, more than approximately 10% of the body surface.

Nail involvement. Psoriasis may also affect the nails, leading to discoloration, formation of small pits, and separation of the nail plate from the nail bed (onycholysis). Nail joints may also be involved, causing pain during movement. The nail surface becomes dull, thickened, and deformed.

Clinical forms of psoriasis: Seborrheic psoriasis**.**This form occurs in areas affected by seborrhea, including the face, scalp, upper chest, and back. Lesions in these areas are distinguished from healthy skin by well-demarcated plaques covered with yellowish scales.Intertriginous psoriasis:Lesions develop on flexural surfaces of the joints, in the axillary regions, and in the groin. This type is more common in individuals with rheumatism or diabetes, in infants, obese individuals, and those with hypersensitive skin. It is often accompanied by itching and burning sensations.

Psoriatic erythroderma: This form occurs in approximately 1–3% of patients with psoriasis. It is considered a severe complication of psoriasis and may develop as a result of exposure to adverse factors or improper treatment. The use of herbal baths or high-concentration topical ointments may trigger this condition. Clinical manifestations include generalized skin inflammation, enlarged lymph nodes, nail destruction, purulent paronychia, and hair loss.

Arthropathic psoriasis (psoriatic arthritis). This form primarily affects both large and small joints. Pathological changes are commonly observed in the small joints of the hands and feet, as well as in the knees and spine. It is characterized by a severe course and relatively low prevalence. Patients experience intense joint pain, which increases during movement.

Psoriasis in children. In recent years, an increasing prevalence of psoriasis among children has been noted. Approximately 50% of cases occur in children under the age of 10. In individuals with psoriasis, genes involved in immune signaling pathways become dysregulated. Globally, genes associated with a predisposition to psoriasis are found in approximately 10 out of every 100 people. During severe exacerbations, psoriasis may spread over the entire body. Self-medication may lead to adverse outcomes; therefore, patients are strongly advised to seek consultation from a qualified dermatologist as soon as symptoms appear.

REFERENCES

1. Eshboyev E.H. Skin and Venereal Diseases. Ilm-Ziyo Publishing House, Tashkent, 2017.
2. Vaisov A. Skin and Venereal Diseases. Yangi Asr Avlodi, Tashkent, 2009.
3. Mannanov A.M., Khayitov Q.N. Pediatric Skin and Venereal Diseases. Iqtisod-Moliya, Tashkent, 2016.
4. Shodiyev K., Ahmedov R. Skin and Venereal Diseases. Tashkent, 2010.
5. Olimkho‘jayev P.R., Inog‘omova D.R. Medical Genetics. Ilm-Ziyo, Tashkent, 2017.
6. Aripov S.S. Skin and Venereal Diseases. National Encyclopedia of Uzbekistan, State Scientific Publishing House, 2010.
7. Karimov A., Mirahmedov U. Skin and Venereal Diseases. Ibn Sina Publishing and Printing Association, Tashkent, 1995.
8. Sachdev P. (Reviewer), Stuart A., Ellis R.R. 7 Types of Psoriasis. April 11, 2024.
9. Internet sources
10. https://www.aad.org
11. https://www.who.int
12. https://www.hopkinsmedicine.org
13. https://www.psoriasis.org
14. https://www.pennmedicine.org
15. https://www.healthline.com
16. https://www.webmd.com
17. https://www.wikipedia.com
18. https://www.nhs.uk