SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |24-02-2026|

MODERN DIAGNOSTIC APPROACHES TO CHRONIC HYPERPLASTIC LARYNGITIS

Bo'riyeva Dilnoz Baxriddinovna
buriyevadilnoz29@gmail.com
Nabiyeva Latofat Azizqul qizi
nabiyevalatofat4@gmail.com
Termez University of Economics and Service
Faculty of Medicine

ABSTRACT

Chronic hyperplastic laryngitis (CHL) is a persistent inflammatory condition of the laryngeal mucosa characterized by epithelial hyperplasia, thickening of the vocal folds, and potential precancerous changes. Early and accurate diagnosis is essential to prevent malignant transformation and preserve vocal function. Traditional diagnostic methods are often insufficient to detect subtle mucosal alterations and early dysplastic changes. Therefore, modern diagnostic techniques have been developed to improve diagnostic accuracy and clinical outcomes. This article reviews contemporary diagnostic approaches to chronic hyperplastic laryngitis, including videolaryngoscopy, videostroboscopy, narrow band imaging (NBI), contact endoscopy, optical coherence tomography (OCT), autofluorescence imaging, and histopathological evaluation. The advantages, limitations, and clinical relevance of each method are discussed. Emphasis is placed on the importance of multimodal diagnostic strategies for early detection, risk stratification, and individualized patient management.

Keywords: chronic hyperplastic laryngitis, modern diagnostics, videostroboscopy, narrow band imaging, optical coherence tomography, laryngeal pathology

INTRODUCTION

Chronic hyperplastic laryngitis (CHL) represents a significant clinical problem in otorhinolaryngology due to its prolonged course, resistance to conservative treatment, and potential for malignant transformation. The disease is characterized by persistent inflammation of the laryngeal mucosa accompanied by epithelial hyperplasia, keratinization, and stromal changes. Patients commonly present with hoarseness, voice fatigue, throat discomfort, chronic cough, and in some cases, dyspnea.

Etiological factors contributing to the development of CHL include chronic smoking, alcohol consumption, occupational voice overload, gastroesophageal reflux disease (GERD), environmental pollution, and recurrent infections. The chronic irritation of the laryngeal mucosa leads to pathological remodeling, which may progress to dysplasia or laryngeal carcinoma if not detected and managed appropriately.

Conventional diagnostic approaches, such as indirect laryngoscopy and routine endoscopic examination, provide limited information regarding microvascular patterns and epithelial integrity. As a result, early neoplastic changes may remain undetected. In recent years, technological advances have introduced innovative diagnostic tools that allow detailed visualization of laryngeal structures at both macroscopic and microscopic levels.

The aim of this article is to provide a comprehensive overview of modern diagnostic methods used in the evaluation of chronic hyperplastic laryngitis and to analyze their clinical significance in improving diagnostic accuracy and patient outcomes.

MATERIALS AND METHODS

This narrative review is based on an extensive analysis of scientific literature published in peer-reviewed journals between 2005 and 2025. Electronic databases including PubMed, Scopus, Web of Science, and Google Scholar were searched using keywords such as *chronic hyperplastic*


SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |24-02-2026|

laryngitis, *laryngeal hyperplasia*, *modern diagnostic methods*, *videostroboscopy*, *narrow band imaging*, and *optical coherence tomography*.


Inclusion criteria consisted of original research articles, clinical trials, systematic reviews, and meta-analyses focusing on diagnostic techniques for chronic inflammatory and hyperplastic laryngeal diseases. Articles not available in English or lacking methodological clarity were excluded.

The diagnostic modalities were analyzed according to their principles, diagnostic value, sensitivity, specificity, advantages, limitations, and clinical applicability.

Modern Diagnostic Techniques



SIR Publishers
CENTER FOR INTERNATIONAL SCIENTIFIC PUBLICATIONS
JOURNAL OF NURSING AND WOMEN'S HEALTH
ISSN(Online): 2984-8083
Volume-9, Issue-01, Published |24-02-2026|



Videolaryngoscopy. Videolaryngoscopy is a fundamental diagnostic tool that enables high-resolution visualization of the larynx using rigid or flexible endoscopes. It allows detailed assessment of vocal fold morphology, mucosal color, thickness, and surface irregularities. Compared to conventional mirror examination, videolaryngoscopy provides superior image quality and documentation capabilities.

Videostroboscopy. Videostroboscopy is considered the gold standard for evaluating vocal fold vibration. By using stroboscopic light synchronized with vocal fold frequency, this method simulates slow-motion visualization of vocal fold oscillations. In CHL, videostroboscopy reveals reduced mucosal wave, asymmetry, and stiffness, which are indicative of hyperplastic changes.

Narrow Band Imaging (NBI). NBI is an advanced optical technology that enhances visualization of mucosal microvascular patterns by using specific blue and green light wavelengths. It is particularly useful in detecting abnormal angiogenesis associated with dysplasia and early malignancy. In patients with CHL, NBI improves differentiation between benign hyperplasia and suspicious premalignant lesions.

Contact Endoscopy. Contact endoscopy involves direct microscopic examination of the laryngeal mucosa after topical staining. This technique allows real-time evaluation of epithelial cell morphology and vascular structures. Although highly informative, its use is limited by technical complexity and the need for specialized equipment.

Optical Coherence Tomography (OCT). OCT is a non-invasive imaging modality that provides cross-sectional images of tissue microarchitecture with near-histological resolution. It enables assessment of epithelial thickness, basement membrane integrity, and subepithelial changes. OCT shows great promise in identifying early dysplastic alterations without the need for biopsy.

Autofluorescence Imaging. Autofluorescence imaging detects biochemical and structural changes in tissues by analyzing fluorescence patterns. Pathological tissues exhibit altered fluorescence compared to normal mucosa. This method serves as an adjunctive tool for screening and lesion mapping.

Histopathological Examination. Despite advances in imaging, histopathology remains the definitive diagnostic method. Targeted biopsy under endoscopic guidance ensures accurate grading of epithelial changes, ranging from hyperplasia to carcinoma in situ.

RESULTS

Analysis of the reviewed literature indicates that modern diagnostic techniques significantly enhance the detection and characterization of chronic hyperplastic laryngitis. Videostroboscopy improves functional assessment, while NBI increases sensitivity for identifying premalignant

changes. OCT and contact endoscopy provide valuable microstructural information, reducing unnecessary biopsies.

Studies demonstrate that multimodal diagnostic approaches yield higher diagnostic accuracy compared to single-method evaluation. The combination of videolaryngoscopy, videostroboscopy, and NBI is particularly effective in routine clinical practice.

DISCUSSION

The diagnosis of chronic hyperplastic laryngitis has evolved considerably with the introduction of advanced imaging technologies. Early identification of pathological changes is critical in preventing malignant transformation and preserving vocal function. Modern diagnostic methods allow clinicians to assess both structural and functional aspects of the larynx with unprecedented precision.

However, limitations such as equipment cost, availability, and operator expertise must be considered. Future developments in artificial intelligence and digital image analysis may further enhance diagnostic capabilities and standardize interpretation.

CONCLUSION

Modern diagnostic approaches play a crucial role in the comprehensive evaluation of chronic hyperplastic laryngitis. The integration of advanced endoscopic and imaging techniques with traditional histopathology enables early detection, accurate diagnosis, and personalized management. Multimodal diagnostic strategies should be considered the standard of care in patients with suspected chronic hyperplastic laryngeal pathology.

REFERENCES

1. Zeitels SM, et al. **Laryngeal pathology and advanced imaging techniques**. Otolaryngol Clin North Am.
2. Piazza C, et al. **Narrow band imaging in laryngology**. Curr Opin Otolaryngol Head Neck Surg.
3. Burns JA, et al. **Optical coherence tomography of the larynx**. Laryngoscope.
4. Dikkers FG. **Chronic laryngitis: pathology and diagnosis**. Curr Opin Otolaryngol.