ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

UDC: 617.55-089.844/616-056.5+616-08-039.71

OBESITY CORRECTION BY SURGICAL METHODS

Khamdamov I.B.

PhD, docent at the Department of Surgical Diseases

Bukhara State Medical University named after Abu Ali Ibn Sina

Summary. The LRBG operation in the correction of DLP in patients with MS is limited in application. In order to correct disorders in the lipid spectrum, the operation is advisable to use only in young age groups, among women, with an initial BMI of no more than 43 kg / m2, not "sweet tooth", capable of following all the recommendations of the attending physician for a long period of time. The use of this type of operation in men is not advisable, since the effectiveness of the operation, after a year of observation, returns to the original data, while their quality of life is significantly reduced.

Key words: obesity, obesity treatment, bariatric surgery, laparoscopic longitudinal resection of the stomach, bariatric surgery, sleeve gastrectomy.

Relevance. The WHO European Commission report (2007) "The Obesity Problem in the WHO European Region and a Strategy for Addressing It" noted that the obesity epidemic in Europe is an unprecedented, underestimated, poorly understood health problem with significant economic consequences [1,2,6,8,10,12,14,16,18]. In European countries, 30-80% of adults and up to one third of children suffer from problems associated with excess body weight [3, 4,20,21,22,23,24,25,26]. There is a trend towards an increasing prevalence of obesity among children and adolescents, which intensifies the obesity epidemic in the next generation. Overweight and obesity are the main risk factors for the development of metabolic syndrome (MS), the main components of which are arterial hypertension, type 2 diabetes mellitus, dyslipidemia (DLP) and other serious diseases [11, 13,15, 17,19,27,28,29,30].

The components of MS have a negative impact on health, quality of life of patients, and their complications are the main causes of premature mortality of the population. Elimination of obesity and related complications is a determining factor in treatment. Often, they are guided by the erroneous idea that some weight loss is “enough” to ensure a positive effect. The results of conservative methods of treating obesity allow you to reduce body weight, as a result of which improvements are observed in the course of a number of concomitant diseases, but it is almost impossible to achieve a sustainable result, especially in people with morbid obesity.

Currently, doctors of various specialties are involved in the treatment of MS. In Russia, there is no single service that includes multidisciplinary teams of specialists with the appropriate qualifications, which would allow the most effective approach to solving this problem with long-term treatment and rehabilitation of patients.

Surgical treatment of the main components of MC has already proven its effectiveness and is the most effective way to treat this category of patients. However, surgical treatment methods are effective when all other therapeutic methods have proven ineffective.

Many aspects of surgical treatment remain unresolved. The effects of bariatric procedures on the course of MC have not been clarified or remain controversial. Improvement of technology and a

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

deeper understanding of the processes occurring after the use of surgical methods of treating such patients are required, which is relevant.

The aim of the work: to evaluate the effectiveness of bariatric surgeries in the correction of DLP, to improve the treatment results of patients with MC.

Materials and methods. Of the 457 operated patients who underwent various types of bariatric interventions (2014-2023), a group of 128 patients with a diagnosis of MC was formed. In all cases, there was obesity of varying degrees in combination with DLP. Further, the results of the influence of various types of bariatric surgeries on DLP and obesity were analyzed and evaluated. During the clinical and biochemical research methods, the results were subjected to multifactorial assessment taking into account the type of surgery performed, age and gender over the S-year observation period. All the results obtained were systematized and analyzed using statistical and mathematical methods.

Research results. As shown by the study, the effectiveness of the RBG operation on the level of DLP in patients with MC in terms of the TC indicator is statistically insignificant ($p < 0.05$). In some patients, the indicator has a separate character, correlated with the initial body weight. This pattern is dominant in the analysis of changes in the total cholesterol in individuals in the older age group. There are no statistically significant changes in the lipid spectrum, which is also typical for the Ka and TG indicators. Consequently, heavy fatty acids do not undergo significant changes throughout the observation period, although the indicators decrease in the first year of observation, but subsequently restore their original values. Thus, energy adaptation of the body occurs due to "light" fats. This type of response of the body does not allow the use of RBZh surgery to correct DLP. BMI reduction is achieved by the 1st year of observation and is 31.6 ^ -36.4 with Me = 34.02. In the 3rd and 5th years, the indicator increases again, which is confirmed by the dynamics of % EBMIL. The most significant indicators equal to 53% were noted after a year of observation among patients under 40 years of age, female, and extremely insignificant - in the older age group. Thus, the operation of the LRGB in young age groups, in women, has limitations in use, in other cases it is ineffective and cannot be recommended for use to correct DLP in patients with MS. After the operation of the LRGB, the absence of statistically significant differences in relation to the initial data was established, revealed regardless of age only in the group of men by the BMI indicator ($p = 0.188134$), at the age of up to 40 years ($p = 0.103114$). At the age over 40 years - after 5 years ($p = 0.057$) and by the HDL indicator ($p = 0.062$) after 3 months and 5 years ($p = 0.07$). In other cases, statistically significant differences were found for all indicators in women and for other indicators in men, regardless of their age group. Changes in TC have no gender differences, except for long-term observation periods. From the 3rd month of observation, regardless of gender, a sharp decrease in the TC level is noted, reaching a maximum by the 1st year of observation (a sharp restriction of nutrient intake and weight loss). In the 3rd year, in men, the TC level begins to increase and reaches the upper limit of the reference values, and manifestations of DLP appear again. The integral indicator, Ka, in men, starting from the 3rd year of observation increases, reaching a maximum by the 5th year, and stabilization of the lipid spectrum is not achieved, but minimization of DLP is preserved. Thus, it can be considered that the LPRG operation has a relatively positive effect on the normalization of the lipid spectrum and neutralization of DLP. The decrease in the TG indicator is not systemic. Thus, among women, significant changes are achieved by the first year of observation. In men, throughout the entire observation period, there is an insignificant range. However, the dynamics of changes in the indicator, with a decrease in its level in the period from 1 to 3 years is the most significant. Then, in women, stabilization of the indicator is noted for 3-5 years. BMI clearly decreases in both men and women.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

At the same time, in women, as a result of the operation, by the 1st year of observation, a decrease in body weight to 35 kg / m2 is achieved. By the 5th year of observation - 31.28 kg / m2. In men, the maximum decrease in BMI is achieved by the first year of observation and is 34.74 kg / m2. In 3 - 5 years - there is an increase in body weight, reaching 36.55 kg / m2 by 5 years. This dynamics is confirmed by the %EBMIL index. Among men, the best results are observed in 1-3 years of observation, in women the effectiveness is slightly lower, with maximum effectiveness by 5 years of observation.

In the future, a similar picture is observed in men, with the exception of a decrease in the index by the 5th year of observation. That is, the index indicates the maximum possible effectiveness of the operation in men only by the 3rd year of observation. Thus, the presented data indicate a moderate effectiveness of this type of operation in normalizing the lipid spectrum. At the same time, weight loss helps to minimize the level of DLP, especially in women in the young age group, which determines the need for careful selection of patients for this type of operation. The obtained data determine the need for individualization in the choice of the LP operation for the purpose of correcting DLP in patients with MS, and the criteria for selecting patients should be age, gender and changes in the initial indicators.

When assessing the effectiveness of the LP operation on the level of DLP in patients with MS, it was found that in the 5-year observation period, taking into account gender and age gradations, the effectiveness of the operation as a whole is optimal, especially among women, regardless of age.

A clearly expressed tendency to a decrease in the TCS, Ka indicator is noted, especially in women. TGs are already sharply reduced by 3 months of observation. The %EBMIL indicator exceeds 160%. Among all the analyzed types of operations, these are the highest efficiency indicators. Based on the data obtained, it can be stated that the LBPSh operation has a significant effect on both obesity of varying degrees and the initial DLP, which allows us to define this type of surgical treatment as the "operation of choice" in the treatment of DLP in patients with MS. The effectiveness of the LBPSh operation, according to the main indicators, is the most significant. TC is sharply reduced by the 3rd month of observation and is maintained throughout the observation period with minimization of DLP. A sharp change in the level of TC and HDL has a direct effect on the value and dynamics of Ka, which determines its leading role in the dynamic monitoring of the lipid spectrum. An effective decrease in the TG level is also characteristic, reaching maximum values already from the 3rd month and maintaining the entire observation period. Characteristically, there is an effective and smooth decrease in BMI, with stabilization of the indicator by the 3rd year of observation. The above is confirmed by the dynamics of the %EBMIL indicator, reaching 120% already in the 1st year of observation. In subsequent years, the effectiveness of the operation only increases. Thus, based on the obtained results, it can be argued that the LRGB operation is characterized by low efficiency in relation to the impact on the level of DLP in patients with MC. Only in young age groups, in women, during the first year of observation after the operation, with an initial body weight not exceeding 43 kg / m2, with initial disturbances in the lipid spectrum, the operation can be effective. This pattern can only be associated with weight loss. In other cases, the operation is ineffective. It is important that the effect of the operation directly depends on the degree of weight loss, initial changes in the lipid spectrum, age and sex of patients. The effect of the LRGB operation on the degree of obesity and the level of DLP occurs from the first weeks after it. Apparently, such an effect of the operation is associated with a sharp (shock) restriction of the intake of nutrients into the body. The effectiveness of the operation decreases during the period of 3-5 years of observation, especially in patients of older

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

age groups and depends on the degree of weight loss, the severity of changes in the initial data. This pattern is characteristic of all restrictive types of operations and is directly related to the process of weight loss, as a result of which the volume of visceral fat depots decreases. In addition, the effect of the LPRZH operation, in addition to a significant gastro-restrictive component, is associated with the removal of the ghrelin-producing zone of the fundus of the stomach, which helps suppress the feeling of hunger and improves the indicators. As a result of the analysis, we can assume that the basis of the early effect of the LPRZH operation in relation to the correction of indicators towards their improvement is not a decrease in body weight, but a "shock" transition of patients to a low-calorie diet, and only subsequently (by the year and beyond) the effect of the operation is associated with a decrease in visceral fat mass. Subsequently, starting from the 3rd year of observation, the effectiveness of the operation decreases, which is associated with partial restoration of body weight and adaptation of the body to a limited supply of nutrients, or with the patient's inability to adapt to a new way of eating and living.

When assessing the effectiveness of combined surgeries (LZhS and LBPSh) on the level of DLP in patients with MC, it can be argued that these surgeries are the most effective throughout the observation period, in all categories of patients. At the same time, the best results are typical for young age groups.

Thus, the effectiveness of bariatric surgeries in correcting DLP in patients with MC varies and requires an individual approach to choosing one or another type of surgical intervention.

The most effective surgeries are LZhS and LBPSh, which suggests an exceptional opportunity to treat this category of patients. Bypass surgeries are highly effective and can potentially represent an alternative treatment for all patients with MC.

Conclusions: 1. The LZhS operation has significant effectiveness in correcting DLP in patients with MC throughout the observation period, especially in young age groups. It is advisable to recommend the operation for wide application as a correction of DLP in patients with MC

2. The LBPSh operation in the correction of DLP in patients with MC has the highest efficiency throughout the entire observation period. It is advisable to recommend the operation as an indication for the treatment of extreme obesity and significant DLP in patients with MC.

Literature

1. Spaziani T.S., Di Cello P., Lo Bianco G. "All In One Mesh Hernioplasty" device for inguinal hernia repair. Results of 400 cases // Ann. Ital. Chir. – 2018. – Vol. 89. – P. 438–442.

2. Steensel S., Hil L., Bloemen A. Prevention of incisional hernia using different suture materials for closing the abdominal wall: a comparison of PDS, Vicryl and Prolene in a rat model // Hernia. – 2020. – Vol. 24, № 1. – P. 67-78.

3. Trippoli S., Caccese E., Tulli G. Biological meshes for abdominal hernia: Lack of evidence–based recommendations for clinical use // Int. J. Surg. – 2018. – Vol. 52. – P. 278–284.

4. Zhang L. Incidence of abdominal incisional hernia in developing country: a retrospective cohort study // Int J Clin Exp Med. – 2015. – Vol. 8, № 8. – P. 13649-13652.

5. Zucker B.E., Simillis C., Tekkis P., Kontovounisios C. Suture choice to reduce occurrence of surgical site infection, hernia, wound dehiscence and sinus/fistula: a network meta-analysis // Ann R Coll Surg Engl. – 2019. – Vol. 101, № 3. – P. 150-161.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

6. Khamdamov I.B. Improving tactical approaches in the treatment of hernias of the anterior abdominal wall in women of fertile age // Tibbiyotda Yangi kun. Bukhoro, 2022.-№10(48)- pp. 338-342.

7. Khamdamov I.B. Morphofunctional features of the abdominal press in women of reproductive age // Tibbiyotda Yangi kun. Bukhoro, 2022.-№3(41)- pp. 223-227.

8. Khamdamov I.B. Clinical evaluation of the effectiveness of the traditional approach to the treatment of hernias of the anterior abdominal wall in women of fertile age // Doctor's Bulletin. – Samarkand 2022. No. 2.2 (104).-P.65-70.Khamdamova M.T. Ultrasound features of three-dimensional echography in assessing the condition of the endometrium and uterine cavity in women of the first period of middle age using intrauterine contraceptives // Biology va tibbyot muammolari. - Samarkand, 2020. - No. 2 (118). - P.127-131.

9. Khamdamov I.B., Khamdamov A.B. Endovideosurgical hernioplasty in women of fertile age // New day in medicine. 2021. №6 (38/1).P.25-27.

10. Khamdamov I.B., Khamdamov A.B. Classification and properties of mesh explants for hernioplasty of hernial defects of the anterior abdominal wall (review) // Biology and integrative medicine. ISSN 2181-8827 2021. №5 – March, April (52).C.12-22.

11. Khamdamova M.T., Zhaloldinova M.M., Khamdamov I.B. The state of nitric oxide in blood serum in patients with cutaneous leishmaniasis // New day in medicine. Bukhara, 2023. - No. 5 (55). - P. 638-643.

12. Khamdamova M.T., Zhaloldinova M.M., Khamdamov I.B. The value of ceruloplasmin and copper in blood serum in women wearing copper-containing intrauterine device // New day in medicine. Bukhara, 2023. - No. 6 (56). - pp. 2-7. 37.Khamdamova M. T. Bleeding when wearing intrauterine contraceptives and their relationship with the nitric oxide system // American journal of pediatric medicine and health sciences Volume 01, Issue 07, 2023 ISSN (E): 2993-2149. P.58-62

13. Khamdamova M. T. The state of local immunity in background diseases of the cervix // Eurasian journal of medical and natural sciences Innovative Academy Research Support Center.Volume 3 Issue 1, January 2023 ISSN 2181-287X P.171-175.

14. Khamdamova M.T., Khasanova M.T. Various mechanisms of pathogenesis of endometrial hyperplasia in postmenopausal women (literature review) // New day in medicine. Bukhara. 2023. - No. 8 (58). - P. 103-107.

15. Khamdamova M.T. Reproductive Health of Women Using Copper-Containing Intrauterine Contraception // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 39-45.

16. Khamdamov I.B. Advantages Of Laparoscopic Hernioplasty in Obesity Women of Fertile Age // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 33-38.

17. Khamdamova M.T., Akramova D. E. Genetic aspects of genital prolapse in women of reproductive age // New day in medicine. Bukhara, 2023. - No. 5 (55). - P. 638-643.

18. Khamdamova M.T., Teshaev Sh.Zh., Hikmatova M.F. Morphological changes of the thymus and spleen in renal failure in rats and correction with pomegranate seed oil // New day in medicine. Bukhara, 2024. - N. 3(65). - P. 167-187.

19. Khamdamova M. T. The state of local immunity in background diseases of the cervix // Eurasian journal of medical and natural sciences Innovative Academy Research Support Center.Volume 3 Issue 1, January 2023 ISSN 2181-287X R.171-175.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

20. Khamdamova M.T., Khasanova M.T. Various mechanisms of pathogenesis of endometrial hyperplasia in postmenopausal women (literature review) // New day in medicine. Bukhara. 2023. - No. 8 (58). - P. 103-107.

21. Khamdamova M.T. Reproductive Health of Women Using Copper-Containing Intrauterine Contraception // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 39-45.

22. Khamdamov I.B. Advantages Of Laparoscopic Hernioplasty in Obesity Women of Fertile Age // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 33-38.

23. Khamdamova M.T., Zhaloldinova M.M., Khamdamov I.B. The state of nitric oxide in blood serum in patients with cutaneous leishmaniasis // New day in medicine. Bukhara, 2023. - No. 5 (55). - P. 638-643.

24. Khamdamova M.T., Zhaloldinova M.M., Khamdamov I.B. The value of ceruloplasmin and copper in blood serum in women wearing copper-containing intrauterine device // New day in medicine. Bukhara, 2023. - No. 6 (56). - P. 2-7.

25. Khamdamova M. T. Bleeding when wearing intrauterine contraceptives and their relationship with the nitric oxide system // American journal of pediatric medicine and health sciences Volume 01, Issue 07, 2023 ISSN (E): P. 2993-2149. R.58-62

26. Khamdamova M.T. Reproductive Health of Women Using Copper-Containing Intrauterine Contraception // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 39-45.

27. Khamdamov I.B. Advantages Of Laparoscopic Hernioplasty in Obesity Women of Fertile Age // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 33-38.

28. Khamdamova M. T. Bleeding when wearing intrauterine contraceptives and their relationship with the nitric oxide system // American journal of pediatric medicine and health sciences Volume 01, Issue 07, 2023 ISSN (E): 2993-2149. R.58-62

29. Khamdamova M. T. The state of local immunity in background diseases of the cervix // Eurasian journal of medical and natural sciences Innovative Academy Research Support Center.Volume 3 Issue 1, January 2023 ISSN 2181-287X R.171-175.

30. Khamdamova M. T., Khasanova M.T. Genetic mechanisms of development of endometrial hyperplastic processes in women in menopacteric age // American Journal of Medicine and Medical Sciences 2025, 15(2): 372-375 DOI: 10.5923/j.ajmms. 20251502.22