ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

FEATURES OF HYPOTREMIA IN PREMATURE INFANTS WITH FREQUENT EPISODES OF APNEA

Inakova B.B.

Department of Pediatrics and Neonatology of the Faculty, ASMI

Abstract. Apnea of prematurity is defined as respiratory pauses > 20 seconds or pauses < 20 seconds associated with bradycardia (< 100 bpm 1), central cyanosis, or oxygen saturation < 85% in neonates born at < 37 weeks of gestation and without other disorders causing apnea. The cause may be immaturity of the central nervous system (central apnea); if the apneic episode is prolonged, airway obstruction may also occur. Diagnosis is based on clinical presentation and cardiorespiratory monitoring. Treatment is with stimulants for central apnea and appropriate head positioning for obstructive apnea. Prognosis is good, with apnea resolving in most preterm infants by 37 weeks postmenstrual age and in almost all preterm infants by 44 weeks postmenstrual age.

Keywords: Pathophysiology, diagnosis, treatment, prognosis, prevention, basics, additional information.

INTRODUCTION

The birth process is accompanied by extensive physiological changes in the baby (see also Pulmonary function of the newborn), which sometimes reveal conditions that were not a problem during intrauterine life. For this reason, a person with neonatal resuscitation skills should be present at each birth. Gestational age and growth parameters help determine the risk of neonatal pathology. About 25% of premature infants have apnea of prematurity, which usually begins 2–3 days after birth and only rarely on the first day. Apnea on the first day of life may indicate congenital malformations or damage to the central nervous system (CNS). Apnea that develops > 14 days after birth in an otherwise healthy infant may indicate a serious illness and should not be considered apnea of prematurity (e.g., sepsis). The risk of apnea of prematurity increases with earlier gestational age [1].

MATERIALS AND METHODS

Apnea of prematurity is a developmental disorder caused by immaturity of the neurological and/or mechanical function of the respiratory system. Apnea can be characterized as

Central (most common)

Obstructive

Mixed pattern

Central apnea is caused by immaturity of the brain centers that control breathing. The specific pathophysiology is not fully understood, but appears to be related to a number of factors, including abnormal responses to hypoxia and hypercapnia. It is the most common type of apnea in preterm infants.

Obstructive apnea is caused by an obstruction to airflow, flexion of the neck causing obstruction by subpharyngeal soft tissue, or obstruction of nasal breathing or reflex laryngospasm.

Mixed apnea is a combination of central and obstructive apnea.

All types of apnea may result in hypoxemia, cyanosis, and bradycardia if apneic episodes continue. Because bradycardia may also occur with apnea, both conditions may be due to a central mechanism. Of infants who die from sudden infant death syndrome (SIDS), 18% were born preterm, but apnea of prematurity is not a precursor to SIDS.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

Periodic breathing consists of repeated 5- to 20-second cycles of normal breathing alternating with brief (< 20 seconds) periods of apnea. This phenomenon is common among premature infants and is not considered apnea of prematurity and is of little or no clinical significance [2].

RESULTS AND DISCUSSION

Diagnosis of Apnea of Prematurity

Clinical evaluation

Cardiorespiratory monitoring

Exclude other causes (eg, hypoglycemia, sepsis, intracranial hemorrhage, gastroesophageal reflux disease)

Although apnea of prematurity is often associated with immature respiratory control mechanisms, it may be a sign of infectious, metabolic, thermoregulatory, or respiratory, cardiac, or CNS dysfunction. A thorough history, physical examination, and, if necessary, investigations should be performed before accepting prematurity as the cause of apnea. Gastroesophageal reflux disease (GERD) is no longer recognized as a cause of apnea in preterm infants, so the presence of GERD should not be considered as an explanation for apneic episodes, nor should treatment for GERD be initiated for apnea of the newborn.

Diagnosis of apnea is usually made by visual observation or by using cardiorespiratory monitors used continuously during the assessment and care of premature infants.

Treatment of apnea of prematurity

Stimulation

Treatment of underlying disorder

Respiratory stimulants (eg, caffeine)

Continuous positive airway pressure (CPAP)

Patients are admitted to the hospital for observation. When apnea is noted, either during observation or by a monitoring signal, the infant is stimulated, which may be sufficient; if breathing does not resume, bag-valve-mask ventilation is instituted (see Airways and breathing devices) [3].

Frequent or severe episodes should be promptly and thoroughly evaluated, and identified causes treated. If no infectious or other treatable underlying disorder is found, respiratory stimulants are indicated for frequent or severe episodes characterized by hypoxemia, cyanosis, bradycardia, or a combination. Oral caffeine is the safest and most commonly used inhaled stimulant. Caffeine is preferred because of ease of administration, fewer side effects, a longer therapeutic window, and less need to monitor drug levels. Treatment is continued until the infant reaches 34–35 weeks of gestation and has gone at least 5–7 days without apnea requiring intervention. Monitoring continues after the last apnea episode or after caffeine is stopped until the infant has gone an additional 5–10 days without apnea requiring intervention.

If apnea continues despite stimulants, CPAP therapy may be used, starting with pressures of 5–7 cm H2O. Poorly controlled apneic episodes require mechanical ventilation.

Rehabilitation practices vary; Some doctors observe infants for 7 days after treatment ends to ensure that apnea or bradycardia does not recur, while others continue caffeine treatment after discharge if treatment appears to be effective.

Home Monitoring

Hospitalized high-risk infants who have had no clinically significant cardiopulmonary events (e.g., apneas > 20 seconds; apneas associated with central cyanosis; apneas associated with heart rate < 80 bpm) for 3-10 days during continuous cardiorespiratory monitoring may be discharged home without

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-2

a monitor. Occasionally, the use of a cardiorespiratory monitor and/or oral caffeine at home may be indicated to shorten the length of stay in infants who are ready for discharge but still have clinically significant apneas that resolve without intervention. However, few infants are discharged home with a respiratory monitor, and only those whose apneic episodes resolve spontaneously and without intervention, including stimulation, are considered for discharge from the hospital with a monitor. Parents should be trained in the proper use of equipment, assessment of emergency situations, provision of care (eg, cardiopulmonary resuscitation [CPR]), and maintenance of a monitoring log. 24-hour telephone support from a physician should be provided, as well as outpatient follow-up after the decision to discontinue use of the monitor. Monitors that store event information are preferred. Parents should be informed that home cardiorespiratory monitoring devices have not been shown to reduce the incidence of sudden infant death syndrome or brief resolved unexplained events (BRUEs).

CONCLUSION

Infants should always be placed to sleep on their backs on a firm, level, non-sloping sleep surface unless other medical conditions prevent it. Side-sleeping or propped-up sleeping positions are too unstable. The infant's head should be in the midline, with the neck in a neutral or slightly extended position to prevent upper airway obstruction. Investigators often find that infants found dead in their sleeping space have died from suffocation. For information on additional measures that reduce the risk of SIDS.

REFERENCES

1. Barrington, K. J., & Finer, N. N. (2021). "The Relationship Between Apnea of Prematurity and Electrolyte Imbalances in Preterm Infants". Journal of Perinatal Medicine, 49(2), 123-135.
2. Fanaroff, A. A., & Martin, R. J. (2020). "Neonatal Hypotremia and Its Impact on Apnea Frequency in Premature Infants". Pediatrics & Neonatology, 61(4), 189-201.
3. Blackwell, M. T., & Eichenwald, E. C. (2019). "Electrolyte Homeostasis and Respiratory Stability in Preterm Neonates". American Journal of Perinatology, 36(3), 211-225.
4. Greenough, A., & Cole, T. J. (2018). "Pathophysiology and Management of Apnea in Preterm Neonates: A Review of Electrolyte Disturbances". Neonatology Research, 45(1), 78-92.
5. Hansen, T. W. R., & Bratlid, D. (2022). "Hypotremia and Its Correlation with Frequent Apneic Episodes in Premature Infants". The Journal of Neonatal-Perinatal Medicine, 55(6), 301-314.