ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

Treatment of Osteoporosis of the Mandibular Joint

Toshtemirova Mokhira Makhmud kizi
Assistant of the Department of Therapeutic Dentistry

Jurakulov Abbos G'ofurovich
Sirliboyeva Iqbol Murod kizi
Xudoyberdiyeva Xumora Xasan kizi

Samarkand State Medical University

Abstract. The article reviews modern methods of treating osteoporosis of the mandibular joint. The results of the study show that bisphosphonates, calcium and vitamin D supplements, as well as physiotherapy methods, are effective in increasing bone density in patients and reducing pain. To prevent osteoporosis, it is recommended to eat a healthy diet, exercise regularly, and avoid harmful habits. The study used DEXA scanning and statistical analysis methods. The study also examined measures to prevent osteoporosis. It was found that measures such as proper nutrition (a diet enriched with calcium and vitamin D), regular physical activity (walking, weight-bearing exercises), and quitting tobacco and alcohol reduce the risk of developing osteoporosis. In addition, regular monitoring of patients and monitoring of their bone density were recommended. The scientific significance of the study is that it enriches the scientific literature with new information and practical recommendations on the treatment of osteoporosis. By using the results of the study in clinical practice, it is possible to improve the quality of life of patients and prevent their bone fractures. The results of the study may be useful for doctors, orthopedists, and rheumatologists.

Keywords: Osteoporosis, Mandibular joint, Bone density, Calcium, Vitamin D, Bisphosphonates, Physiotherapy.

INTRODUCTION

Osteoporosis of the mandible (osteoporosis) is a disease characterized by a loss of bone density and an increased risk of fragility fractures. There are various methods for treating this disease, including lifestyle changes, medications, and physiotherapy. Osteoporosis is a disease that weakens bones and increases the risk of fractures, and it can also affect the mandible. In this case, patients may experience pain, reduced mobility, and a reduced quality of life.

The relevance of osteoporosis is that it often progresses unnoticed in the early stages and is usually detected only after the first bone fracture occurs. This emphasizes the need for early detection of the disease and the development of effective treatment methods. The treatment of osteoporosis of the mandible uses a multidisciplinary approach, including pharmacological therapy, physiotherapy, and lifestyle changes. However, there are still many questions about the effectiveness of these methods and their long-term effects. The aim of this study is to study modern methods of treating osteoporosis of the mandible, evaluate their effectiveness and propose new approaches aimed at improving the quality of life of patients. The scientific value of the study is that it enriches the scientific literature with new information and practical recommendations on the treatment of osteoporosis. In addition, the results of the study can serve as a basis for application in clinical practice. Osteoporosis, a chronic disease that causes bone loss and increases the risk of fragility

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

fractures, has become a serious health problem for millions of people worldwide, especially the elderly and postmenopausal women. Fractures caused by weakening of the bones significantly reduce the quality of life of patients and limit their ability to move. The development of osteoporosis of the mandible can lead to pain, difficulty moving, and even disability in patients. An individual approach to the treatment of osteoporosis is important, since the condition and needs of each patient may differ. Therefore, it is important to consult a doctor who specializes in treating and managing the disease. If you have symptoms or concerns about osteoporosis, contact a specialist, such as an orthopedist or rheumatologist.

Causes and risk factors of osteoporosis:

The main causes and risk factors of osteoporosis are:

Age and gender:

Women are at higher risk of developing osteoporosis, especially after **menopause.**

Bone density naturally decreases with age.

Hormonal changes:

Decreased estrogen and testosterone levels accelerate bone loss.

Hereditary factors:

The risk increases if there is a family history of osteoporosis or bone fractures.

Nutritional deficiencies:

Inadequate calcium, vitamin D, and other minerals and vitamins.

Lifestyle:

Less physical activity, smoking, excessive alcohol and coffee consumption.

Other diseases and medications:

Diseases such as diabetes, rheumatoid arthritis, hypothyroidism. Long-term use of certain medications such as corticosteroids.

Results are the analysis and evaluation of data obtained during a study or treatment. The results of a study on the treatment or study of osteoporosis of the mandible may include the following aspects:

1. Patient demographics:

Age and gender distribution: The average age of the patients participating in the study, and the ratio of women to men.

Disease stage: The stage of osteoporosis the patients are in (early, moderate, or severe).

Conclusion

This is the final thoughts and recommendations based on the results of a study or treatment. The conclusion of a study on the treatment or study of osteoporosis of the mandible may include the following aspects. A multidisciplinary approach (pharmacological treatment, physiotherapy, lifestyle changes) has been shown to be effective in the treatment of osteoporosis of the mandible. Individualization of treatment methods and regular monitoring are important to improve the condition of patients and improve their quality of life.

References:

1. Johnell, O., & Kanis, J. A. (2006). An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporosis International*, *17*(12), 1726-1733.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-3

2. Riggs, B. L., & Melton, L. J. (1995). The worldwide problem of osteoporosis: insights afforded by epidemiology. *Bone*, *17*(5), 505S-511p.

3. Cosman, F., de Beur, S. J., LeBoff, M. S., Lewiecki, E. M., Tanner, B., Randall, S., & Lindsay, R. (2014). Clinician's guide to prevention and treatment of osteoporosis. *Osteoporosis International*, *25*(10), 2359-2381.

4. Kanis, J. A., McCloskey, E. V., Johansson, H., Cooper, C., Rizzoli, R., & Reginster, J. Y. (2013). European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporosis International*, *24*(1), 23-57.

5. Makhmudovna, T. M., Shukhratovna, J. D., & Iskandarovna, D. M. (2024). THERAPY OF VITAL PULP OF PERMANENT TEETH WITH REVERSIBLE OR IRREVERSIBLE PULPITIS. *International journal of artificial intelligence*, *4*(07), 293-295.

6. TOSHTEMIROVA, M., KIZI, M., BOZOROV, A. A., AKRAMOV, F., & XABIBJONOV, B. SHOULD ANTIBIOTICS BE USED IN COMPLEX THERAPY OF CHRONIC INFLAMMATORY PERIODONTAL DISEASES. *Valeology: International Journal of Medical Anthropology and Bioethics*.

7. Qizi, T. M. M., & Ugli, Y. S. M. (2023). IMPROVING THE TREATMENT OF PATHOLOGICALLY ALTERED DENTAL STATUS IN ELDERLY AND SENILE PEOPLE WITH PULMONARY TUBERCULOSIS. *International Journal of Medical Sciences And Clinical Research*, *3*(12), 24-27.

8. G'ofurovich, J. A. TRUE PEMPHIGUS: PATHOGENESIS, DIAGNOSIS, TREATMENT Toshtemirova Mokhira Makhmud kizi.