ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

THE SPREAD OF DIARRHEAL DISEASES IN THE KHOREZM REGION

OTAJANOV SH.Z., ARTIQOV I.A., ABDULLAYEVA D.K.

Urgench branch of Tashkent Medical Academy

✓ **Resume**

The study of the etiological structure of intestinal diseases in the Kharezm region was based on a retrospective of the analyzed facts for more than 12 years, from 2012 to 2023. The study of the etiological structure revealed that Sh.Flexneri was identified in 80% of cases from the serovars of E.coli types 026,055 and 0111 were identified. S. Typhimurium was found in Salmonelle bacille in the most common enterobacteria.

Key words: intestines, antibiotic, diarrhea, ecology, shigella, salmonella, enterobacteria.

Introduction.

Diarrhea is one of the most common bowel dysfunctions. According to WHO, diarrheal diseases cause more than 2 million deaths per year. Therefore, the issues of pathogenesis, diagnosis and treatment of diseases occurring with this intestinal dysfunction do not lose their relevance. Any diarrhea is a clinical manifestation of the predominance of water and electrolyte secretion over absorption in the intestine. The ability of the small and large intestines to absorb water and electrolytes is surprising in its complexity and precision in achieving the final result. A person consumes about 2 liters of water daily with food. The volume of endogenous fluid entering the intestinal cavity as part of digestive secretions reaches an average of 7 liters. From 70 to 80% of it is absorbed in the small intestine. From 1 to 2 liters of water enters the colon during the day, 90% of it is absorbed and only 100-200 ml is lost with feces. Even a small change in the volume of fluid in the stool leads to constipation or diarrhea. The creation of new laboratory methods and their implementation in practice gives us a chance to discover the etiological factors of infectious diseases. But humanity is still far from completely defeating infectious diseases. In the Republic of Uzbekistan, in a region with a hot climate, there is a decrease in diarrheal diseases, but the incidence among children remains high. Khorezm region is located in the northern part of the Republic of Uzbekistan, in the last 50 years in the Khorezm region, due to the drying of the Aral Sea, an increase in the concentration of heavy metal salts and various organic compounds in the composition of flowing waters and in the ground has been observed in the ecosystem. This in turn increases the morbidity of the population of the Khorezm region. The main task of infectology at the present time is to study acute diarrheal diseases and their complications, causes of occurrence and mortality. It is known that infectious diseases are formed due to the virulence of the pathogen, toxicity, immunogenicity, resistance of microorganisms, direct action of the external environment and resistance to antibiotics. In particular, the causative agents of diarrheal diseases are being studied, the definition of new antibiotics that are polyresistant to salmonella and shigella. It is difficult to study these diseases from all sides in the Khorezm region, the determination of the properties of the pathogen is determined by complications of the consequences of diseases and the mortality rate is not fully solved.

The purpose of the study: To study the features of the etiological structure of acute diarrheal diseases in the Kharezm region.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

Materials and methods. To study the etiological structure of acute diarrheal diseases in the Kharezm region, we retrospectively analyzed the material of the bacteriological service of the Kharezm region for 12 years (2012-2023). The material is taken from the report of the bacteriological laboratories of the region, the journal of registration of grown and isolated cultures. Statistical processing of the material was carried out by Student and Fisher.

Results and discussions. During the analyzed period, 3755 shigella were isolated, of which 116 (4.4%) strains were identified as Sh.Dysenteriae, 3040 (80%)-Sh.Boydi. At the same time, it was noted that in recent years the specific gravity of Sh.Flexneri has significantly increased, compared with other shigella species. So if in 2023 Sh.Flexneri accounted for 72% of all shigella, then in 2023 this figure reached 94%. The isolation of Sh.Dysenteriae and Sh.Boydii began to decrease from 2015 and by 2012-2023 they were isolated in isolated cases. A similar pattern was observed with respect to Sh.Sonni, so if in 2012-2023 Sh.Sonni among all shigella accounted for 23%, then in 2019 this indicator decreased to 2.1%. The total seeding rate of shigella compared to 2012-2014, in 2016 decreased by 4.4 and 5.9 times. The study of the antigenic structure of shigella showed that in each species there were dominant antigenic variants. Thus, among Sh.Dysenteria, almost half (53 out of 116) belonged to Serovar 1, as well as among Sh.Flexneri more than serovar 1. In Sh.Flexneri, serovar 2a (27.9%) and serovar 6 (27.6%) were detected in more than half of the cases. Among other serovars stand out for (8.6%) t 2b (6.7%). The analysis of the frequency of detection of different serovars and subserovars of shigella in different years did not reveal a pronounced trend towards an increase in different serovars in dynamics, although fluctuations in their specific gravity in different years were quite high. Thus, the detection rate of Sh.Flexneri subserovar 2a ranged from 18.2 to 34.9% of serovar 6 from 20.3% to 40%, subserovar 3a from 3.2% to 16.6, subserovar 2b from 1.9% to 12.5% and so on.

The results of serotyping of diareogenic E.Coli showed that enteropathogenic E. Coli serovariants prevailed in our region, accounting for 90.6% of strains; 797 strains (9.4%) belonged to serogroup 0124 (enteroinvasive E.Coli), enterotaxigenic and enteroimorogic E. Coli were not detected by us. Serovars 026,055,0111 (enteropathogenic E. Coli) sharply prevailed over other serovars, being found in 58.8% of the samples studied. Most often, the causative agents of acute diarrheal diseases were serogroups 055 (24.9%), 0111 the specific gravity of the serogroup 0126 (6%), 020 (5,3%), 044 (4,8%) and 0151 (4.8%). The leading serovar of salmonella causing acute diarrheal diseases S. Typhimurium dominated almost all the years of follow-up and in total salmonella amounted to 91.7%. In the first year of observation (2010-2015), S. Heidelberg was also detected (3.4%), in later periods S. Enteritidis (1.7%); other serovars were sown in isolated cases and in the total amount for 12 years did not exceed 1%. At the same time, attention is drawn to the fact that in some years some of the rare serovars dominated. So, in 2016, 10 strains of S. Derby (2.5%) were isolated, in 2017 12 strains of S.Stanley (3.8%).

Considering that conditionally pathogenic bacteria (UPB) have the greatest etiological significance in acute diarrheal diseases in children, we analyzed their dynamics. A total of 14927 strains of UPB were isolated during the observation period. Of these, representatives of the Enterobacteria family were 13534 (49%); St.aureus 1316 (8.8%); a group of non-fermenting gram-negative bacteria 77 (0.5%). When considering the generic structure of the identified enterobacteria causative agents of acute diarrheal diseases, it was found that the highest percentage of excretion was in proteus (34% of all UPB). The detectability of enterobacteria (20%) and citrobacteria (18%) was almost at the same level, Klebsiella was sown in 14% of cases. The remaining representatives of UPB

were detected less frequently from 0.1% to 2.3%. Analysis of these data in the dynamics of observation showed that there were no significant differences in the frequency of detection of various representatives of this group of bacteria.

Conclusions:

A retrospective analysis of the bacteriological study of pathogens showed that the most significant place in their etiology is occupied by UPB, mainly enterobacteria. Enteropathogenic, diarrheogenic E.Coli prevailed among pathogenic intestinal bacteria, in particular serovars 026, 055, 0111. The specific gravity of shigella was at the lowest level and tends to constantly decrease.

The establishment of the etiological factor of acute diarrheal diseases in patients living in the Kharezm region shows the following: Sh.Flexneri was sown in 80% of cases, while in 27.9% of cases sub-serovars were caused, and in 27.6% serovar 6. Of serovars E.Coli were sown 026 (11,3%), 055 (24,9%) and 0111 (22.6%) and as conditionally pathogenic enterobacteria Proteus (34%).

List of literature.

1. Masharipov, S., Sadullaev, S. E., & Sh, M. D. (2023). THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS. Scientific Impulse, 2(15), 65-70.
2. Masharipova Sh.S, Masharipov S, Sadullaev S.E, & Matyakubova D.Sh. (2023). THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS. Scientific Impulse, 2(15), 65–70. Retrieved from
3. Sadullayev S. E. et al. THE COURSE OF NOSOCOMIAL PNEUMONIA IN PATIENTS ON LONG-TERM ARTIFICIAL LUNG VENTILATION //O'ZBEKISTONDA FANLARARO INNOVATSIYALAR VA ILMIY TADQIQOTLAR JURNALI. – 2024. – Т. 2. – №. 26. – С. 80-84.
4. Машарипова Шохиста Сабировна, Артиков Икром Ахмеджанович, Матякубова Ойша Уриновна РАСТРОЙСТВА ПСИХИКИ У БОЛЬНЫХ ДЕСТРУКТИВНЫМИ ФОРМАМИ ТУБЕРКУЛЕЗА В УСЛОВИЯХ ПАНДЕМИИ COVID-19 // JCRR. 2021. №3.
URL: https://cyberleninka.ru/article/n/rastroystva-psihiki-u-bolnyh-destruktivnymi-formami-tuberkuleza-v-usloviyah-pandemii-covid-19
5. Машарипова Шохиста Сабировна МОРФОЛОГИЧЕСКОЕ СТРОЕНИЕ ЛЕГОЧНЫХ АРТЕРИЙ ПОД ВЛИЯНИЕМ САХАРНОГО ДИАБЕТА // JCRR. 2022. №1.
URL: https://cyberleninka.ru/article/n/morfologicheskoe-stroenie-legochnyh-arteriy-pod-vliyaniem-saharnogo-diabeta
6. Туйчиев Л. Н. и др. A study of the factors affecting the effectiveness of COVID-19 rehabilitation. – 2023.
7. Машарипова Ш. С. О ‘PKA ARTERIYALARINING QANDLI DIABET TA’SIRIDA MORFOLOGIK TUZILISHI //Журнал кардиореспираторных исследований. – 2022. – Т. 3. – №. 1.
8. Yusupov Sh.Sh, Ibrakhimova H.R, & Masharipova Sh.S. (2023). IMMUNOLOGICAL CHARACTERISTICS OF PATIENTS WHOSE BODY IS INFECTED WITH CATTLE SOLITAIRE. Новости образования: исследование в XXI веке, 2(15), 120–124. извлечено от http://nauchniyimpuls.ru/index.php/noiv/article/view/12430
9. Ibrakhimova H.R, Masharipova Sh.S, & Artikov I.A. (2023). CHANGES IN THE IMMUNE STATUS OF PATIENTS WITH PARASITIC DISEASE. Новости образования: исследование в

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

XXI веке, 2(15), 103–108. извлечено от http://nauchniyimpuls.ru/index.php/noiv/article/view/12427

10. Masharipova Sh.S, Masharipov S, & Matyakubova D.Sh. (2023). UDC:616.36-005:75-642 TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. Scientific Impulse, 2(14), 95–99. Retrieved from http://nauchniyimpuls.ru/index.php/ni/article/view/12215

11. Masharipova Sh.S, Ibrakhimova H.R, & Nurllayev R.R. (2023). A METHOD FOR OBTAINING PRECIPITATING SERUMS FOR THE DETECTION OF HUMAN SEMINAL FLUID USED IN THE STUDY OF PHYSICAL EVIDENCE IN FORENSIC BIOLOGICAL LABORATORIES. World Bulletin of Management and Law, 19, 42-44. Retrieved from https://scholarexpress.net/index.php/wbml/article/view/2119

12. IMMUNE STATUS IN PATIENTS WITH PARASITIC DISEASES IN KHOREZM REGION. (2025). Multidisciplinary Journal of Science and Technology, 5(1), 514-517. http://www.mjstjournal.com/index.php/mjst/article/view/2518

13. Машарипов С.М., Юсупов Ш.Р., Машарипова Ш.С., Матякубова О.У. КЛИНИЧЕСКОЕ ТЕЧЕНИЕ ТУБЕРКУЛЕЗА У БОЛЬНЫХ ГЕПАТИТОМ В / Вестник ТМА.uz. 2023 г. № 3 (2) – стр. 155-157.

14. Аскарова Р.И., Юсупов Ш.Р., Машарипова Ш.С., Машарипова Х.К. Эпидемиология легочного туберкулеза/EUROPEAN RESEARCH: INNOVATION IN SCIENCE, EDUCATION AND TECHNOLOGY LVII International correspondence scientific and practical conference. 2019/Издательство: PROBLEMS OF SCIENCE / United Kingdom, 06–07 ноября 2019 год, стр. 96-100

15. S., M. S., E., S. S. ., J., I. S. ., I., Q. N. ., & B., B. Y. . (2024). THE SPREAD OF CHICKENPOX AMONG THE POPULATION. Research Journal of Trauma and Disability Studies, 3(3), 48–54. Retrieved from http://journals.academiczone.net/index.php/rjtds/article/view/2302

16. Masharipova Sh.S, Ibadullayeva S.S, Yakubov K.Y, & Artikov I.A. (2024). FEATURES OF INFECTIOUS MONONUCLEOSIS IN CHILDREN. Scientific Impulse, 2(16), 1165–1171. Retrieved from http://nauchniyimpuls.ru/index.php/ni/article/view/14208

17. Masharipova Sh.S, Ibadullayeva S.S, Yakubov K.Y and Artikov I.A 2024. FEATURES OF INFECTIOUS MONONUCLEOSIS IN CHILDREN. Scientific Impulse. 2, 16 (Jan. 2024), 1165–1171.

18. Садуллаев, С. Е., Машарипова, Ш. С., & Машарипов, С. (2023). КЛИНИКО-ЛАБОРАТОРНЫЕ ОСОБЕННОСТИ ТЕЧЕНИЯ ПНЕВМОНИИ, АССОЦИИРОВАННОЙ С COVID-19, У ДЕТЕЙ РАННЕГО ВОЗРАСТА. Международный журнал образования, социальных и гуманитарных наук. Finland Academic Research Science Publishers, 11(9), 851-856.

19. Сабировна, Ш., & Машарипова, А. И. А. Садуллаев Сирож Эрназарович, и Абдуллаева Дилфуза Кадамовна. 2022.«ТЕЧЕНИЕ КОРОНАВИРУСНОЙ ИНФЕКЦИИ НА ФОНЕ ГЕПАТИТОВ». Новости образования: исследование в XXI веке 1 (5): 573-77. ТЕЧЕНИЕ КОРОНАВИРУСНОЙ ИНФЕКЦИИ НА ФОНЕ ГЕПАТИТОВ». Новости образования.

20. Masharipova Sh.S, Masharipov S, & Matyakubova D.Sh. (2023). UDC:616.36-005:75-642 TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. Scientific Impulse, 2(14), 95–99. Retrieved from https://nauchniyimpuls.ru/index.php/ni/article/view/12215

21. Masharipova Sh.S, Masharipov S, & Matyakubova D.Sh. (2023). UDC:616.36-005:75-642 TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. Scientific Impulse, 2(14), 95–99. Retrieved from http://nauchniyimpuls.ru/index.php/ni/article/view/12215

22.Masharipova Shokhista Sabirovna, & Masharipov Sobir. (2023). UDC: 619:616.995.132.6 IMMUNE STATUS OF ADULTS AND CHILDREN WITH AN ALLERGIC BACKGROUND DIAGNOSED WITH ENTEROBIOSIS. Новости образования: исследование в XXI веке, 2(14), 24–28. извлечено от https://nauchniyimpuls.ru/index.php/noiv/article/view/11911

23.CHARACTERISTICS OF PATHOMORPHOLOGICAL CHANGES IN LYMPHOCYTIC LEUKOSIS IN CHILDREN. (2023). Western European Journal of Medicine and Medical Science, 1(4), 21-26. https://westerneuropeanstudies.com/index.php/3/article/view/122

24.RESULTS OF STUDIES ON THE LEVEL OF POPULATION KNOWLEDGE ABOUT PARASITIC DISEASES AND ITS PREVENTION. (2023). Western European Journal of Medicine and Medical Science, 1(4), 15-20. https://westerneuropeanstudies.com/index.php/3/article/view/121

25. THE STRUCTURE OF COMORBID PATHOLOGY IN CHILDREN WITH COVID-19. (2024). CONFERENCE ON THE ROLE AND IMPORTANCE OF SCIENCE IN THE MODERN WORLD,1(2), 27-28.

26. Sadullaev S. E., Sh M. D. THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS Masharipova Sh. S Masharipov S //Научный импульс. – 2023. – С. 78.

27. Ibrakhimova, H. R., Sadullaev, S. E., & Nurllayev, R. R. (2023). SPREAD OF MYOCARDIAL INFARCTION AMONG THE POPULATION OF THE KHOREZM REGION. IMRAS, 6(7), 328-332.

28. Ibraximova H. R., Sadullayev S. E. AHOLI ORASIDA O ‘TKIR ICHAK KASALLIKLARINING TARQALISHI //Новости образования: исследование в XXI веке. – 2023. – Т. 2. – №. 15. – С. 115-119.

29. Ибрахимова ХР, Матьякубова ОУ, Садуллаев СЭ и Абдуллаева ДК (2023). ГЕЛЬМИНТЫ У ДЕТЕЙ СРЕДИ НАСЕЛЕНИЯ УЗБЕКИСТАНА. IMRAS, 6(7), 323–327. Получено с https://journal.imras.org/index.php/sps/article/view/521

30. Artikov, I. A., Sadullaev, S. E., Ibrakhimova, H. R., & Abdullayeva, D. K. (2023). *RELEVANCE OF VIRAL HEPATITIS EPIDEMIOLOGY. IMRAS, 6 (7), 316–322*.

31. Tuychiev, L. N., Khudaykulova, G. K., Eraliev, U. E., Djuraeva, N. K., & Sadullaev, S. E. (2023). A STUDY OF THE FACTORS AFFECTING THE EFFECTIVENESS OF COVID-19 REHABILITATION.

32. Sadullaev S. E., Sh M. D. THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS Masharipova Sh. S Masharipov S //Научный импульс. – 2023. – С. 78.

33. Туйчиев, Л. Н., Худайкулова, Г. К., Джураева, Н., & Эралиев, У. Э. (2023). A study of the factors affecting the effectiveness of COVID-19 rehabilitation.

34. Sadullaev M., Sh S. M. D. THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS. – 2023.

35. Nurullayev R. R., Sadullayev S. E. DIAREYALI KASALLIKLARNING EPIDYEMIOLOGIK XUSUSIYATLARI //World of Science. – 2023. – Т. 6. – №. 9. – С. 64-67.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

36. Ералиев, У. Э., & Садуллаев, С. Э. (2020). Molecular genetic characteristics of rotavirus infection in children. Молодой ученый, (38), 46-48.

37. ANALYSIS OF THE EPIDEMIOLOGICAL FEATURES OF DIARRHEAL DISEASES IN CHILDREN IN THE SOUTHERN ARAL REGION. (2024). *Multidisciplinary Journal of Science and Technology*, *4*(2), 345-351. http://www.mjstjournal.com/index.php/mjst/article/view/870

38. Рахматуллаева, Ш.Б. Особенности течения COVID-19 у детей с коморбидной патологией // Ш.Б. Рахматуллаева, М.Т. Муминова, А.Х. Нурматов и др. // Педиатрия. Восточная Европа. - 2023. № 3 12. - 436-442. - https://recipe.by/wp-content/uploads/2024/10/436-442_ped_3-2024-v12.pdf (дата обращения: 26.03.2025). doi: 10.34883/PI.2024.12.3.006

39. INTESTINAL IMMUNITY. (2025). *Multidisciplinary Journal of Science and Technology*, *5*(2), 485-488. https://mjstjournal.com/index.php/mjst/article/view/2691

40. PSYCHOLOGICAL REHABILITATION DURING THE CORONAVIRUS PANDEMIC. (2025). *Multidisciplinary Journal of Science and Technology*, *5*(2), 429-433. http://www.mjstjournal.com/index.php/mjst/article/view/2678

41. THE STRUCTURE OF COMORBID PATHOLOGY IN CHILDREN WITH COVID-19. (2024). *CONFERENCE ON THE ROLE AND IMPORTANCE OF SCIENCE IN THE MODERN WORLD*, *1*(2), 27-28. https://universalconference.us/universalconference/index.php/crismw/article/view/641