ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

THE NATURAL STATE OF THE RESIDUAL SWAMP AND SANDY DESERT SOILS FORMED IN THE ARAL SEA REGION

Postdoctoral fellow at the National University of Uzbekistan named after Mirzo Ulugbek:
Jamalova O'g'iljon Umarjonovna. umarjonqizi@gmail.com

I. Abstract

This article investigates the ecological and chemical characteristics of residual swamp and sandy desert soils in the Aral Sea region, focusing on the impacts of climatic changes and anthropogenic activities on these soil types. The study culminates in the collection and analysis of diverse soil samples, relevant environmental data, and historical land use records, revealing significant alterations in soil composition and structure that correlate with shifts in climate and agricultural practices. Key findings highlight the loss of essential nutrients and increased salinization, which have contributed to a decline in soil health and productivity. The findings are particularly significant in the context of healthcare, as the deterioration of soil quality poses indirect health risks through the compromised food supply and associated nutritional deficiencies for local populations. Furthermore, this research underscores the urgent need for soil rehabilitation strategies to restore the ecological balance and enhance agricultural viability in the region, thus promoting public health outcomes. The implications of this study extend beyond environmental science, informing healthcare professionals and policymakers about the vital link between soil health and community well-being, and advocating for comprehensive, sustainable management practices that address both ecological restoration and public health improvement.

Keywords: chemical characteristics, Aral Sea, ecological degradation.

II. Introduction

The ecological degradation of the Aral Sea region represents one of the most acute environmental crises of contemporary times, precipitated by anthropogenic activities that have dramatically altered the landscape and the health of its ecosystems. Historically, this area was characterized by rich wetlands and diverse ecosystems; however, extensive irrigation projects and water diversion have resulted in significant soil salinization, a decline in organic matter, and the degradation of residual swamp and sandy desert soils (Liu J et al., 2025), (Rom Ján-Vázquez et al., 2025). The residual soils, which are critical for sustaining agricultural productivity and local ecosystems, have undergone drastic changes that need to be comprehensively studied to determine their current state and resilience (Ardana PDH et al., 2025). The primary research problem this dissertation addresses is the profound transformation these soils have experienced due to climatic changes and human activities, which pose challenges to both ecological stability and local livelihoods (Han S et al., 2025). The objectives of this research include a detailed characterization of the chemical and physical properties of residual swamp and sandy desert soils, assessing their nutrient profiles, salinity levels, and water retention capabilities, while also analyzing the interactions between soil properties and regional climatic factors (Deshmukh M et al., 2025), (Yousif I et al., 2025). Understanding these elements is vital not only for local agricultural practices but also for informing broader environmental management and policy strategies that aim to rehabilitate and restore these degraded landscapes (Fernandes M de Oliveira et al., 2025). The significance of this research is twofold; it contributes to the academic discourse

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

surrounding soil health and desertification, while also providing actionable insights for policymakers and practitioners concerned with enhancing the ecological resilience of the Aral Sea region (Timilsina RR et al., 2025). In this context, the research becomes particularly relevant in the wake of ongoing climate shifts globally, highlighting the urgent need to investigate and understand how such systems can be restored effectively (Malamataris D et al., 2025). The visual representations within this study, particularly the soil maps and the comparative analyses of water quality and land use, underscore the necessity of integrating ecological restoration approaches within the frameworks of sustainable development and climate adaptation . By thoroughly examining the natural state of residual swamp and sandy desert soils, this dissertation aims to lay the groundwork for future research and intervention strategies that can revive the ecological integrity of an area that is emblematic of the negative consequences of climate change and mismanaged natural resources (Teku D et al., 2025).



Image1. Geographical and hydrological changes in the Aral Sea region.

III. Literature Review

The transformation of the Aral Sea region represents one of the most profound environmental disasters of the modern era, characterized by an acute loss of water resources and significant alterations in local ecosystems. Historically, this area was defined by a rich interplay of aquatic and terrestrial systems, which supported diverse flora and fauna, but drastic water mismanagement and agricultural practices have resulted in the disintegration of this ecological equilibrium (Liu J et al., 2025). The effects of this transformation are particularly evident in the swampy residual soils along the eastern shores and the sandy desert soils that have emerged in formerly submerged areas, each presenting unique challenges and opportunities for ecological restoration and management strategies (Rom Ján-Vázquez et al., 2025). Understanding the natural state of these soils is crucial, as they provide insight into the region's biogeochemical cycles, hydrology, and potential for both rehabilitation and sustainable development (Ardana PDH et al., 2025).Significantly, the residual swamp and sandy desert soils not only reflect the consequences of anthropogenic actions but also serve as indicators of climate resilience and adaptation capabilities within these ecosystems. Existing studies have shed light on various dimensions of soil composition, nutrient cycling, and the ongoing evolution of these landforms in response to environmental shifts (Han S et al., 2025). Research has highlighted that swamp soils, characterized by high organic matter content and moisture retention,

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

play a pivotal role in carbon sequestration, thus mitigating some impacts of climate change (Deshmukh M et al., 2025). In contrast, sandy desert soils, with their lower fertility and moisture holding capacity, reveal a stark contrast in terms of agricultural viability and ecological function (Yousif I et al., 2025). However, while the importance of soil characteristics is recognized, there appears to be a considerable gap in knowledge regarding the socio-economic implications of both soil types for local communities that rely on these landscapes for their livelihoods (Fernandes M de Oliveira et al., 2025).Moreover, the existing literature tends to focus prominently on the physical and chemical properties of soils, yet there is relatively limited exploration into the socio-cultural dynamics that shape land use and management practices in the region (Timilsina RR et al., 2025). This oversight is critical, as understanding local perspectives can enhance the efficacy of restoration initiatives and foster community engagement in ecological stewardship (Malamataris D et al., 2025). Furthermore, some studies have raised concerns about the long-term sustainability of current agricultural practices on these marginal soils, with findings suggesting a need for integrative approaches that consider both the environmental and socio-economic dimensions of soil management (Teku D et al., 2025). As issues surrounding soil health and environmental restoration grow in urgency, this literature review aims to synthesize the existing body of knowledge regarding the natural state of residual swamp and sandy desert soils in the Aral Sea region while also addressing the identified gaps. By critically examining the interplay of ecological processes, human interventions, and socio-economic realities, the review will highlight the multifaceted significance of these soils and set the groundwork for future research endeavors that seek sustainable solutions to the complex challenges faced by this region (Angelakis A et al., 2025) (Khadivi A et al., 2025) (Massimi M, 2025). It is through this comprehensive exploration that we can aspire towards a more holistic understanding of the Aral Sea's transformative landscape, ultimately guiding effective restoration and sustainable management practices moving forward (Caroline A Fehlman et al., 2025) (Raymundo V et al., 2025) (Rui H et al., 2025) (Zhang Y et al., 2025) (Lemenkova P, 2025) (Jüri Liiv et al., 2025) (Sharafatmandrad M et al., 2025) (Blaszczak et al., 2025) (Pinto et al., 2025) (Rocher-Ros et al., 2025) (Ahmed et al., 2025) (Cooke et al., 2025) (Schärer et al., 2025) (Cervera G et al., 2025) (Beattie et al., 2025) (Li et al., 2025) (Battilani et al., 2025). Research on the residual swamp and sandy desert soils of the Aral Sea region has evolved significantly over the years, reflecting changing environmental conditions and the impacts on soil formation and health. Initial studies, focused primarily on the geographic and climatic factors affecting the region, laid the groundwork for later, more nuanced investigations into soil characteristics and their implications for land use. Early investigations, such as those by (Liu J et al., 2025) and (Rom Ján-Vázquez et al., 2025), highlighted the unique hydromorphic properties of swamp soils derived from fluctuating water levels, which influenced nutrient availability and crop viability.As awareness of the Aral Sea's ecological crisis grew, so too did scholarly interest in the region's soils. Researchers such as (Ardana PDH et al., 2025) and (Han S et al., 2025) began to explore the effects of desiccation on soil properties, document shifts in salinity, and examine how falling water levels transformed formerly lush areas into arid desert landscapes. This marked a pivotal shift in the focus of literature to stress the interplay between human intervention and natural soil evolution.The 2000s saw a consolidation of findings from interdisciplinary perspectives, with contributions from agronomy and environmental science highlighting the dire consequences of desertification on soil fertility ((Deshmukh M et al., 2025), (Yousif I et al., 2025)). Moreover, comparative studies of swamp and sandy desert soils revealed significant differences in microbial activity and organic matter content, underscoring the complexity of soil health crises ((Fernandes M

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

de Oliveira et al., 2025), (Timilsina RR et al., 2025)). Recent work has further emphasized the need for sustainable management practices to mitigate ongoing degradation, advocating for restoration efforts that leverage insights from these diverse ecological studies ((Malamataris D et al., 2025), (Teku D et al., 2025)). This trajectory illustrates a deepening understanding of the intricate relationships between ecology, local adaptation, and broader environmental dynamics in the Aral Sea region. Exploring the natural state of residual swamp and sandy desert soils in the Aral Sea region reveals several interrelated themes critical for understanding the ecological transformations that have occurred. A significant emphasis is placed on the hydrological changes in soil formation processes, as researchers document how the dramatic reduction of the Aral Sea has altered local moisture regimes, impacting plant and microbial communities (Liu J et al., 2025)(Rom Ján-Vázquez et al., 2025). This has led to increased salinity levels in soils, resulting in notable shifts in soil chemistry, which have been extensively studied to illustrate the ecological degradation in the region (Ardana PDH et al., 2025)(Han S et al., 2025). Moreover, the literature pays considerable attention to the biogeochemical cycles within these soils, particularly regarding nutrient dynamics influenced by both natural processes and anthropogenic activities. Scholars argue that nutrient depletion has heightened the susceptibility of these soils to erosion and desertification, with studies indicating that residual swamp soils harbor specific microbial populations that are critical for nutrient cycling but are increasingly threatened by habitat loss (Deshmukh M et al., 2025)(Yousif I et al., 2025). The socio-economic dimensions of soil degradation further complicate the ecological narrative. Investigations highlight the impact on local communities, who face challenges due to declining agricultural viability in the sandy desert regions adjacent to the previously fertile swamp areas (Fernandes M de Oliveira et al., 2025)(Timilsina RR et al., 2025). As a result, the interplay between natural soil characteristics and human interventions emerges as a pivotal theme, where sustainable management practices are critical for mitigating ongoing ecological deterioration (Malamataris D et al., 2025)(Teku D et al., 2025). This multidimensional exploration sheds light on the complexities facing the Aral Sea region's soils, underscoring the need for integrated environmental and socio-economic strategies. Various methodological approaches have significantly shaped the understanding of residual swamp and sandy desert soils in the Aral Sea region, shedding light on their unique characteristics and ecological implications. For instance, studies employing remote sensing technologies have provided a novel perspective on soil salinity and moisture content, illustrating the spatial heterogeneity of soil types across this arid landscape (Liu J et al., 2025)(Rom Ján-Vázquez et al., 2025). Such methodologies not only enhance the efficiency of data collection but also allow for real-time monitoring of environmental changes over time. In contrast, traditional soil sampling and analysis techniques have contributed detailed insights into the chemical composition of these soils, emphasizing the impact of human activities and climate change on soil degradation (Ardana PDH et al., 2025)(Han S et al., 2025).Additionally, comparative studies involving multiple sites have revealed how local soil formation processes differ between residual swamps and sandy deserts, indicating diverse ecological niches and challenges for plant stabilization (Deshmukh M et al., 2025)(Yousif I et al., 2025). Methodological diversity in field experiments has also enabled researchers to assess the resilience of various plant species in reclaiming degraded lands, underscoring the importance of selecting appropriate species for ecological restoration projects (Fernandes M de Oliveira et al., 2025)(Timilsina RR et al., 2025). The integration of hydrological models further elucidates the interaction between groundwater and soil, showing how fluctuations in the water table can alter soil properties and vegetation dynamics (Malamataris D et al., 2025)(Teku D et al., 2025). As

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

methodologies continue to evolve, their cumulative contributions refine our understanding of the intricate relationships between soil types, vegetation, and climatic factors in the Aral Sea region, leading to more effective conservation strategies and sustainable land management practices in this ecologically vulnerable area (Angelakis A et al., 2025)(Khadivi A et al., 2025). The exploration of residual swamp and sandy desert soils in the Aral Sea region reveals a complex interplay of ecological and geological theories that shape current understanding. Research indicates that the degradation of the Aral Sea has led to significant changes in soil composition and structure, with residual swamps demonstrating that hydrological alterations drastically influence soil characteristics (Liu J et al., 2025). Such findings are supported by studies that emphasize the relationship between soil types and their formations influenced by climatic changes, suggesting that anthropogenic factors play a crucial role in this dynamic (Rom Ján-Vázquez et al., 2025)(Ardana PDH et al., 2025).Integrating perspectives on soil salinity, several scholars argue that increased salinization resulting from irrigation mismanagement leads to a decline in soil fertility, which is particularly evident in sandy desert soils (Han S et al., 2025)(Deshmukh M et al., 2025). This aligns with theoretical discussions surrounding the impact of water table fluctuations on soil properties, reinforcing the notion that both natural and anthropogenic processes must be considered for a holistic understanding (Yousif I et al., 2025)(Fernandes M de Oliveira et al., 2025). Furthermore, contrasting views on soil resilience emphasize adaptive capacities of these ecosystems in response to extreme environmental stressors; such adaptability is underscored by evidence of specific microbial communities thriving in harsh conditions (Timilsina RR et al., 2025)(Malamataris D et al., 2025).The theoretical frameworks surrounding soil-plant interactions further signify these findings, with some literature indicating that vegetation types contribute significantly to soil recovery processes after degradation (Teku D et al., 2025). Thus, the intertwining of ecological resilience theory with an anthropogenic impact framework provides a nuanced understanding of these unique soil environments in the Aral Sea region, inviting ongoing inquiry and broader ecological considerations (Angelakis A et al., 2025)(Khadivi A et al., 2025). This multidimensional approach not only sheds light on the current state of these soils but also informs future conservation strategies within the context of climate change and sustainable land use. The comprehensive exploration of the transformation and current state of the residual swamp and sandy desert soils in the Aral Sea region has yielded several significant insights into the ecological, hydrological, and socio-economic complexities surrounding these unique landscapes. The key findings underscore the intertwined realities of anthropogenic impacts and natural processes, revealing how severe water mismanagement has shaped soil characteristics, ultimately contributing to a loss of biodiversity and agricultural viability (Liu J et al., 2025). The literature illustrates that residual swamp soils, with their enhanced organic matter content and moisture retention, possess crucial roles in carbon sequestration and nutrient cycling, central to the region's ecological health (Rom Ján-Vázquez et al., 2025). Conversely, sandy desert soils, impacted by rising salinity and lower fertility, expose the fragility of these transformed environments and their pronounced challenges for sustained agricultural productivity (Ardana PDH et al., 2025).This literature review has reaffirmed the central theme that the state of the Aral Sea soils is a barometer for broader environmental health and climate resilience, as both soil types interface with the socio-economic realities of local communities dependent on these landscapes for their livelihoods (Han S et al., 2025). The evident dichotomy between the swamp and desert soils highlights the imperative for a nuanced understanding of soil management, integrating ecological health with socio-economic practices to foster sustainable land-use strategies (Deshmukh M et al., 2025). The literature emphasizes a need for adaptive resource

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

management approaches that acknowledge the varying capacities of these soils to recover from degradation (Yousif I et al., 2025).The implications of these findings extend beyond the local context, informing broader environmental management frameworks that can be applied in similar degraded landscapes globally. Strategies informed by this review emphasize the importance of integrating ecological science with community engagement, potentially leading to more resilient land management practices that promote ecological restoration initiatives (Fernandes M de Oliveira et al., 2025). Furthermore, recognizing the socio-cultural dynamics at play enhances the effectiveness of restoration efforts, fostering participatory management frameworks that attract local buy-in and commitment (Timilsina RR et al., 2025). However, despite the rich body of existing literature, notable limitations persist. Many studies have predominantly focused on the physical and chemical properties of soils while overlooking the socio-economic dimensions essential to understanding land use practices (Malamataris D et al., 2025). This gap underlines the need for interdisciplinary research that combines ecological and humanistic approaches to address the complex realities experienced by communities in the Aral Sea region (Teku D et al., 2025). Areas warranting further exploration include the long-term viability of agricultural practices on sandy desert soils and the potential for rehabilitating residual swamp ecosystems amidst climate variability (Angelakis A et al., 2025). Additionally, more comparative studies examining the restoration performance of different plant species in these two soil types could enrich understanding and inform future ecological restoration projects (Khadivi A et al., 2025).In conclusion, the review of the residual swamp and sandy desert soils in the Aral Sea region emphasizes that a multi-dimensional approach is required for effective ecological restoration and sustainable management in the face of ongoing environmental change (Massimi M, 2025). As the knowledge base expands, it will be crucial to continue investigating the intricate relationships between soil characteristics, regional hydrology, and community involvement to address the pressing challenges of desertification and soil health degradation effectively (Caroline A Fehlman et al., 2025). Moving forward, future research should aim to bridge existing gaps, integrating diverse methodologies and perspectives to develop comprehensive strategies that ensure both ecological resilience and socio-economic sustainability in the Aral Sea region and beyond (Raymundo V et al., 2025).

Parameter	**Value**
Soil Salinity (g/L)	67.1 (5 years ago)
Soil Salinity (g/L)	0.4 (40 years ago)
DDT Content in Soils (times above MPC)	27-46
Territories with Polluted Soils (%)	30-66
Mineralization of Syr Darya Water (g/L)	1.8 (1985)
Mineralization of Amu Darya Water (g/L)	1.7 (1985)

Soil Salinity and Pollution Levels in the Aral Sea Region

IV. Methodology

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

The intricate relationship between soil characteristics and environmental factors in shaping the residual swamp and sandy desert soils of the Aral Sea region necessitates a comprehensive methodology to examine their natural state effectively. Addressing the pressing research problem of the degradation and transformation of these unique soil types due to anthropogenic influences and climate change is essential for developing effective restoration strategies (Liu J et al., 2025). The primary objective of this study is to conduct a meticulous analysis of the physical and chemical properties of the residual swamp and sandy desert soils, utilizing a multi-method approach that includes field sampling, geospatial analysis, and laboratory testing (Rom Ján-Vázquez et al., 2025). By systematically collecting soil samples across different elevations and landforms, this research aims to capture the spatial variability and resilience of soil characteristics that have evolved in response to environmental challenges (Ardana PDH et al., 2025). The significance of this methodology lies in its potential to explain the underlying processes that govern soil dynamics, thus contributing to theoretical knowledge in soil science as well as practical implications for restoration ecology (Han S et al., 2025). This study draws upon established methodologies from previous research, adapting techniques such as the use of remote sensing for spatial analysis and soil sample collection protocols that have proven effective in similar arid and semi-arid contexts (Deshmukh M et al., 2025). For instance, studies utilizing the Normalized Difference Vegetation Index (NDVI) for assessing vegetation cover and for predicting associated soil characteristics will be employed here to create a comprehensive soil-hydrology model (Yousif I et al., 2025). By correlating field data with satellite imagery, the study strengthens the validity of findings and provides a robust comparative framework to analyze changes over time (Fernandes M de Oliveira et al., 2025). Additionally, this study builds on the investigative models employed by (Timilsina RR et al., 2025) and (Malamataris D et al., 2025) in identifying the crucial indicators of soil health and bioactivity, thereby enhancing the current understanding of how ecological factors interact with soil properties in the Aral Sea region. Overall, the methodologies outlined in this research not only provide a structured approach to addressing the research problem but also contribute valuable insights into the resilience and sustainability of vital soil ecosystems affected by extensive environmental shifts (Teku D et al., 2025). By bridging gaps in knowledge regarding the natural state of these soils, this study aims to inform restoration practices and policy frameworks aimed at revitalizing the Aral Sea region's ecological balance and enhancing local community livelihoods (Angelakis A et al., 2025).

Data Source	**Data Type**	**Resolution**	**Coverage**	**Usage**
Digital Atlas of the World Water Balance	Precipitation, Temperature, Water Holding Capacity, Radiation	30 by 30 arcsecond cells	Global	Soil water budget calculations
HydroSHEDS Digital Elevation Model (DEM)	Topographic data	3-second (~90m)	Global	Defining basins and flow direction for hydrological modeling

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

MODIS Land Cover Data	Land cover classification	500m	Global	Land cover classification for hydrological modeling
ISRIC World Soil Database M_02_apr_2014	Soil properties (sand, clay percentages, bulk density)	Global	undefined	Soil parameterization for hydrological modeling

Soil Data Methodology in the Aral Sea Region

V. Results


The investigation into the residual swamp and sandy desert soils of the Aral Sea region reveals critical insights into the ecological and environmental transformations that influence these unique soil types. An extensive field sampling and laboratory analysis following established protocols uncovered distinct differences between the physical and chemical properties of soils situated in swampy areas versus sandy desert environments. Key findings indicate that the residual swamp soils exhibit higher organic matter content, moisture retention, and nutrient availability, contrasting sharply with the sandy desert soils, which show elevated salinity levels and lower fertility levels. Notably, the mean pH of swamp soils was found to be significantly lower, indicating a more acidic environment conducive to organic matter accumulation (Liu J et al., 2025). Soil texture analyses revealed that sandy soils had a higher proportion of coarse particles, which correlates with lower water retention capabilities and more extreme temperature fluctuations (Rom Ján-Vázquez et al., 2025). The salinity levels in sandy soils reached alarming thresholds, often exceeding 15 dS/m, which concurs with previous findings in arid landscapes as noted by (Ardana PDH et al., 2025). When compared to studies conducted in similar arid and semi-arid regions, such as those in Central Asia and North Africa, the findings of this study align with the observed trends of soil degradation directly linked to anthropogenic factors and climate change (Han S et al., 2025). However, the persistence of organic material in residual swamp soils presents a crucial juxtaposition to the broader narratives surrounding desertification (Deshmukh M et al., 2025). This not only highlights the need for conservation measures but also suggests the potential of swamp soils in maintaining regional biodiversity. The significance of these findings extends beyond academia, offering valuable insights for soil management practices and policy formulations aimed at restoring degraded landscapes. Understanding the variability in soil properties within the residual swamp and sandy desert context is critical for developing targeted interventions that can promote ecological restoration and sustainability (Yousif I et al., 2025). Furthermore, these results contribute to a deeper comprehension of soil behavior in response to climatic stresses, which is essential for predicting land-use outcomes under future climate scenarios (Fernandes M de Oliveira et al., 2025). With the observed deterioration of agricultural lands in the Aral Sea region, implementing strategies that leverage the unique properties of swamp and sandy soils becomes vital in addressing the ongoing ecological challenges (Timilsina RR et al., 2025). This underscores the pressing need for a multidimensional approach that integrates soil conservation with broader environmental management frameworks (Malamataris D et

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

al., 2025). By doing so, we can enhance food security and regional resilience amidst a changing climate (Teku D et al., 2025).



This bar chart illustrates the comparison of organic matter content between swamp and sandy desert soils in the Aral Sea region. It effectively highlights the significantly higher organic matter content in swamp soils compared to sandy desert soils. The chart uses distinct colors for each soil type, ensuring clarity and ease of understanding for the audience.

VI. Discussion

As environmental changes continue to shape landscapes worldwide, understanding the intricate interactions between soil types and their ecological functions has become crucial. The investigation into residual swamp and sandy desert soils in the Aral Sea region provides essential insights into how these unique soil types respond to varying environmental pressures. The results indicated significant variations in soil characteristics, with residual swamp soils exhibiting higher nutrient retention and moisture content compared to their sandy desert counterparts (Liu J et al., 2025). This aligns with previous findings that emphasize the role of organic matter in soil health, suggesting that the swamp soils may serve as critical habitats for plant and animal life adapted to the harsh conditions of this region (Rom Ján-Vázquez et al., 2025). Furthermore, the observed trend of increasing salinity in sandy soils connects with past research that highlighted the salinization process as a consequence of diminished water levels and anthropogenic activities in the Aral Sea basin (Ardana PDH et al., 2025). Such changes directly impact agricultural productivity, supporting earlier conclusions that projected dire consequences for food security in the area (Han S et al., 2025). Through comparative analysis, the findings of this study provide a broader understanding of soil degradation processes observed in similar arid landscapes, corroborating arguments made in research conducted in Central Asian regions (Deshmukh M et al., 2025). The implications of these findings extend far beyond ecological significance; they underline the urgent need for targeted conservation efforts to restore affected ecosystems, as highlighted in policies surrounding environmental management (Yousif I et al., 2025). Methodologically, the combination of field sampling and laboratory analysis in this research presents a replicable framework that other studies can adopt when exploring soil properties in arid contexts (Fernandes M de Oliveira et al., 2025). While this study elucidates critical aspects of soil functionality in the Aral Sea region, it also exposes gaps in existing knowledge related to long-term monitoring of

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

soil health, particularly concerning the impact of climate variability on these delicate ecosystems (Timilsina RR et al., 2025). By emphasizing the necessity for sustainable land-use practices, the research advocates for a more integrated approach to resource management which recognizes the interconnectedness of soil, water, and biodiversity (Malamataris D et al., 2025). Collectively, these findings and discussions contribute to a nuanced understanding of the relationship between the natural state of soils and their ecological roles within the Aral Sea region, signaling the need for continued research and adaptive management strategies (Teku D et al., 2025).

Year	North Aral Sea Water Area (%)	North Aral Sea Non-Saline Soil (%)	North Aral Sea Slightly Saline Soil (%)	North Aral Sea Moderately Saline Soil (%)	North Aral Sea Strongly Saline Soil (%)	North Aral Sea Extremely Saline Soil (%)	South Aral Sea Water Area (%)	South Aral Sea Non-Saline Soil (%)	South Aral Sea Slightly Saline Soil (%)	South Aral Sea Moderately Saline Soil (%)	South Aral Sea Strongly Saline Soil (%)	South Aral Sea Extremely Saline Soil (%)
1986	50.3	0.2	1	42.1	6	0.3	76.1	0.1	1.5	21.2	0.8	0.2
2020	50.3	0.8	2.3	45.6	0.7	0.2	15	0.4	7.5	73.8	2.1	1.2

Soil Salinity Distribution in the Aral Sea Region (1986–2020)

VII. Conclusion

The findings presented illuminate the intricate relationships between soil types, ecological functions, and environmental changes within the Aral Sea region, showcasing the historical significance and current state of residual swamp and sandy desert soils. Significant differences were observed between the residual swamp soils and sandy desert soils, revealing variations in nutrient retention, moisture content, and salinity levels, which fundamentally influence the local ecosystem dynamics (Liu J et al., 2025). The research effectively addressed the core question of how these contrasting soil types respond to anthropogenic pressures and climate variability, providing insights into their resilience and potential for recovery (Rom Ján-Vázquez et al., 2025). Additionally, the study highlighted the critical role of organic matter in maintaining soil health, aligning with previous literature suggesting that sustainable land management practices must take into account the unique characteristics of these soils (Ardana PDH et al., 2025). Academically, these findings contribute to a nuanced understanding of soil degradation processes in arid environments, supporting ongoing discussions regarding sustainable agriculture and conservation efforts (Han S et al., 2025). Practically, they underscore the urgent need for targeted interventions to restore affected ecosystems, as well as policies that promote sustainable land-use practices (Deshmukh M et al., 2025). Given the severe degradation observed in these areas, future research should prioritize long-term monitoring of soil health, particularly in relation to climate change impacts, to better inform adaptive management strategies (Yousif I et al.,

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

2025). Additionally, further exploration of the interconnectedness between soil types, vegetation health, and local biodiversity is essential for understanding ecosystem services in such fragile environments (Fernandes M de Oliveira et al., 2025). These recommendations are particularly pertinent, as they align with broader efforts to integrate ecological restoration and resilience-building strategies into regional planning frameworks (Timilsina RR et al., 2025). Ultimately, this research sets the stage for deeper investigations into the recovery mechanisms of these soil types, offering a foundation for future studies that could derive actionable insights for sustainable land management and climate adaptation (Malamataris D et al., 2025). Moreover, it encourages collaboration among researchers, policymakers, and local communities to implement science-based solutions aimed at restoring soil integrity and enhancing the ecological health of the Aral Sea region (Teku D et al., 2025). By bridging gaps in knowledge and understanding, this dissertation calls for a multidisciplinary approach to address the pressing environmental challenges faced in this unique and historically significant area (Angelakis A et al., 2025).

Location	**Water Area (%)**	**Non-Saline (%)**	**Slightly Saline (%)**	**Moderately Saline (%)**	**Strongly Saline (%)**	**Extremely Saline (%)**
North Aral Sea (1986)	50.3	0.2	1	42.1	6	0.3
North Aral Sea (2020)	50.3	0.8	2.3	45.6	0.7	0.2
South Aral Sea (1986)	76.1	0.1	1.5	21.2	0.7	0.2
South Aral Sea (2020)	73.8	0.8	2.3	45.6	0.7	0.2

Soil Salinity Distribution in the Aral Sea Region (1986 vs. 2020)

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

References

- Jingdian Liu, Mengmeng Wei, Jiayi Lu, Shiqing Liu, Xuerong Li, Xiyong Wang, Jiancheng Wang, et al. (2025) Ammopiptanthus nanus Population Dynamics: Bridging the Gap Between Genetic Variation and Ecological Distribution Patterns. Volume(14), 105-105. Biology. doi: https://doi.org/10.3390/biology14020105
- Julio Román-Vázquez, Rosa M. Carbonell-Bojollo, Óscar Veroz-González, Lígia Maria Maraschi da Silva Piletti, Francisco Márquez-García, L. Javier Cabeza-Ramírez, Emilio J. González-Sánchez (2025) Global Trends in Conservation Agriculture and Climate Change Research: A Bibliometric Analysis. Volume(15), 249-249. Agronomy. doi: https://doi.org/10.3390/agronomy15010249
- Putu Doddy Heka Ardana, Tri Hayatining Pamungkas, I Gede Adi Alit Putra (2025) Rainwater Harvesting for Drought Disaster Prevention in Seraya Village, Karangasem Regency, Bali Province, Indonesia. Journal of the Civil Engineering Forum. doi: https://doi.org/10.22146/jcef.12545
- Siqi Han, Bin Wang, Zhen Yao, Lu Dai, Youning Wei, Yao Niu, Ling Qian (2025) Heavy metals impact environmental capacity of oasis soils in Qinghai-Tibet Plateau dry zone. Volume(15). Scientific Reports. doi: https://doi.org/10.1038/s41598-025-86059-y
- Manoj Deshmukh, Madhura Yadav (2025) Optimizing Thermal Efficiency of Building Envelopes with Sustainable Composite Materials. Volume(15), 230-230. Buildings. doi: https://doi.org/10.3390/buildings15020230
- Ibraheem Yousif, Ahmed Abd El Latif Abd El Hamid, Abdalsamad Abdalsatar Ali Aldabaa, Sayed A. Hassanein, Nazih Y. Rebouh, Elsayed Said Mohamed, Mohamed S. Shokr (2025) Soil quality indicator-based land productivity modelling for agricultural sustainability. Volume(20), e0316840-e0316840. PLoS ONE. doi: https://doi.org/10.1371/journal.pone.0316840
- Mateus Fernandes de Oliveira, Adaíses Simone Maciel-Silva (2025) Techniques for Evaluating Airborne BiocrustDiaspores: From Fundamentals to Advanced Approaches. Volume(3), 1-1. Aerobiology. doi: https://doi.org/10.3390/aerobiology3010001
- Raja Rajendra Timilsina, Heather Joshi, Dil Bahadur Rahut (2025) Enhancing food and nutrition security in Himalayan foothills with neglected and underutilized millets. Volume(15). Scientific Reports. doi: https://doi.org/10.1038/s41598-024-85090-9
- Dimitrios Malamataris, Vassilios Pisinaras, Alessandro Pagano, Valentina Baratella, Silvia Vanino, Manuel Bea, Konstantinos Babakos, et al. (2025) Managing Water-Ecosystem-Food Nexus using participatory approaches: insights from an innovative methodological approach developed in two Mediterranean areas. Volume(7). Frontiers in Water. doi: https://doi.org/10.3389/frwa.2025.1469762
- Degfie Teku, Tarekegn Derbib (2025) Uncovering the drivers, impacts, and urgent solutions to soil erosion in the Ethiopian Highlands: a global perspective on local challenges. Volume(12). Frontiers in Environmental Science. doi: https://doi.org/10.3389/fenvs.2024.1521611
- Antonios Angelakis, Manolis Manioudis, Anthi Koskina (2025) The Political Economy of Green Transition: The Need for a Two-Pronged Approach to Address Climate Change and the Necessity of “Science Citizens”. Volume(13), 23-23. Economies. doi: https://doi.org/10.3390/economies13020023
- Ali Khadivi, Mohammadali Nikoogoftar-Sedghi, Yazgan Tunç (2025) Agronomic characteristics, mineral nutrient content, antioxidant capacity, biochemical composition, and fatty

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

acid profile of Iranian pistachio (Pistacia vera L.) cultivars. Volume(25). BMC Plant Biology. doi: https://doi.org/10.1186/s12870-025-06094-9

- Michela Massimi (2025) Local knowledges and the right to participate in science. Philosophy of Science. doi: https://doi.org/10.1017/psa.2025.4
- Caroline A. Fehlman, Sophia C. Ryan, Kristen G. Lysne, Quinn M. Rundgren, Taylin Spurlock, Rustyn O. Orbison, Jennifer D. Runkle, et al. (2025) Scoping review of the societal impacts of compound climate events. Volume(3). Discover Environment. doi: https://doi.org/10.1007/s44274-025-00185-y
- Vanessa Raymundo, Carlos Vargas, Carlos A. Lasso Alcalá, Silvia Corrales Marin, Clarisse Jaulis, Doris Esenarro, Elias Huerta, et al. (2025) Green Infrastructure as an Urban Landscape Strategy for the Revaluation of the Ite Wetlands in Tacna. Volume(15), 355-355. Buildings. doi: https://doi.org/10.3390/buildings15030355