ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

ARLY DIAGNOSIS OF CHRONIC SINUSITIS: MODERN DIAGNOSTIC METHODS AND SCIENTIFIC FOUNDATIONS

Tokhirjon Abdunazarov
Assistant, Department of Surgical Diseases, Central Asian Medical University, Fergana, Uzbekistan
Otaqulov Otabek
Assistant, Department of Surgical Diseases, Central Asian Medical University, Fergana, Uzbekistan
Kamoldinov Bobur
Assistant, Department of Surgical Diseases, Central Asian Medical University, Fergana, Uzbekistan

Keywords: Maxillary sinus, ENT combine, CT, MRI.

Abstract

Chronic sinusitis is a long-lasting inflammatory disease of the maxillary sinus. If not diagnosed in a timely manner, it can negatively affect the patient's quality of life and lead to various complications. This article discusses effective methods for early diagnosis of chronic sinusitis, modern diagnostic technologies, and efficient examination techniques.

1. Introduction

Sinusitis is an inflammation of the paranasal sinuses that can present in acute or chronic forms. Chronic sinusitis lasts for more than 12 weeks, and symptoms include nasal congestion, mucous or purulent discharge, headache, and a sensation of pressure in the face.
If not diagnosed in time, chronic sinusitis can lead to the following severe complications:

- Orbital complications (spread of infection to the eye area).
- Intracranial infections (meningitis, brain abscess).
- Respiratory and cardiovascular system-related problems.

The purpose of this article is to analyze effective methods for early diagnosis of chronic sinusitis and to provide recommendations based on modern research.

2. Pathophysiology and Risk Factors of Chronic Sinusitis

2.1. Pathophysiology

The development of chronic sinusitis is influenced by the following factors:

- Persistent inflammation of the mucous membrane of the nasal cavity.
- Disruption of the sinus's natural drainage (accumulation of mucous or purulent secretions).
- Weakened immune system or allergic inflammation processes.

2.2. Risk Factors

- Allergic rhinitis and asthma.
- Deviated septum.
- Decreased immunity (e.g., diabetes, HIV).
- Harmful habits (smoking, exposure to dust or chemicals).
- Chronic infectious diseases.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

3. Diagnosis

Various clinical, imaging, and laboratory methods are used to diagnose chronic sinusitis.

3.1. Clinical Symptoms and Physical Examination

The analysis of patient complaints and the examination by a doctor are crucial.

- **Headache** – pain in the forehead, around the eyes, and in the cheek area.
- **Nasal congestion and discharge** – mucous or purulent secretion.
- **Reduced sense of smell.**
- **Fever** – rarely observed, only when inflammation intensifies.

The doctor examines the external nose and face area and checks for pain through palpation.

3.2. Endoscopic Diagnosis

Nasal endoscopy allows for the inspection of the inner parts of the nasal cavity using a special device (nasal endoscope).

- Narrowing of the natural openings of the sinuses.
- Hyperemia and swelling of the mucous membrane.
- Presence of purulent discharge.

3.3. Radiological Diagnosis

Computed Tomography (CT)

CT is considered the "gold standard" in diagnosing chronic sinusitis. It reveals:

- The degree of mucous accumulation in the sinus cavities.
- Thickening of the mucous membrane.
- Deviated septum or presence of polyps.

Magnetic Resonance Imaging (MRI)

MRI is recommended if complications (orbital or intracranial infections) are suspected.

Radiographic Examination

This method provides less detailed information but is still used in some cases.

3.4. Laboratory Diagnosis

- **Bacteriological examination of nasal secretions** – to detect bacterial infections.
- **C-reactive protein (CRP) and general blood test** – to assess inflammation.
- **Allergic tests** – if allergic factors are suspected.

---

4. Differential Diagnosis of Chronic Sinusitis

Chronic sinusitis must be differentiated from the following conditions:

- Allergic rhinitis.
- Nasal polyposis.
- Nasal internal tumors and growths.
- Neuralgia (trigeminal nerve neuralgia).

Thus, it is not appropriate to base a diagnosis solely on symptoms, and a combination of diagnostic tests must be used.

---

5. Treatment and Prevention

Both conservative and surgical methods are used in the treatment of chronic sinusitis.

5.1. Conservative Treatment

- **Antibiotic therapy** – if a bacterial infection is detected.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

- **Nasal corticosteroids** – to reduce allergic or inflammatory processes.
- **Nasal irrigation** (with saline or seawater solutions).
- **Mucolytics and antihistamines.**

5.2. Surgical Treatment

If conservative treatment is ineffective or if sinus drainage is poor, the following surgical methods are employed:

- **Endoscopic sinus surgery (FESS).**
- **Maxillectomy** – drainage of the sinus cavity.

5.3. Prevention

- Preventing allergic reactions.
- Keeping the nasal cavity clean regularly.
- Strengthening immunity.
- Protecting the nose in cold weather.

---

6. Conclusion

Early diagnosis of chronic sinusitis is crucial for effective treatment and prevention of complications. A combination of clinical symptoms, endoscopic examinations, CT, and laboratory tests provides the optimal diagnostic approach.

References:

1. Brook, I. (2016). Chronic sinusitis in children: Pathogenesis and management. Pediatric Drugs, 18(4), 275–281.

2. Fokkens, W. J., Lund, V. J., Hopkins, C., et al. (2020). EPOS 2020: European Position Paper on Rhinosinusitis and Nasal Polyps 2020. Rhinology Supplement, (29), 1–464.

3. Rosenfeld, R. M., Piccirillo, J. F., Chandrasekhar, S. S., et al. (2015). Clinical practice guideline (update): Adult sinusitis. Otolaryngology–Head and Neck Surgery, 152(2_suppl), S1–S39.

4. Klossek, J. M., et al. (2005). Epidemiology of chronic rhinosinusitis in France. Rev Laryngol Otol Rhinol (Bord), 126(1), 27–31.

5. Kennedy, D. W. (1992). Prognostic factors, outcomes and staging in ethmoid sinus surgery. Laryngoscope, 102(12 Pt 2 Suppl 57), 1–18.