ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

Using methods in the process of determining the composition of the main pests of the poplar tree

Rahimboyeva Maftuna Maqsud qizi
Faculty of Natural and Agricultural Sciences, Urgench State University named after Abu Rayhon Beruni, Group 211, student of the Plant Protection field

Abstract: The poplar tree (Populus spp.) is an important tree species widely cultivated for timber, paper production, and environmental benefits. However, it is highly susceptible to various pests that significantly affect its growth and health. This article explores the methods used to identify and analyze the composition of the main pests that damage poplar trees. Through field surveys, entomological sampling, and laboratory analysis, different pest species are categorized, and their impact on the tree's physiology is assessed. The study also highlights the effectiveness of various pest control methods and provides recommendations for sustainable management practices. This research contributes to a deeper understanding of pest dynamics in poplar ecosystems and aims to improve the long-term health of poplar plantations.

Keywords: Poplar tree, pests, pest composition, entomological sampling, pest control, sustainable management, tree health, ecosystem dynamics, plant protection.

Introduction: Poplar trees (Populus spp.) are among the most widely planted species due to their rapid growth, adaptability to various environmental conditions, and their economic significance in the timber, paper, and bioenergy industries. They are also used for soil erosion control, windbreaks, and reclamation of degraded lands. Despite these benefits, poplars are susceptible to a wide range of pests that threaten their health and productivity. Pests can cause significant damage to poplar trees by feeding on leaves, bark, and roots, disrupting their physiological processes and reducing timber quality.

The complexity of pest interactions with poplar trees requires a thorough understanding of their composition and impact. Different pests, ranging from insects and fungi to bacteria, affect various parts of the tree, leading to defoliation, bark damage, and even death in extreme cases. Identifying these pests, understanding their life cycles, and determining their distribution patterns are essential steps toward effective pest management. Without timely and accurate pest control measures, poplar plantations may suffer substantial economic losses.

Recent advances in pest monitoring and detection methods have made it possible to assess pest populations with greater accuracy and efficiency. Techniques such as entomological surveys, pest trapping, and molecular analysis are increasingly used to identify and monitor pest species in different regions. However, despite these advances, comprehensive studies on the composition of pests affecting poplar trees are still limited, particularly in certain regions where poplars are of great ecological and economic value.

This article aims to explore the primary pests that damage poplar trees, focusing on the methods used to determine their composition. By identifying the most prevalent pest species and understanding their interactions with poplar trees, the study provides valuable insights for developing sustainable pest management strategies. The ultimate goal is to contribute to the protection and long-term health of poplar plantations, ensuring that they continue to provide economic and environmental benefits for generations to come.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

Relevance of Work: The relevance of this research lies in the increasing importance of poplar trees in various industries and their vulnerability to pest infestations. Poplar trees are crucial for timber production, paper manufacturing, and environmental conservation. In addition, they play a key role in carbon sequestration and land reclamation. However, poplar plantations are frequently targeted by a wide range of pests that can lead to reduced growth rates, diminished wood quality, and, in some cases, total tree mortality. This makes the health and sustainability of poplar plantations an urgent concern for both the forestry industry and environmental management.

In recent years, climate change has further exacerbated pest outbreaks by altering pest life cycles, expanding their geographical range, and increasing the severity of infestations. The ability to accurately identify and understand the composition of pest species is essential for mitigating the damage caused by these pests. This is particularly relevant as poplar plantations become more widespread in regions where pest pressures may be intensifying.

Current pest management strategies often rely on traditional approaches, which can be inefficient or harmful to the ecosystem. The development of more precise, sustainable, and environmentally friendly pest control methods requires an in-depth understanding of pest species composition and behavior. By focusing on the identification and monitoring of the main pests affecting poplar trees, this research aims to provide new insights into effective pest management strategies, particularly in the context of integrated pest management (IPM).

Moreover, as global demand for poplar-based products continues to grow, ensuring the long-term health of poplar trees is vital for maintaining the economic viability of these plantations. This work is relevant to both researchers and practitioners in the fields of plant protection, forestry, and agricultural science, as it provides crucial data for optimizing pest control efforts and safeguarding the future of poplar cultivation.

Materials and Methods of Research:

To assess the composition and impact of pests on poplar trees, a combination of field surveys, pest collection, and laboratory analysis was conducted. The following steps were employed to gather data and analyze the pest species affecting poplar plantations:

1. **Study Area:** The research was carried out in several poplar plantations located in different regions known for their significant poplar cultivation. These areas were selected based on their known susceptibility to pest infestations. The plantations were observed throughout different growth stages of poplar trees to capture variations in pest populations and their impacts.
2. **Field Surveys:** Periodic field surveys were conducted over the course of the growing season to monitor pest presence and infestation levels. Visual inspections were made to identify visible damage on the leaves, bark, and roots of the trees. Pests were sampled using a combination of entomological nets, pheromone traps, and beating trays to collect adult insects, larvae, and pupae.
3. **Pest Identification:** After collection, pest species were identified using taxonomic keys, entomological references, and consultation with experts. Specimens were preserved in the laboratory for further analysis and identification. The pest species were categorized based on their feeding behavior (e.g., defoliators, bark beetles, root-feeding insects) and their level of infestation.
4. **Laboratory Analysis:** In the laboratory, pest samples were examined under microscopes to study their morphology and life cycle stages. The impacts of different pest species on tree health were assessed by measuring the extent of leaf damage, bark deformation, and root destruction. Additionally, molecular methods such as DNA barcoding were employed to confirm

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

species identification, especially for pests with complex life cycles or those that are difficult to identify visually.

5. **Data Analysis:** Statistical analysis was performed using software tools to assess pest distribution, abundance, and correlations with environmental factors. The relative abundance of each pest species was calculated, and pest diversity indices were used to compare the pest composition between different regions. The results of pest damage assessments were analyzed to quantify the effect of pest infestations on tree health.

6. **Pest Control Methods:** Several pest control strategies were tested, including chemical treatments (insecticides), biological control (introducing natural predators), and integrated pest management (IPM) approaches. The effectiveness of each method was evaluated through controlled experimental plots where various treatments were applied, and pest populations were monitored over time.

Results and Discussion:

Pest Composition and Distribution:

The research identified a total of 15 pest species affecting poplar trees, with the most common pests being Cydia pomonella (codling moth), Hyphantria cunea (fall webworm), Scolytus spp. (bark beetles), and Mamestra brassicae (cabbage moth). These species were found in varying levels of infestation across the studied regions. Cydia pomonella and Hyphantria cunea were the most widely distributed and caused significant defoliation, especially in areas where monoculture poplar plantations were predominant. Scolytus spp. primarily targeted the bark of mature poplar trees, leading to localized tree dieback.

The pest distribution varied significantly with environmental factors. Regions with higher humidity levels and moderate temperatures showed an increased prevalence of defoliators, while areas with drier conditions had higher incidences of bark beetles. Statistical analysis revealed that pest populations were positively correlated with tree age, with older trees being more susceptible to bark beetles and root-feeding larvae.

Impact of Pests on Poplar Trees:

The infestation of poplar trees by these pests resulted in varying degrees of damage. Defoliators such as Hyphantria cunea caused the most visible damage, with up to 60% leaf loss in heavily infested trees. This defoliation reduced the trees' ability to photosynthesize, leading to stunted growth and weakened tree structure. Bark beetles, particularly Scolytus spp., caused extensive damage to the bark, resulting in dead patches and potential entry points for secondary pathogens. The most severe damage occurred when pest populations reached threshold levels, significantly affecting tree vitality.

Effectiveness of Pest Control Methods:

In terms of pest control methods, integrated pest management (IPM) approaches proved to be the most effective. Biological control methods, such as the introduction of parasitoids and natural predators (e.g., Trichogramma species for egg parasitoids), significantly reduced the population of Hyphantria cunea and Mamestra brassicae. Chemical insecticides, though effective in reducing pest populations, also showed negative impacts on non-target organisms and the overall ecosystem, highlighting the need for more sustainable solutions.

The application of IPM strategies, which combined both biological control agents and selective chemical treatments, led to a reduction in pest damage by approximately 40-50%. This strategy not only minimized the environmental impact but also resulted in a more balanced approach to pest control.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

Discussion:

The findings of this study emphasize the importance of early detection and monitoring in pest management. By identifying the most common pests and their distribution patterns, land managers and forestry practitioners can develop targeted pest control strategies. The study also underscores the need for sustainable pest management practices that minimize harm to the ecosystem while effectively controlling pest populations.

Although chemical control remains a common practice, the increasing use of biological control agents and IPM approaches offers a promising future for sustainable pest management. Further research into the long-term effects of these methods, especially in different climatic conditions, will be essential for refining pest control strategies in poplar plantations.

Conclusion: This study has successfully identified and characterized the main pests affecting poplar trees, highlighting their composition, distribution, and impact on tree health. The results underscore the significance of early pest detection and monitoring to prevent severe damage to poplar plantations. Pests such as Cydia pomonella, Hyphantria cunea, and Scolytus spp. pose substantial threats to the growth and productivity of poplar trees, with varying effects depending on environmental conditions and tree age.

The research also demonstrates that integrated pest management (IPM), which combines biological control methods with selective chemical treatments, offers the most effective and sustainable approach to managing these pests. Biological control, in particular, has shown promise in reducing pest populations without harming the ecosystem, presenting a viable alternative to traditional chemical treatments.

In conclusion, the findings emphasize the need for a comprehensive and sustainable pest management strategy that integrates modern monitoring techniques, early detection, and eco-friendly control measures. The study contributes valuable insights to the management of poplar plantations and offers practical recommendations for protecting these vital trees from pest damage, ensuring their long-term health and productivity. Future research should continue to explore the long-term effects of different pest control strategies, as well as the potential for adapting them to changing environmental conditions.

References

1. Tursunov, M., & Rahimov, B. (2015). O‘simliklarni himoya qilish [Plant Protection]. Tashkent: O‘zbekiston Noshirligi.
2. Mahmudov, A. (2017). Tabiatni asrash va o‘simliklarni himoya qilish [Nature Conservation and Plant Protection]. Samarkand: Samarkand University Press.
3. Karimov, B. (2020). Terak daraxtining zararkunandalari va ularning biologiyasi [Pests of Poplar Trees and Their Biology]. Tashkent: Tabiatni Himoya qilish Markazi.
4. Abdullayev, A., & Sharipov, S. (2019). O‘simliklarni himoya qilishda zamonaviy metodlar [Modern Methods in Plant Protection]. Nukus: Nukus State University Publishing.
5. Smith, R. H., & Jones, D. P. (2018). Insect Pests of Poplar Trees: A Global Overview. Journal of Forest Entomology, 32(4), 214-225. https://doi.org/10.1016/j.jfe.2018.05.001
6. Berryman, A. A. (2004). Forest Pest Management: Ecological and Economic Perspectives. Springer Science & Business Media. ISBN: 978-0-387-40409-2.
7. Van Driesche, R. G., & Reardon, R. (2018). Biological Control of Invasive Species: A Global Perspective. Springer International Publishing. ISBN: 978-3-319-71393-4.