ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

EPISOOTHIC STATE OF FASCIOLOSIS OF CATTLE IN SOME DISTRICTS OF THE SURKHANDARYA REGION

Assistant of the Department of Zooengineering, Veterinary Medicine and Sericulture of the Termez State University of Engineering and Agrotechnologies
Achilov Shamsiddin Amanovich

Assistant of the Department of Zooengineering, Veterinary Medicine and Sericulture of the Termez State University of Engineering and Agrotechnologies
Chorshambiyev Akbar Bozorovich

student of the Department of Veterinary Medicine of the Termez State University of Engineering and Agrotechnologies
Boypulatova Sevinch Azam qizi

Abstract. *This article analyzes the prevalence of trematodoses such as fascioliasis, dicrocoeliasis, and paramphistomiasis, which are widespread among agricultural animals, including cattle, in various natural and climatic conditions of Termiz, Jarkurgan, and Angor districts of the Surkhandarya region of our Republic. The study examines the distribution of these diseases in relation to the age of animals and the seasons of the year, as well as the extensiveness of helminthic invasion.*
Keywords. Trematodesis, fascioliasis, dicrocoeliosis, paramfi stomatosis, extensiveness of invasion, helminth coprology.

Relevance of the topic. There are a number of factors hindering the accelerated development of the livestock sector, increasing the number of livestock in all economic entities, increasing their productivity, implementing measures to combat and prevent various infectious, non-infectious, and parasitic diseases of livestock, and the development of this sector. In particular, among parasitic diseases of cattle, some helminthiases occupy an important place in terms of their spread and economic damage. These diseases include fascioliasis, dicrocoeliosis, and paramphistomatosis.

Degree of study of the problem. To determine the degree of helminthiasis spread in a specific area, the geographical location of that area (landscape natural climatic conditions), the presence of all links for disease spread, and the presence of an epizootic chain (definitive, intermediate, additional hosts) are taken into account. Economic damage from cattle trematodes is characterized by a decrease in milk and meat productivity of animals, forced slaughter, damage to the liver and gastrointestinal tract, unfitness for consumption, abortion, deterioration of breeding characteristics, and a certain decrease in the quality and energy value of dairy and meat products [6].

As a result of fascioliasis in cattle, a decrease in the immune properties of the body, a decrease in T- and B-lymphocytes, a decrease in the level of IgM due to a decrease in IgL and IgA, as well as the activity of AsAT, AlAT, and alkaline phosphatase were observed. At the same time, a decrease in the amount of protein, biogenic elements such as calcium, phosphorus, zinc, iron, copper, and manganese in blood plasma was observed. All this combined causes immunodeficiency in the animal organism. According to the authors, it is recommended to use anthelmintic drugs in combination with immunostimulants in combination with an early immunodiagnostic analysis in fascioliasis. The measures taken allow maintaining milk productivity of cows by 23.4-37.3 percent [12].

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

According to the analysis of research conducted by our scientists in various biocenoses of Uzbekistan, it was established that 186 species of trematodes parasitize vertebrates. Of these, 30 species were identified in the fish class, 14 species in the amphibian class, 2 species in the reptile class, 113 species in the bird class, and 27 species in the mammal class [8; 9].

According to the research conducted by the authors, trematodes of representatives of the order Artiodactyla - even-toed ungulates: Fasciola hepatica (L.,1758), F.gigantica (Cobbold, 1856), Dicrocoelium dendriticum (Rudolphi, 1819); Paramphistomum ichikawai (Fukui, 1922), Liorchis scotiae (Willmott, 1950); Liorchis hiberniae (Willmott, 1950); Calicophoron calicophorum (Fischoeder, 1906); Calicophoron erschowi (Davydova, 1959), Gastrothylax crumenifer (Creplin, 1874); Orientobilharzia turkestanica (Skryabin, 1913); *Hasstilesia ovis*, *Eurytrema pancreaticum*, *E. caelomaticum*, *Brachylaemus suis*, *Gastrodiscoides hominis*. Of the above 14 species of trematodes, 11 species were found in cattle (Bost aurus), 10 species in sheep (Ovis aries), 8 species in goats (Capra hircus), 2 species in domestic pigs (Sus scrofa domesticus), 3 species in wild boars (Sus csrofa), and 5 species of parasites were identified in representatives of the deer (Cervidae) family of wild herbivorous mammals [10].

According to the results of research conducted in livestock farms of the Samarkand region, located in the central part of the Republic of Uzbekistan, and the Syrdarya region, located in the northeastern part of the republic, 58.8% of cattle raised in the Samarkand region and 54.4% of cattle examined in the Surkhandarya region were found to have fascioliasis [4]. In each dairy cow affected by liver trematodes, there is an average decrease in milk productivity by 20-30 percent, miscarriage, decreased resistance to other diseases, and delays in lambing [2]. Up to 50-100% of young cattle, sheep, and goats grazed in pastures in areas susceptible to paramphistomatosis have been observed to die [7]. The acute course of paramphistomatosis was observed in livestock up to 2 years of age, and in sheep older than 2 years [3].

According to the author, in special studies conducted in the territory of the Republic of Karakalpakstan in 1989, 2000, 2001, the presence of 3 types of paramfi stomatomes - P.ichikawai, C.calicophorum, G.crumenifer - in cattle was noted, 16.0% of cattle were infected with gastrotiliacosis, 12.1% with paramfi stomosis, and 9.2% with calicophosis. At the same time, literature data from 2000 indicate that 4.6% of cattle and 2.6% of sheep in the Aral Sea region were affected by paramphigus stomatosis (G.cru mennifer) [4;5].

Purpose of the research. Analysis of the epizootic state of the main trematodes of cattle in some districts of the Surkhandarya region by seasons.

Place and object of research. Studies were conducted on cattle kept by the population in the Termez, Djarkurgan, and Angor districts of the Surkhandarya region, and samples of their feces were examined in the "Parasitology Laboratory" at the State Center for Animal Disease Diagnostics and Food Safety of the Surkhandarya region.

Research methods. The study used epizootological, clinical, helminthological, and partially pathoanatomical methods. Research results. Among the helminthiases caused by trematodes with the participation of two hosts, which are widespread among farm animals in Uzbekistan, fascioliasis, paramfistomatosis, and dicrocoeliosis are relatively rare, including schistosomiasis, eurytrematosis, and hapstylesiosis.

Two species of the causative agent of fascioliasis are widespread in the territory of the Republic of Uzbekistan: Fasciola hepatica (L., 1758) and Fasciola gigantica (Cobbold, 1856).

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

F.hepatica is considered a cosmopolitan species, as it is distributed in almost all countries of the world. The body length is 20-40 mm, width 8-14 mm, the shape is leaf-like, the shoulder parts are developed, and it is found in all regions of Uzbekistan, except for the Khorezm region and the Republic of Karakalpakstan. [1]

F.gigantica is distributed only in countries with a warm climate, including among the CIS countries in the Central Asian region, in the warm climates of Kazakhstan, Azerbaijan, Georgia, Armenia, and Russia, and in all irrigated, plain, and partially foothill-mountain agrobiocenoses of Uzbekistan. Its body length averages 30-70 mm, and its width is 4.0-11.0 mm. Both types of fascioli are hermaphrodite. [8]

When analyzing the epizootological state of fascioliasis in the Termez, Djarkurgan, and Angor districts of the Surkhandarya region, it was established that in the Termez district, up to 27.7% of calves under one year of age, 55.5% of heifers under two years of age, and 61.1% of cattle over three years of age were affected by fascioliasis. Of the 54 head of cattle examined, 26 were diagnosed with fascioliasis. The extent of invasion averaged 48.1%.

Accordingly, in the Djarkurgan district, the invasion extensiveness of fascioliasis was 31.25; 56.25; 62.5 percent. Of the 48 head of cattle examined, 24 were affected by fascioliasis, and the extensiveness of the invasion was 50%.

In the Angor district, the extensiveness of the invasion was 41.6; 50; 58.3 percent. Of the 36 head of cattle examined, 18 were infected with fascioliasis, and the average extent of invasion in the district was 41.5%. Out of 136 head of cattle in the region, 66 were diagnosed with fascioliasis, and the prevalence of the invasion was 45.5%.

Distribution of fascioliasis among cattle depending on their age in some districts of the Syrdarya region (coprological examination)

T/r	Districts	Animal age	Number of examined animals	Affected by fascioliasis	
				Number of affected animals	IE (%)
	Termez District	Up to 1 year old	18	5	27,7
		Up to 2 years old	18	10	55,5
		over 3 years old	18	11	61,1
total			54	26	48,1
	Jarkurgan District	Up to 1 year old	16	5	31,25
		Up to 2 years old	16	9	56,25
		over 3 years old	16	10	62,5
total			48	24	50
	Angor District	Up to 1 year old	12	5	41,6
		Up to 2 years old	12	6	50
		over 3 years old	12	7	58,3
total			36	18	49,9
By district			138	68	49,3

Analyzing the studies in 3 districts where the above studies were conducted, we believe that the extensiveness of invasion among animals of different ages is different, which is associated with the maintenance, feeding, watering of animals, and other factors. According to the results of our research,

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

it was observed that the extensiveness of trematode invasion among cattle increases proportionally to the age of the animal.

The main reason for this is that young animals are often kept at home, and it is impossible to detect them by helminth ovoscopy from the moment of initial infection until the parasite reaches sexual maturity.

Conclusion.

1. It was established that in irrigated agrobiocenoses of the Surkhandarya region, the prevalence of trematodoses of various ages among cattle is 47.5% due to fascioliasis, 28.75% due to dicrocoeliosis, and 28.5% due to paramphiostomatosis.

2. When analyzing cattle trematodes depending on the age of the animals, it was noted that the incidence of trematodes is high in cattle aged 3 years and older.

List of references:

1. Daminov A.S. “Respublikaning turli biogeosenozlarida qoramollar trematodozlarining epizootologik va immunologik xususiyatlari”. Vet.fan.dok...., diss. Samarqand: SamVMI, 2016. - 200 b.

2. Salimov B.S., Daminov A.S., Qurbonov Sh.X., Iza tullayev Z.X. Qishloq xo‘jalik hayvonlarining jigar trematodozlari, ularni davolash va oldini olish chora-tadbirlari (tavsiyanoma). Toshkent, 2009. –B. 7.

3. Ерболатов, К.М. Парамфистоматозы. – Алма-Ата: Кайнар, 1982. – С.183-238.

4. Кожабоев М. «Ассоциация инвазия трематода крупного рогатого скота «Приаралья». //Автореферат. дисс. канд. биол. наук. Институт Зоологии Ан.Уз 2001. -С. 24.

5. Кожабоев М.К. другие. «Биология возбудителя и эпизоотологии ориентобильгорциоизы животных». Тезисы науч. конф. «Гельминтология сегодня, проблемы и перспективы». Москва, 1989. -С. 158.

6. Кумышева, Ю.А. Морфологическая характеристика говядины при дикроцелиозе крупного рогатого скота / Ю.А. Кумышева, А.М. Мазихова // Вестник КрасГАУ. - 2009. - №5. – С.122-125.

7. Подлесный, Г.В. Экономический ущерб при лиорхозе крупного рогатого скота / Г.В. Подлесный // Ветеринария. – 1960. – № 5. – С. 18-19.

8. Султанов М.А., Азимов Д.А., Гехтин В.И., Муминов П.А. Гельминты домашних млекопитающих Узбекистана. Тошкент, «Фан», 1975. - 186 с.,

9. Шакарбоев Э.Б., Шакарбоев У.А., Дадаев С.Д., Махамадиев З.С. Жиззах вилоятининг турли минтақаларида қўйлар гельминтозларининг фаунаси ва экологияси. // Зоо ветеринария. - Тошкент, 2015. №6. - Б.17-19.

10. Шакарбоев Э.Б., Акрамова Ф.А., Азимов А.А. Тематоды - паразиты позвоночных Узбекистана. // Ташкент, 2012. -215 с.

11. Шихалиева, М.А. Сезонная динамика зараженности овец и крупного рогатого скота смешанной инвазией, вызванной Fasciola hepatica и Dicrocoelium lanceatum, в условиях Северного Кавказа // Российский паразитологический журнал. - 2014. - № 1. - С. 81-84.

12. Якубовский М.В., Шемелева Н.Ю. Особенности иммунитета и профилактика фасциолеза крупного рогатого скота. //Ветеринария.Москва, 2011. -С. 34-38.