ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

METHODS OF PREVENTING DISPEPSIA IN CALVES.

assistant of the Department of Zooengineering, Veterinary Medicine and Sericulture of the Termez State University of Engineering and Agrotechnologies
Davlatov Karim Mamadmuso ugli

Termez State University of Engineering and Agrotechnologies
Usmanov Rustam Dilmurod ugli,
student of the Department of Veterinary Medicine

Termez State University of Engineering and Agrotechnologies
Anorov Sardor Botir ugli
, student of the Department of Veterinary Medicine

Abstract: The article presents data on metabolic disorders in the body of cows excluded from milk as a consequence of etiological factors in calves dyspepsia and it's prevention by adding sprouted verdure of the wheat, monocalcium phosphate and LPP-1 to the diet of cows.

Keywords:hypotrophy, dyspepsia, diarrhea, hemoglobin, carotene, calcium, phosphorus, monocalciphosphate, LPP-1, trace elements, vitamins, prophylaxis, glucose, protein, leukocyte**.**

Meeting the population's demand for livestock products through the accelerated development of animal husbandry is a priority of modern agrarian policy.

In recent years, many pedigree heifers and heifers have been imported to our republic from foreign countries, and the breed of cattle has been improved.

The next task facing livestock farmers and veterinary specialists is to preserve the genetic potential of animals for high productivity through proper care and replenish the herd with healthy calves.

Without fully satisfying the physiological needs of the cow's body, it is impossible to raise healthy and viable calves. Calves born in a hypotrophic state are more susceptible to dyspepsia.

Analysis of literature data shows that to date, in the conditions of our republic, dyspeptic disease among newborn calves is widespread (Eshburiev B.M., 2003, 2014), and despite the fact that in some farms the incidence rate reaches 46% and the mortality rate up to 32%, effective methods for preventing dyspepsia in calves have not been developed in cattle farms.

In order to improve methods for the prevention of dyspepsia in calves, research was conducted on 10 Holstein cows of 3-4 years of age and calves born from them at the "Janub elita nasilchilik" farm in the Termez district. Clinical and hematological studies were conducted on the experimental cows once a month, and the incidence of dyspepsia in the calves born from them was studied.

The composition of the cows' ration was checked for feed units, digestible protein, sugar, carotene, calcium, and inorganic phosphorus. Through clinical examination of the experimental cows, their appetite, skin and skin covering, visible mucous membranes, condition of lymph nodes, body temperature, respiratory rate and pulse rate per minute, and abdominal wall movement rate per 2 minutes were determined. The physiological development of the calves, their response to external stimuli, milk intake, mucous membranes, skin and skin covering, lymph nodes, functional state of the respiratory, cardiovascular, digestive and excretory systems, frequency of manuring, consistency and color of manure were determined. The calves' body weight was determined by weighing on scales.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

In the blood of the experimental cows, the number of erythrocytes and leukocytes (in the Goryaev counting grid), hemoglobin (Hemoglobin-cyanide method), glucose (color reaction with ortho-toluidine), and total protein in blood serum (Refractometric method) were determined by laboratory testing.

The cows selected for the experiments were divided into two groups of 5 heads, one of which was designated as the experimental group and the other as the control group. In addition to the ration of cows of the experimental group at 8-9 months of pregnancy, grown after milking, 1 kg of sprouted wheat, 50 g of monocalcium phosphate, and 40 g of LPP-1 per head per day were added. Cows of the control group were fed only on the farm ration. The experiments lasted 60 days.

When determining the economic efficiency of the work, the methodological guide "Determination of the Economic Efficiency of Veterinary Measures" (M.Kh. Shaykhamanov et al., 1987) was used. In this case, the body weight of calves obtained from cows and the milk productivity of cows were taken into account as criteria for economic efficiency.

Obtained results. The cows' ration is of the silage-haylage-concentrate type, consisting of corn silage, haylage, alfalfa hay, cotton hulls, cottonseed meal, and mixed feed. The nutritional value of the ration is 8.60 feed units. Compared to feeding norms, the nutritional value of the ration was 1.4 units less, there was a deficiency of digestible protein by 110 g, sugar by 400 g, carotene by 261 mg, phosphorus by 15.3 g, and an excess of calcium by 15.2 g and fiber by 1363 g. The satisfaction of the cows' body's needs for nutrients was 84.3%, digestible protein - 89%, sugar - 52.9%, carotene - 46.3%, phosphorus - 74.5%, calcium - 116%, and fiber - 133%.

The carbohydrate part of the diet was characterized by a deficiency of sugar and an excess of fiber compared to feeding norms. An imbalance of protein and energy components was also revealed, i.e., the sugar-protein ratio in the diet was 0.51 instead of 0.8-1.5:1.

The macroelement part of the diet was characterized by an imbalance in their ratio due to excess calcium and phosphorus deficiency. That is, the ratio of phosphorus to calcium was 0.42:1 (norm - 2.0:1). In some animals, the occurrence of hypotonia in the peritoneum is explained by the confinement of animals in one place, the absence of motility, the feeding of fiber-rich food, its difficult digestion, and low nutritional value.

Cows' feed ration

Types of food	Quantity (kg)	Food unit	Digested protein (g)	Carotene, mg	Sugar (g)	Fiber (g)	Sa, g	R, g
Corn silage	9	2,22	186	65	126	1860	36,0	14,0
Alfalfa Hay	3	1,20	2,05	100	210	1160	50,6	9,8
Haylage	3	2,40	194	60	92	320	2,4	9,0
Cotton husk	4	1,12	84	-		1840	9,2	2,8
Cottonseed meal	2	1,66	221	-	22	300	7,0	8,8

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

Total:		8,60	890	225	450	5480	105,2	44,7
Norm		10,2	1000	486	850	4117	90	60
Difference ±		- 1,4	-110	-261	-400	+1363	+15,2	-15,3

Some morphobiochemical indicators of the blood of the experimental cows before the start of the experiments were characterized by the same indicators in the experimental and control groups, while in the control group, morphobiochemical indicators decreased compared to normal indicators until the end of the experiments.

By the end of the experiments, an increase in the number of erythrocytes in the blood of cows in the experimental group by 0.91 million/µl, hemoglobin - by 10.4 g/l, glucose - by 0.28 mmol/l, total protein - by 6.2 g/l was noted compared to the initial indicators. An increase in the number of erythrocytes and the amount of hemoglobin in cows of this group can be explained by an increase in hematopoiesis, the amount of glucose, total protein, and an improvement in metabolism in the body. In the control group, by the end of the experiments, a decrease in the number of erythrocytes in the blood by an average of 0.20 million/µl, hemoglobin by 2.6 g/l, glucose by 0.09 mmol/l, and total protein by 0.7 g/l was revealed compared to the initial indicators. We explain such changes by the fact that the needs of the cow's body for nutrients are not met.

The average body weight of calves born from cows in the experimental group was 31 kg, and the average body weight of calves born from cows in the control group was 27 kg. The growth and development of the calves were monitored for a month. Calves born from cows of the control group were affected by dyspepsia by the 3rd day after birth. One of the calves in the experimental group suffered from dyspepsia. After one month, the average body weight of calves born from cows in the experimental group was 41 kg, and the average body weight of calves born from cows in the control group was 37 kg. Thus, it has been experimentally proven that the body weight of calves born from cows in the experimental group is 10.8% higher than the body weight of calves in the control group. The milk productivity of cows in the first month of lactation averaged 10.85 kg in the experimental group and 8.84 kg in the control group.

Conclusion: In order to prevent dyspepsia in calves, the addition of 1 kg of wheat grass grown per cow, 50 g of monocalcium phosphate and 40 g of LPP-1 to the ration of pregnant cows that have been weaned for 60 days improves the processes of metabolism in the body of cows and leads to the birth of healthy calves.

References.

1. Abramov S.S., Arestov I.G., Karput I.M. "Prevention of non-communicable diseases of young people." M.: Agropromizdat. 1990 - 92 p.

2. Norboev Q.N., B.Bakirov, B.Eshbo'riyev: Pathology and Therapy of Non-Infectious Internal Diseases of Young Animals.

5. Norboev Q.N., B.Bakirov, B.Eshbo'riyev: "Internal Non-Infectious Diseases of Animals" Samarkand, 2007.

6. Eshburiev B.M. Clinical and hematological changes in neonatal dyspepsia of calves "Zooveterinary" 2009. No2. P. 16-17.

7. Eshburiev B.M., Norboev K.N., Eshburiev S.B. Recommendations for the treatment and prevention of dyspepsia in calves. DVBB RUz. 06.03.2009 "N.Doba" XT. Samarkand, 2009.