ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

FERTILITY CHARACTERISTICS OF GOLSHTIN BREED COWS OF DIFFERENT BREEDINGS.

Termez State University of Engineering and Agrotechnologies Faculty of Economics, Forestry and Veterinary Medicine Assistant of the Department of Zooengineering, Veterinary Medicine and Sericulture
Kurbanov Shaymardon Bobokulovich

Termez State University of Engineering and Agrotechnologies
Faculty of Economics, Forestry and Veterinary Medicine
Assistant of the Department of Zooengineering, Veterinary Medicine and Sericulture
Gulbutaev Bekmurod Khushmurod ugli
Kamolova Nozliya Jahongirovna,
1st year student of group 117, Zooengineering (by type)

Annotation: *the study of the age of the animals in the experimental groups at first insemination showed that the age of the animals in group II at first insemination was I and*
They were 5.5 (P>0.99) and 2.1 (P>0.95) days shorter than their peers in groups II, respectively, and the live weight at the first fertilization was 10.0 (P>0.99) and 15.5 (P>0.999) kg higher in group I than in groups II and III, respectively. The gestation period of heifers in group I was 16 days shorter than in group II, while in group III there was no significant difference Also, the age of first calving in cows of group II was on average 20.9 (P>0.999) 16.9 days (P>0.99) shorter than in cows of groups I and III, respectively, and their live weight in cows of group I was 26.7 (P>0.99) and 35.1 (P>0.999) kg higher than in cows of groups II and III, which indicates that the reproductive characteristics of the cows were at the level of biological norms.

Keywords: *Fertility, pregnancy, cattle, cow, of various breeds, service life, fertility rate, escape index.*

Introduction. In cattle breeding, especially in dairy cattle breeding, the study of the reproductive characteristics of cows is of great importance in improving the breeding qualities of cows and improving the herd composition.

Therefore, inseminating cows after calving at optimal times is of great practical importance. Considering this

achieving high milk productivity of cows with minimal labor costs and minimal consumption of material resources is one of the most important goals facing dairy cattle breeding. Considering that cows start producing milk only after calving, the required level of their reproductive characteristics indicates that the reproductive characteristics of cows in the entire herd are the basis of milk production efficiency [1; 2].

When assessing the fertility characteristics of cows, along with their service period, the period from weaning to calving is also of great importance. During the lactation period, their diet often fails to meet the body's nutrient requirements, especially when feeding high-yielding cows. Therefore, most nutrients, especially minerals, are consumed at the expense of reserves in the cow's body. Therefore, during this period, it is especially important to create a reserve of nutrients to cover the consumption

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

of these reserves. During this period, the cow can produce an additional 15-20 kg of milk per kilogram of live weight gain during postpartum lactation. Therefore, it is advisable to give cows rest for the next calving period of 60 days. In dairy cattle breeding, the study of the reproductive characteristics of cows plays an important role in improving the breeding qualities of cows and improving the herd composition [3].

Also, the level of productivity and fertility indicators of cows depend on their age at first calving and live weight. Therefore, the study and improvement of their reproductive characteristics is of particular importance for breed improvement [4]. **The purpose of the research is to study the breeding characteristics of Holstein cows of various breeds in the conditions of the Surkhan oasis.**

Location of research. Research In 2024-2025, it was carried out at the dairy farm of the "Navruz-naslchilik" livestock complex under JSC "Uzbekistan Railways," located in the Termez district of Surkhandarya region.

Discussion of the obtained results. The origin of the cows of the experimental groups and their belonging to different breeds were studied from primary documents, that is, from the breeding certificates of each cow (form MOL-2). For the experiment, based on similar traits, 3 groups of Holstein cows of three or more lactations of different breeds were selected, group I - cows of German breeds, group II - cows of Polish breeds, and group III - cows of Belarusian breeds of three or more lactations. We studied the reproductive characteristics of cows of the Holstein breed of different breeds (Table 1).

Table 1

Fertility indicators of cows of different breeds (p=10)

Indicators	Groups					
	I		II		III	
	$\overline{X} \pm S\overline{x}$	Sv, %	$\overline{X} \pm S\overline{x}$	Sv, %	$\overline{X} \pm S\overline{x}$	Sv, %
Age at first conception, days	552,2 + 5,8	5,78	557,2 + 4,7	2,67	555,1 + 7,2	4,07
Live weight at first fertilization, kg	372,5 + 1,7	1,45	362,5 + 3,3	2,85	357,0 + 2,4	2,11
Duration of pregnancy, days	283,3 + 8,5	9,53	299,3 + 11,6	9,53	284,4 + 8,2	9,15
Age at first delivery, days	835,5 + 11,4	4,31	856,4 + 9,4	3,46	839,5 + 3,7	1,40
Live weight at first birth, kg	465,0 + 7,1	4,84	438,1 + 6,5	4,69	429,9 + 3,4	2,49
Service period duration, days	78,8 + 1,3	5,28	81,1 + 1,6	6,16	79,5 + 1,5	5,85
fertility rate from 1st insemination, %	76	-	72	-	70	-
Escape Index	1,38	-	1,53	-	1,41	-

As can be seen from Table 1, the study of the age of cattle in the experimental groups at first insemination showed that the age of cattle in group II at first insemination was 5.5 (P>0.99) and 2.1 (P>0.95) days shorter than their peers in groups I and II, respectively, and the live weight of cattle in group I at first insemination was 10.0 (P>0.99) and 15.5 (P>0.999) kg higher than their peers in groups II and III, respectively. The gestation period of heifers in group I was 16 days shorter than in group II, while in group III there was no significant difference. Also, the age of first calving in cows

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-4

of group II was on average 20.9 (P>0.999) 16.9 days (P>0.99) shorter than in cows of groups I and III, respectively, and their live weight in cows of group I was higher than in cows of groups II and III. 26.7 (P>0.99) and 35.1 (P>0.999) kg higher.

The duration of the service period of cows in group I was shorter by 2.3 (P>0.99) and 0.7 days, respectively, than in cows of groups II and III, which indicates that the cows of all groups had good fertility characteristics.

Thus, the reproductive characteristics of the cows of the experimental group were within biological norms**.** This indicates that the results of the conducted research have positive indicators.



At the "Navruz-naslchilik" livestock complex
Feeding of breeding cows.

List of used literature.

1. Ashirov M.E. Breeding of Dairy Cattle. //Tashkent. 2017, "Navruz" Publishing House. 371 p.
2.Ashirov M.I., Ashirov B.M., Yuldashev A.A. Breeding of Holstein cattle in Uzbekistan. //Tashkent. 2020, "Navruz" Publishing House. p. 264.
3. Gridina S.L. Reproductive ability of black-feathered Ural type cows. //J. "Zootechnics." 2005, No. 3, p. 30-31.
4. Amirov Sh.K., Boybulov B.Sh., Pardaev R., Shodieva U. Fertility characteristics of cows in the experimental group. // "Prospects for the Development of Animal Husbandry in the Republic," Materials of the Republican Scientific and Practical Conference. Tashkent, 2019, pp. 81-83.