ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

REAGENTS IN THE ANALYSIS OF ALKALOIDS: A COMPREHENSIVE REVIEW

RAKHMATILLAEVA INDIRA ANVAROVNA[1]
OLTIEVA SHOKHISTA POLVANNAZIROVNA[1]
Urgench branch of Tashkent medical academy[1],
Assistant of the department of medical biology and pharmacy

Abstract. Alkaloids represent a diverse class of naturally occurring nitrogen-containing compounds with significant pharmaceutical and toxicological properties. This review examines the various reagents employed in the qualitative and quantitative analysis of alkaloids, focusing on their chemical principles, applications, and limitations. The classical precipitation reagents, color-forming reagents, and modern analytical techniques are discussed in detail. Special attention is given to the selectivity, sensitivity, and specificity of these reagents for different alkaloid classes. This comprehensive overview serves as a valuable resource for researchers, pharmacognosists, forensic scientists, and quality control specialists working with alkaloid-containing natural products and pharmaceutical preparations.

Keywords: alkaloids, precipitation reagents, color reagents, chromatography, spectroscopy, analysis

Introduction

Alkaloids constitute one of the most diverse groups of natural products with over 12,000 structures identified to date. These nitrogen-containing compounds, predominantly derived from plants but also found in fungi, bacteria, and animals, exhibit remarkable structural diversity and biological activities. The analytical determination of alkaloids is essential in pharmaceutical research, quality control of herbal medicines, forensic toxicology, and ecological studies.[1,2]

The history of alkaloid analysis dates back to the early 19th century when Friedrich Sertürner isolated morphine from opium in 1805. Since then, numerous methods have been developed for alkaloid detection, identification, and quantification. These methods rely heavily on specific reagents that interact with alkaloids to produce characteristic precipitates or color reactions, or facilitate their detection through modern instrumental techniques.

This review provides a comprehensive examination of the reagents used in alkaloid analysis, discussing their chemical basis, applications, advantages, and limitations. The information compiled herein aims to guide researchers in selecting appropriate reagents for specific analytical needs in alkaloid research.

General principles of Alkaloid analysis

Alkaloid analysis typically involves extraction from the natural matrix, followed by detection, identification, and quantification. The selection of appropriate reagents depends on several factors:

1. The chemical structure of the alkaloid(s) under investigation
2. The complexity of the sample matrix
3. The required sensitivity and specificity
4. The availability of analytical equipment
5. The purpose of the analysis (qualitative screening vs. precise quantification)

Precipitation reagents

Mayer's reagent (Potassium Mercuric Iodide)

Composition: Potassium mercuric iodide solution (K_2HgI_4)

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

Principle: Mayer's reagent forms complexes with alkaloids through coordination with the nitrogen atom, resulting in white or yellowish precipitates.

Applications: This reagent is widely used for preliminary alkaloid screening in plant extracts. It reacts positively with most alkaloids but does not distinguish between different types.

Sensitivity: The detection limit varies between 1-10 µg of alkaloid, depending on the specific compound.

Limitations: Non-specific reactions can occur with proteins, amines, and other nitrogen-containing compounds.

Wagner's reagent (Iodine-Potassium Iodide)

Composition: Iodine dissolved in potassium iodide solution (I_2-KI)

Principle: Wagner's reagent forms brown precipitates with alkaloids through the formation of periodides.

Applications: Used for alkaloid detection in plant extracts and pharmaceutical preparations, particularly effective for quaternary and tertiary alkaloids.

Sensitivity: Detection limit typically ranges from 5-50 µg.

Limitations: May give false positives with other nitrogenous compounds.

Dragendorff's reagent (Potassium Bismuth Iodide)

Composition: Potassium bismuth iodide solution ($K[BiI_4]$)

Principle: Forms orange-red precipitates with alkaloids through coordination complexes.

Applications: Widely used in thin-layer chromatography (TLC) for visualizing alkaloids and for preliminary screening in plant extracts.

Sensitivity: Can detect as little as 1 µg of some alkaloids.

Limitations: Gives positive reactions with quaternary ammonium compounds and some non-alkaloid substances.

Hager's reagent (Saturated Picric Acid Solution)

Composition: Saturated aqueous solution of picric acid (2,4,6-trinitrophenol)

Principle: Forms yellow crystalline precipitates with alkaloids through salt formation.

Applications: Used for detection and semi-quantitative determination of alkaloids.[3]

Sensitivity: Varies considerably depending on the alkaloid structure.

Limitations: Less selective than other precipitation reagents.

Tannic acid

Composition: Aqueous solution of tannic acid

Principle: Forms tannates with alkaloids, resulting in brown precipitates.

Applications: Used primarily for qualitative detection of alkaloids in crude extracts.

Limitations: Low specificity, as tannins precipitate many other compounds including proteins.

Color-Forming reagents

Color reactions are crucial in alkaloid analysis, providing valuable information about structural features and facilitating detection in chromatographic methods.

Marquis reagent

Composition: Formaldehyde in concentrated sulfuric acid

Principle: Produces characteristic colors with various alkaloids through condensation reactions, followed by oxidation.

Applications: Primarily used for detection of opium alkaloids, phenethylamines, and indole alkaloids. Morphine produces a purple color that changes to deep violet.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

Sensitivity: Can detect microgram quantities of many alkaloids.
Specificity: Different alkaloid classes produce distinct colors, aiding in preliminary identification.
Froehde's reagent
Composition: Molybdic acid in concentrated sulfuric acid
Principle: Forms colored complexes through oxidation reactions and complex formation.
Applications: Particularly useful for opium alkaloids; morphine gives a violet color, while codeine produces a green-blue color.
Limitations: The colors fade relatively quickly and may vary with reagent concentration and age.

Mandelin's reagent

Composition: Ammonium vanadate in concentrated sulfuric acid
Principle: Forms colored complexes through oxidation and coordination.
Applications: Used for detection of opium alkaloids, lysergic acid derivatives, and phenethylamines.
Sensitivity: Typically in the microgram range.
Limitations: Color stability issues and reagent deterioration over time.

Ehrlich's reagent

Composition: p-Dimethylaminobenzaldehyde in acidic medium
Principle: Reacts with indole alkaloids to form colored Schiff bases.
Applications: Specific for indole alkaloids (ergot alkaloids, tryptamines) and compounds with a pyrrole ring.
Specificity: Relatively high for indole-containing structures.
Limitations: Limited to certain structural types of alkaloids.

Van Urk's reagent

Composition: p-Dimethylaminobenzaldehyde in sulfuric acid with ferric chloride
Principle: Modified Ehrlich's reagent that produces more intense colors with indole alkaloids due to the oxidizing effect of ferric ions.
Applications: Detection of ergot alkaloids and other indole derivatives.
Sensitivity: More sensitive than standard Ehrlich's reagent.
Dragendroff-Munier reagent
Composition: Modified Dragendorff's reagent with additional chemicals
Principle: Produces orange-red colors with alkaloids, enhanced sensitivity compared to standard Dragendorff's reagent.
Applications: Used in TLC and paper chromatography for alkaloid detection.
Sensitivity: Can detect nanogram quantities of some alkaloids.

Spectrophotometric reagents

Reagents that facilitate spectrophotometric determination of alkaloids are essential for quantitative analysis.

Bromocresol Green (BCG)

Composition: 3,3',5,5'-Tetrabromo-m-cresolsulfonephthalein
Principle: Forms ion-pair complexes with alkaloids that can be extracted into organic solvents and measured spectrophotometrically.
Applications: Quantitative determination of various alkaloids including quinine, atropine, and berberine.
Sensitivity: Detection limits in the range of 0.5-5 μg/mL.
Limitations: Potential interference from other basic compounds.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

Methyl Orange

Composition: Sodium 4-[(4-dimethylamino)phenylazo]benzenesulfonate

Principle: Forms extractable ion-pairs with alkaloids, measurable by spectrophotometry.

Applications: Quantification of berberine, palmatine, and other quaternary alkaloids.

Sensitivity: Detection limits typically 1-10 µg/mL.

Congo red

Composition: Sodium 3,3'-([1,1'-biphenyl]-4,4'-diyl)bis(4-aminonaphthalene-1-sulfonate)

Principle: Forms colored complexes with certain alkaloids.

Applications: Used for quantitative determination of some isoquinoline alkaloids.

Limitations: Limited application range compared to other dyes.

Reinecke's salt

Composition: Ammonium tetrathiocyanatodiamminechromate(III)

Principle: Forms colored precipitates with alkaloids that can be dissolved and measured spectrophotometrically.

Applications: Quantification of quaternary ammonium alkaloids.[4]

Sensitivity: Moderate sensitivity, typically requiring microgram quantities.

Reagents in chromatographic analysis

Modern alkaloid analysis heavily relies on chromatographic techniques, which require specific reagents for optimal separation and detection.

TLC Visualization reagents

Dragendorff-Munier reagent

As previously described, this modified Dragendorff's reagent provides enhanced sensitivity for TLC detection of alkaloids.

Iodoplatinate reagent

Composition: Chloroplatinic acid and potassium iodide

Principle: Forms blue-violet complexes with alkaloids.

Applications: Detection of alkaloids on TLC plates with high sensitivity.

Sensitivity: Can detect nanogram quantities of some alkaloids.

Cerium(IV) Sulfate

Composition: Cerium(IV) sulfate in sulfuric acid

Principle: Oxidizes alkaloids to form colored compounds.

Applications: General detection reagent for organic compounds, including alkaloids.

Limitations: Lacks specificity for alkaloids.

HPLC Mobile Phase Additives

Ion-Pairing reagents

Examples: Sodium 1-heptanesulfonate, tetrabutylammonium phosphate, triethylamine

Principle: Form ion pairs with charged alkaloids to improve retention and peak shape in reversed-phase HPLC.

Applications: Separation of quaternary alkaloids and other ionic alkaloids.

Advantages: Improves peak symmetry and resolution for basic compounds.

pH Modifiers

Examples: Formic acid, phosphoric acid, ammonia, diethylamine

Principle: Control ionization state of alkaloids to optimize retention and peak shape.

Applications: Essential for reproducible HPLC analysis of alkaloids.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

Considerations: Selection depends on detection method and alkaloid properties.

Derivatization reagents for GC analysis

Silylating reagents

Examples: BSTFA (N,O-bis(trimethylsilyl)trifluoroacetamide), MSTFA (N-methyl-N-trimethylsilyltrifluoroacetamide)

Principle: Substitute active hydrogens in alkaloid molecules with trimethylsilyl groups to improve volatility and thermal stability.

Applications: GC analysis of alkaloids with hydroxyl, carboxyl, or amino groups.

Limitations: Not applicable to quaternary alkaloids.

Acylating Reagents

Examples: Trifluoroacetic anhydride, acetic anhydride

Principle: Form volatile derivatives with functional groups in alkaloids.

Applications: Improve GC properties of alkaloids with hydroxyl or amino groups.

Advantages: Can enhance detector response in addition to improving chromatographic properties.

Reagents in Mass spectrometric analysis

Mass spectrometry has become the method of choice for alkaloid identification, requiring specific reagents to optimize analysis.

Derivatization reagents for Enhanced MS detection

Quaternization reagents

Examples: Methyl iodide, benzyl chloride

Principle: Convert tertiary amine alkaloids to quaternary ammonium salts to enhance ESI-MS detection.

Applications: Improve sensitivity in LC-MS analysis of alkaloids.[5]

Advantages: Can lower detection limits by 1-2 orders of magnitude.

Stable Isotope Labeling reagents

Examples: Deuterated methyl iodide (CD_3I), ^{13}C-labeled reagents

Principle: Introduce stable isotopes into alkaloid molecules for quantitative analysis and structural studies.

Applications: Quantitative determination of alkaloids using isotope dilution MS.[6]

Advantages: High accuracy and precision in quantitative analysis.

Matrix compounds for MALDI-MS

Examples: α-Cyano-4-hydroxycinnamic acid, 2,5-dihydroxybenzoic acid

Principle: Facilitate ionization of alkaloids in matrix-assisted laser desorption/ionization mass spectrometry.

Applications: Analysis of alkaloids in complex matrices.

Limitations: Matrix selection is critical for successful analysis.

Reagents for Immunochemical analysis

Immunochemical methods offer high specificity and sensitivity for alkaloid detection, particularly in complex biological samples.

Reagents for Alkaloid-protein conjugation

Examples: Carbodiimides, glutaraldehyde, succinic anhydride

Principle: Link alkaloids to carrier proteins to produce immunogens for antibody production.

Applications: Development of antibodies against low molecular weight alkaloids.

Challenges: Preserving the epitope structure of the alkaloid during conjugation.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

Enzyme conjugates

Examples: Alkaloid-horseradish peroxidase conjugates, alkaloid-alkaline phosphatase conjugates

Principle: Used in competitive enzyme immunoassays for alkaloid detection.

Applications: High-throughput screening of alkaloids in biological samples.

Advantages: High sensitivity and adaptability to various detection formats.

Modern trends in Alkaloid analysis reagents

Recent advances in analytical chemistry have introduced novel reagents and approaches for alkaloid analysis.[7]

Fluorescent derivatization reagents

Examples: Dansyl chloride, fluorescamine, 9-fluorenylmethyl chloroformate (FMOC-Cl)

Principle: Form fluorescent derivatives with alkaloids for sensitive detection.

Applications: HPLC analysis with fluorescence detection, offering improved sensitivity and selectivity.

Advantages: Detection limits in the nanogram to picogram range.

Molecularly Imprinted Polymers (MIPs)

Principle: Synthetic polymers designed to have specific binding sites for target alkaloids.

Applications: Sample preparation, chromatographic separation, and sensing of alkaloids.[8]

Advantages: High selectivity, stability, and reusability.

Nanomaterial-Based reagents

Examples: Gold nanoparticles, quantum dots, carbon nanotubes

Principle: Interact with alkaloids through various mechanisms to produce detectable signals.

Applications: Colorimetric, fluorescent, and electrochemical detection of alkaloids.[9]

Advantages: Ultra-high sensitivity, potential for multiplexed detection.

Future perspectives

The field of alkaloid analysis continues to evolve with advances in analytical chemistry and materials science. Several promising directions include:

1. Development of more selective and sensitive reagents for specific alkaloid classes
2. Integration of artificial intelligence for optimizing reagent selection and method development
3. Miniaturization and automation of analytical procedures using microfluidic platforms
4. Green analytical chemistry approaches using environmentally friendly reagents
5. Point-of-care alkaloid detection systems for field applications

These developments will likely enhance the capabilities of researchers to detect, identify, and quantify alkaloids in complex matrices with improved accuracy, sensitivity, and throughput.

CONCLUSION

Reagents play a critical role in the analytical chemistry of alkaloids, enabling their detection, identification, and quantification in various matrices. From classical precipitation and color reagents to modern spectroscopic, chromatographic, and immunochemical approaches, the choice of appropriate reagents significantly impacts the success of alkaloid analysis. This review has summarized the principles, applications, advantages, and limitations of major reagent classes used in alkaloid analysis, providing a comprehensive resource for researchers in this field.

As analytical techniques continue to advance, new reagents and methodologies will emerge, further enhancing our ability to study these fascinating natural products. The ongoing development of more selective, sensitive, and environmentally friendly reagents will continue to drive progress in alkaloid analysis and expand its applications in pharmaceutical, toxicological, and natural product research.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

REFERENCES

1. Cordell, G. A. (2013). Fifty years of alkaloid biosynthesis in Phytochemistry. Phytochemistry, 91, 29-51.
2. Mokhtar, S. U., Abdulrahman, A. L., & Al-Raqobah, N. A. (2015). Comparative study of the traditional methods for the quantitative determination of alkaloids. Asian Journal of Chemistry, 27(1), 221-225.
3. Chauhan, A., Goyal, M. K., & Chauhan, P. (2013). GC-MS technique and its analytical applications in science and technology. Journal of Analytical & Bioanalytical Techniques, 4(4), 1-5.
4. Zhong, X., Huang, Y., Zhong, Y., Liang, Y., & Li, X. (2017). A colorimetric method for mercury (II) detection with lysozyme regulated growth of gold nanoparticles. Analytical Methods, 9(13), 1995-2002.
5. Tian, Y., Zhang, C., Guo, M., & Zhang, J. (2014). Comparative analysis of Amaryllidaceae alkaloids from three Lycoris species. Molecules, 19(9), 13909-13919.
6. Ciura, K., Dziomba, S., Nowakowska, J., & Markuszewski, M. J. (2017). Thin layer chromatography in drug discovery process. Journal of Chromatography A, 1520, 9-22.
7. Siddiqui, M. R., AlOthman, Z. A., & Rahman, N. (2017). Analytical techniques in pharmaceutical analysis: A review. Arabian Journal of Chemistry, 10, S1409-S1421.
8. Tiwari, M. P., & Uzun, L. (2017). Molecularly imprinted polymers for selective analysis of chemical compounds in food and beverages. In Biosensors for Sustainable Food-New Opportunities and Technical Challenges (pp. 153-176). Elsevier.
9. Khan, M. Z. H., Liu, X., Tang, Y., Liu, X., & Zhang, H. (2018). Ultra-sensitive electrochemical sensing platform for detection of multiple alkaloids in herbal plants. Sensors and Actuators B: Chemical, 266, 520-532.
10. Anvarovna R. I. POISONS EXTRACTED FROM DIALYSIS OBJECTS USING WATER //Multidisciplinary Journal of Science and Technology. – 2025. – Т. 5. – №. 4. – С. 232-236.
11. Davronovna, A. K., & Bekboevich, K. S. (2025). GENERAL CHEMISTRY PROBLEMS OF MEDICAL STUDENTS AND STRATEGIES FOR THEIR SOLUTION. Multidisciplinary Journal of Science and Technology, 5(4), 726-730.
12. Davronovna, A. K., Yusupovna, A. Z., & Olimboyevna, I. M. (2025). ADDRESSING COMMON CHALLENGES IN CHEMISTRY COURSES FOR MEDICAL STUDENTS AND ENSURING ACTIVE STUDENT PARTICIPATION. Multidisciplinary Journal of Science and Technology, 5(4), 719-725.
13. Абдуллаева, Х. Д., Атаджонова, З. Ю., & Ибодуллаева, М. О. (2025). BLENDED AND DIGITAL TEACHING STRATEGIES FOR CHEMISTRY IN MEDICAL EDUCATION. Multidisciplinary Journal of Science and Technology, 5(4), 714-718.
14. Davronovna, A. H. (2025, January). MEDICINAL SUBSTANCES CONTAINING BIOMETALS AND THEIR APPLICATION IN MEDICINE. In Next Scientists Conferences (pp. 87-89).
15. Davronovna, A. H. (2025). MEDICINAL SUBSTANCES CONTAINING BIOMETALS AND THEIR APPLICATION IN MEDICINE. Next Scientists Conferences, 1 (01), 87–89.
16. Sadullayev, S. E., Matyakubova, O. U., Artikov, I. A., Nurllayev, R. R., Ibadullayeva, S. S., & Yakubov, K. Y. (2023). RESULTS OF STUDIES ON THE LEVEL OF POPULATION

KNOWLEDGE ABOUT PARASITIC DISEASES AND ITS PREVENTION. Western European Journal of Medicine and Medical Science, 1(4), 15-20.

17. Nurllayev R. R., Ibadullayeva S. S., Yoqubov Q. Y. KICHIK QON AYLANISH DOIRASI ARTERIYALARINING MORFOLOGIK TUZILISHI //Научный Фокус. – 2023. – Т. 1. – №. 8. – С. 463-468.
18. Ibadullayeva, S. S., Yakubov, K. Y., Artikov, I. A., Nurllayev, R. R., & Sadullayev, S. E. (2023). CHARACTERISTICS OF PATHOMORPHOLOGICAL CHANGES IN LYMPHOCYTIC LEUKOSIS IN CHILDREN. Western European Journal of Medicine and Medical Science, 1(4), 21-26.
19. ANALYSIS OF THE EPIDEMIOLOGICAL FEATURES OF DIARRHEAL DISEASES IN CHILDREN IN THE SOUTHERN ARAL REGION. (2024). Multidisciplinary Journal of Science and Technology, 4(2), 345-351. https://www.mjstjournal.com/index.php/mjst/article/view/870
20. Yusupov , S., Sadullayev , S., & Yoqubov , Q. (2025). GEPATITLAR FONIDA KORONAVIRUS INFEKSIYASINING KECHISHI. Journal of Science-Innovative Research in Uzbekistan, 3(3), 294–303. Retrieved from https://inlibrary.uz/index.php/journal-science-innovative/article/view/76633
21. Yusupov , S., Sadullayev , S., Yoqubov , Q., & Ibragimov , U. (2025). HOMILADOR AYOLLAR ORASIDAGI QIZILCHA: XUSUSIYATLARI, XAVFLARI VA OLDINI OLISH. Journal of Science-Innovative Research in Uzbekistan, 3(3), 286–293. Retrieved from https://inlibrary.uz/index.php/journal-science-innovative/article/view/76634
22. SH, Y. S., YOQUBOV, Q., & NURLLAYEV, R. (2025). FEATURES OF THE COURSE OF THE HERPES TYPE 4 VIRUS IN CHILDREN. Multidisciplinary Journal of Science and Technology, 5(3), 397-402.
23. NURLLAYEV, R., SH, Y. S., & YOQUBOV, Q. (2025). PECULIARITIES OF THE ETIOLOGICAL STRUCTURE OF ACUTE DIARRHEAL DISEASES IN THE CONDITIONS OF THE SOUTHERN ARAL REGION. Multidisciplinary Journal of Science and Technology, 5(3), 403-408.
24. Saliybayevna, I. S., Yodgor o‘g‘li, Y. Q., & Odilbekovna, R. D. (2025). EPIDEMIOLOGY AND COMPLICATIONS OF COVID-19. Multidisciplinary Journal of Science and Technology, 5(3), 218-224.
25. Saliybayevna, I. S., Yodgor o‘g‘li, Y. Q., & Odilbekovna, R. D. (2025). NOSOCOMIAL PNEUMONIA AND ITS ETIOLOGICAL FACTORS. Multidisciplinary Journal of Science and Technology, 5(3), 225-230.
26. Razzokberganova, D. O., Ibragimov, S. J., Yoqubov, Q. Y., & Atadjanova, O. N. (2025). NUTRITION IN VIRAL HEPATITIS. Multidisciplinary Journal of Science and Technology, 5(3), 59-62.
27. Ibragimov, S. J., Razzoqberganova, D. O., Yoqubov, Q. Y., & Atadjanova, O. N. (2025). EMERGENCIES IN INFECTIOUS DISEASES: EPIDEMIOLOGICAL, MEDICAL AND SOCIAL ASPECTS. Multidisciplinary Journal of Science and Technology, 5(3), 63-67.
28. Razzokberganova, D. O., Ibragimov, S. J., Bobojonov, Y. B., & Yoqubov, Q. Y. (2025). EPIDEMIOLOGICAL AND CLINICAL CHARACTERISTICS OF ENTEROBIASIS. Multidisciplinary Journal of Science and Technology, 5(2), 665-668.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-5

29. Yoqubov, Q. Y., Bobojonov, Y. B., Razzokberganova, D. O., & Ibragimov, S. J. (2025). FEATURES OF THE COURSE AND SPREAD OF TENIARINCHOSIS DISEASE. Multidisciplinary Journal of Science and Technology, 5(2), 669-672.
30. Masharipov, S. M., Matyakubova, O. U., & Yakubov, K. Y. (2025). IMMUNE STATUS IN PATIENTS WITH PARASITIC DISEASES IN KHOREZM REGION. Multidisciplinary Journal of Science and Technology, 5(1), 514-517.