ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

THE IMPORTANCE OF VARIOUS FEEDING METHODS IN INCREASING MEAT PRODUCTIVITY OF GOLISTHENIZED BULLS.

Davronov Baxtiyor Xasanovich Termez State University of Engineering and Agrotechnology, Department of Zooengineering, Veterinary Medicine and Sericulture q.x.f.f.d (PhD) v.b dotsent
baxtiyordavronov95@gmail.com **https://orcid.org/0009-0009-4781-8313**
Rajabov Baxtiyor Musulmonovich Termez State University of Engineering and Agrotechnology, Department of Zooengineering, Veterinary Medicine and Sericulture
baxtiyorrajabov7079@gmail.com https://orcid.org/0009-0000-1417- 3875
Jo‘raboyev Bunyod Sherzod o‘g‘li Student of Termez State University of Engineering and Agrotechnology, Department of Zooengineering, Veterinary Medicine and Sericulture

Keywords: bullfinch, meat, productivity, live Weight, Food, Food unit, daily growth, feed ration, growth rate, digestible protein, blue and succulent foods.

Annotation: it is the correct Organization of feeding animals, taking into account the period of growth. Structured feed ration is harmonized in all respects, animals are given more blue and juicy foods. This is because these foods they are rich in digestible protein, are lightly digested and absorbed by the body. When young bulls are given blue and juicy feeds at the norm level, they ensure the growth of animals durkun, do not suffer from various diseases, the main thing is that less feed is spent per unit of product.

Relevance of the subject; the animal organism is always influenced by various environmental factors. Including natural climate, environmental condition, radiation, living conditions (storage method, placement density, group size, building microclimate, etc.) will be related to feeding level and type, feed preparation method for consumption, biological full value of ration content, etc

Introduction: the program for the development of the livestock sector and its sectors in the Republic of Uzbekistan for 2022 — 2026 (next-the program) is aimed at setting priority goals and objectives for the rapid development of the livestock sector and its sectors, sustainable provision of the population of the Republic with food products and expansion of production opportunities. Decision of the president of the Republic of Uzbekistan on measures to introduce advanced digital technologies into the agricultural sector, n PQ-257, date of adoption 02.08.2023, date of entry into force 03.08.2023. Due to the increase in the production of livestock products in the country, it is being organized to ensure food security, to widely introduce modern methods of production, to create a value-added chain, to develop cooperative relations, to provide state support for the livestock sector and its industries, and to effectively use the achievements of modern information and communication technologies and science in this Cattle breeding has long been practiced in intensive conditions and grazing on the territory of present-day Uzbekistan. Before the end of the 19th century, cammahsul, jussasi Maida jaydari and zebusimon cattle made up the bulk of cattle. Since the 20s of the 20th century, work has been carried out in the Republic on the formation of a group of productive herds based on the interbreeding of local and zebusimon black-cattle with cultural black, red-and Schwyz breeds, the establishment of reproductive farms of breeding purebred cattle, the development of State and cooperative cattle dairy farms and the establishment of dairy complexes, Dairy, dairy, meat and breeding farms of cattle were established. as a result, by the 80s, cattle breeding reached 98-99%. Herds of breeding cows giving 3800-4100 kg of milk were erected. The weight of bakers bred in meat

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

production complexes increased to 400-420 kg at the age of 12-13 months. Intensive breeding and stocking of cattle at Inter-farm cattle farms established in the districts gave positive results. Advanced experience and achievements of Science played an important role in the development of the practice of cattle breeding.

Comments on: Y.G.Abovyan (1990) geneticist kelib kelib tried to distinguish Keskin, 3 tribes of Buzoklar azratib, among them 20 tribes in general composition in reasonable numbers. Analyzing the results, we can conclude that barcha usish is a round (3,6,12,20 oylika) Kara-ola Bilan of the Holstein breed, katishma Balan of the Holstein breed Caucasian Kungur, Kara-ola and Caucasian Kungur Bilan of the Holstein breed chatishtirish as a result of olingan chatishma animal husbandry generally has a moderate odor. In particular, 20 individuals of cuttlefish of the Holstein breed had 77.7 g/l, Caucasian kungir bred 8.9 g/l (12.9%), cuttlefish 2.7 g/l (3.4%) and Caucasian kungir of the Holstein breed bred 5.3 g/l (7.3%). ketishgan. In this verse, Allah Almighty reported that all animals in the womb have the same size. On the territory of the Republic of Uzbekistan, each breed belongs to a clinic, a hematologist, a cadet, and a larynx living in the belgilar foidal. bilan baglikligin Z.T. 1986), V. V. I. Lenin (1986). Sh.Ballasov (1999), H.S.Mukhiddinova (2001), U.Kochchiev (2006), Sh. Karimov (2006), A.Boltaev (2007), A.A.Khushkavatov (2007), A.Kakharov (2008), R.Khayitov (2008) and bashkalar sharganishgan. A.Jehish (1990) noted each genotype of the Kara-Ola tanalarin from 18 tribes of the Jadalard bakib, among them the kursatkichlarini sharganib, the pure-bred breed of the Kara-Ola relative to the ozorbayjon zebus and the Indian zebus bilan olingan hybrid tanalard oxidative and Tiklanish zhara jadal kechganligin. 12 individuals were decorated with olingan duragailar channeling with erythrocyte content relative to 3.7-5.0% of the olage surface. Hemoglobin still has a distinct odor. This is the difference between shundai and shundai. N.Khushkavatov (1998), B.A.Kakhramanov (1999), A.T. (2006), including in Kazan. Yes.Karpova and bashkalar (2005) Saratov region "Abodimovsky" in conditions of keeping in each product a typical Simmental breed of karamollar morphological composition of tukislab sharganib, hemoglobin content in day-cigirlard meat 11.26 g%, day 0.23 (2.1%) and meat containing 0.42 g% (3.9%) of ecanligin. The number of red blood cells varies from 0.25 million/m3 (3.6%) to 0.43 million/mm3 (6.4%). At that time, when they couldn't eat meat, they couldn't eat meat. Day meat belongs to a strong Constitution, while Uzbekistan and Keskin's climatic conditions are changing relatively quickly. They are an organ consisting of a barge, a net and a hull, as well as shunting in the form of a boyd's elementaries. Bu informar S.Lumbunov and bashkalarning (1998). I.Skorkina and bashkalarning (2005) yesishicha fustning biologists in relation to amino acids deganda, undaga almashadigan and almashmaidigan. During the study, scientists analyzed all genotypes of chatishma (simmental x Kyzyl-ola holstein) containing the amino acid uzun musculin. Shunin Natizha kursatganki tyrosine, as in Bali, leucine, phenylalin, histidine, as lisa, kab arganin kyp bulgan uzun bulsa, according to Holstein sciences, as in chatishma xayvonlarning orka musculida amino acids ko 7/8, 3/4, with kuprok bulgan sciences, aksincha ko as their eco saklanar xayvonlarda meti. The total content of amino acids in pure form in simmental animal breeds is 3/4 and 7/8%, respectively, 0.83 and 2.10 mg, respectively. Yukori darajada Asil the adjective kursatkichi (0.79) also literally corresponds to the word 7/8 of chatishma bukachalardy kuzatilgan. Ushba was informed by G. Kalinin, S. Dolgochaev (2000), S.Karimov and Bashkalarning (2006) of the results of scientific research conducted by the Keladi Institute. In the Belgorod Republic, it is very important to eat meat, salmon meat and praise its quality, as well as ensure that meat consumption is balanced. In this verse, Allah Almighty said:

Table 1

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

Bibliographic Institute of Ukraine (co-authored)

Specification	Groups		
	I	II	III
Duration of experience, day	630	630	630
Living weight at the beginning of the experiment, kg	30,0±0,3	32,5±0,5	32,9±0,4
Living weight at the end of the experiment, kg	468,3±4,8	510,1±5,4	528,4±3,8
Extra live weight gained during the experiment, kg	438,3±4,1	477,6±4,7	495,5±3,9
Average one-day growth, g	696,0±5,8	758,0±5,1	787,0±4,9
Total food spent during the experiment, in the food unit	4012,2	4039,6	4109,5
Feed spent for 1 kg of extra live weight, in the food unit	9,15	8,46	8,29

The average weight of 630 rabbits in group I is 438.3 kg, and the average weight in tyrik is olishgan. Bu kursatkich II and III-groups with a relative body weight of 38.9 kg ($p<0.05$) or 8.8% and 55.8 kg ($p<0.01$) by 12.7% less. Srtacha Mutlak usish II is a group of animals, III is a group of tengkurlarig about 16.9 kg ($p<0.05$) or 3.5% lower than the norm of erishgan content. Usish tezligi, i.e. bir kunlik usish has 89 grams or 12.8 percent and 27 grams or 3.6 percent, respectively, as the III group. Group I, and group II, the last difference was 62 grams, or 7.5 percent. The experience of animal breeding is mainly due to the presence of phenotypic factors that influence their phenotypic behavior, and, in turn, an animal breeding experiment was organized during the period of the Olgan Hold Act. At the same time, each of them must be provided with nutrients and nutrients, as well as the nutrients contained in the food. It contains a variety of nutrients that are beneficial for digestion and the digestive tract. Don't think about it.A.Akmalkhonov (1984, 1993), Z.T.Turkulov (1985), P.S.Sobirov (1990), M.E.Ashirov (1994), I.M.Maksudov (1994), A.K.Kakharovlar (1994, 2005, 2007) the results of scientific research are mainly bildirishgan. The authors note that when growing dog food and dairy products, the feed consumption rate decreases, animals do not have enough space in the diet, while the main food products consume significantly less feed.

Conclusion: This is a kilib, and tadkikotlarimizda is a Kazakh view, important things, to shoot uchun yukori bulgan kyshimcha weight 1 kg weight, to shoot Holstein sailgan bukachalar gurularida sarflangan kamayishi kuzatib food, beer, food, as well as when shooting hususiyatlari bilan koplash weight, Ularda yahshilanganligi kuzatilgan. It took time to raise cattle, as the livestock population needed the product to feed their cattle with Galdag rice. Here is the riddle of ijobi etchimina topishda verb-atvor, in general, ethology has both practical and theoretical significance. At the same time, as with other animals, there are many different kinds of animals in this world, such as sheep, sheep, sheep, sheep, sheep, etc.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

LIST OF USED REFERENCES

1. Mirziyoyev S. M. Decree of the President of the Republic of Uzbekistan “On amendments to the Code of the Republic of Uzbekistan on Administrative Offenses” GI Kazakhstan pk-4576 dated January 29, 2020.

2. Mirziyoyev S. M. Decree of the President of the Republic of Uzbekistan “On amendments to the Law of the Republic of Uzbekistan” On amendments to certain legislative acts of the Republic of Uzbekistan "On Amendments to the Law of the Republic of Uzbekistan" On amendments to certain legislative acts of the Republic of Uzbekistan " No. gi Kazakhstan pk-5017 2021-03-March 2021.

3. Abovyan Y. G. immunobiochemical parameters of natural resistance of calves // J. "Zootechnia". 1991, No. 1. pp. 19-21.

4. Jehish Ahmed. Growth, development and meat productivity of hybrid young black-and-white breeds of different genotypes in the hot climate of Uzbekistan. The abstract. Dissertation Candidate of Art History, Agricultural Sciences. Tashkent, 1990. p. 24.

5. Karpova O., Anasimova E. Adaptive approach to the use of simmentals in the Volga region. "dairy and beef cattle breeding". 2004. No. 1. pp. 17-18.

6. Skorkina I., Negreeva A., Khlupov A. meat quality of bulls of different genotypes J."dairy and beef cattle breeding".- 2005. No. 5. pp. 14-15.

7. Karimov Sh., Davronova N., Kakharov A. K. Biological assessment of the quality of young meat of dairy and combined breeds of cattle: Scientific Journal of the Moscow Medical Center. I. M. Sechenov Academy. Moscow, 2006. pp. 242-244.

8. Zhavoronkov O. Z. and Bosch. European journal of Molekular & Klinical Medicine, 2020, Volume 7, Issue 9, Pages 1959-1965.