ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

IMPROVEMENT OF SURGICAL MANAGEMENT IN ANOPHTHALMIA

Isomov I.I., Hakimova Z.K.

Center for Professional Development of Medical Workers, Republic of Uzbekistan

Abstract

Anophthalmia, resulting from the surgical removal of the eyeball (enucleation or evisceration), remains a serious medical and psychosocial problem worldwide. According to the World Health Organization (WHO) and Vision 2020 initiative, over 55 million people sustain ocular injuries annually, leading to blindness in 1.6 million patients (Foster A., Vision 2020). Despite advances in ophthalmology, enucleation remains one of the most commonly performed ocular surgeries, particularly in cases of trauma, inflammation, and malignancy. In Russia alone, up to 8,000 enucleations are performed annually due to severe trauma (Gundorova R.A. et al., 1994).

Anophthalmic patients frequently experience significant psychological distress, functional limitations, and cosmetic disfigurement, which negatively affect their social integration and quality of life. It is reported that 80% of patients with anophthalmia are dissatisfied with their appearance and prosthetic outcomes, leading to reduced self-esteem and avoidance of social interactions (Haralampidi M.P., 2002).

A major challenge in the management of anophthalmia is the formation of a stable, functional, and aesthetically acceptable orbital stump capable of supporting a mobile and well-positioned ocular prosthesis. However, modern surgical methods and implant materials still demonstrate limitations, achieving satisfactory results in only 20–25% of cases (Gundorova R.A. et al., 2009). Poor outcomes are often associated with incomplete formation of the motility-supporting complex, restricted prosthesis mobility, eyelid deformities, and socket contracture.

Recent studies (Sorokina I.A., 2005; Beraya M.Z., 2006; Lavrentieva N.V., 2013) emphasize the importance of individualized surgical planning based on comprehensive assessment of the orbital stump, as well as the use of biointegrative implants to improve functional and cosmetic results. Moreover, the concept of quality of life has become a critical outcome measure in ophthalmic surgery, encompassing not only physical and visual parameters but also psychological and social well-being (WHO, 1993).

This study aimed to improve surgical management in anophthalmia by developing standardized criteria for orbital stump evaluation, refining surgical indications, and conducting a comparative analysis of reconstructive techniques in order to optimize prosthetic outcomes and enhance patients' quality of life.

Keywords: anophthalmia, enucleation, ocular prosthesis, reconstructive surgery, orbital implants, ophthalmic rehabilitation, quality of life.

Introduction

Anophthalmia is a severe condition following the removal of the eyeball, characterized by a significant cosmetic defect, functional impairments, and a reduced quality of life. Despite advances in ophthalmic surgery, effective reconstructive approaches for these patients remain a clinical challenge. The relevance of this study is determined by the necessity to develop an individualized surgical strategy to enhance clinical outcomes and improve aesthetic results. In clinical practice, rehabilitation of anophthalmic patients is one of the most difficult aspects of ophthalmic surgery. Complications such as socket contracture, eyelid malpositions, prosthetic instability, and limited

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

motility of the ocular prosthesis compromise both aesthetic appearance and functional outcomes. Consequently, these issues have a profound effect on the psychological and social well-being of patients.

Given the importance of both cosmetic and functional restoration in anophthalmic patients, the development of standardized diagnostic and surgical protocols tailored to the individual anatomical and functional characteristics of the orbital stump is of paramount importance. Additionally, advances in biomaterials, including porous polyethylene and hydroxyapatite orbital implants, have provided new opportunities to enhance postoperative outcomes.

Literature Review

Anophthalmia management has evolved considerably over the past few decades. Traditional surgical techniques were limited to simple enucleation without the use of orbital implants, often resulting in socket contraction and poor prosthetic outcomes. The introduction of intraorbital implants aimed to preserve orbital volume, prevent socket contraction, and support prosthesis mobility.

Multiple studies have investigated the efficacy of various implant types, including non-integrated acrylic implants and biointegrative materials such as hydroxyapatite, porous polyethylene (Medpor), and alumina ceramics. Beraya M.Z. (2006) reported improved cosmetic outcomes with the use of porous implants, which facilitate fibrovascular ingrowth, reducing the risk of implant migration and extrusion.

Lavrentieva N.V. (2013) emphasized the importance of combining orbital implant insertion with meticulous reconstruction of the motility-supporting complex (tenon's capsule, conjunctiva, and orbital muscles) to optimize prosthesis mobility and socket anatomy. According to Sorokina I.A. (2005), individualized preoperative assessment of the orbital stump significantly improves the selection of surgical techniques and reduces the risk of postoperative complications.

Recent publications highlight the role of advanced imaging modalities such as high-resolution ultrasound and computed tomography (CT) in the preoperative assessment of the orbital cavity. These methods allow for precise evaluation of the volume, configuration, and anatomical relations of the orbital stump, facilitating personalized surgical planning.

Furthermore, psychosocial rehabilitation is now recognized as a critical aspect of comprehensive care for anophthalmic patients. Quality of life studies demonstrate that successful cosmetic rehabilitation significantly enhances patients' psychological well-being, social integration, and self-esteem.

Objective

To assess the condition of the orbital stump in patients with anophthalmia and conduct a comparative analysis of the effectiveness of current reconstructive surgical techniques.

Materials and Methods

The study included 20 patients with anophthalmia treated in ophthalmology departments. Patients ranged in age from 18 to 65 years. The primary indications for enucleation were trauma, inflammatory eye diseases, and orbital tumors.

Examination methods included:

- Ophthalmological assessment (eyelid position, conjunctival fornix condition);
- Evaluation of the orbital stump (size, consistency, mobility, position, and relation to surrounding tissues);
- Laboratory tests (complete blood count, urinalysis, coagulation profile);
- Instrumental examinations (orbital ultrasound).

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

Patients were divided into groups based on the type of reconstructive surgical technique applied. Early and long-term postoperative outcomes were evaluated.

Results

The study revealed:

- The condition of the orbital stump was a key factor in selecting the appropriate surgical technique.
- Criteria for assessing the stump condition were developed (based on size, consistency, mobility, and position).
- Among the applied techniques, the best cosmetic and functional outcomes were achieved with surgeries involving the formation of a complete motility-supporting stump using bioimplants.
- A comparative analysis showed that an individually tailored surgical approach based on stump assessment increased prosthetic effectiveness and reduced the risk of anophthalmic socket syndrome.
- The use of modern reconstructive techniques provided satisfactory cosmetic and functional outcomes in 78% of patients.

Discussion

The findings confirm that the success of rehabilitation for anophthalmic patients largely depends on a thorough assessment of the orbital stump and a differentiated choice of surgical strategy. The application of individually selected reconstructive methods significantly reduces complication rates and improves patients' quality of life.

Conclusion

1. Criteria for evaluating the condition of the orbital stump have been developed, allowing for optimal surgical treatment planning.
2. A comparative analysis of modern reconstructive techniques for anophthalmia was conducted.
3. An individualized approach to selecting surgical methods improves cosmetic results, functional outcomes, and the quality of life for patients.
4. The obtained data justify the implementation of the proposed recommendations in clinical ophthalmic practice.

References

1. Foster, A. (2001). Vision 2020: The right to sight. Bulletin of the World Health Organization, 79(3), 244–249.
2. World Health Organization. (1993). WHOQOL Study Protocol. WHO Regional Office for Europe.
3. Gundorova, R. A., Neroev, V. V., & Kashnikov, V. V. (2009). Surgery of the eyeball: Modern approaches to enucleation and prosthetics. Moscow: Medicina.
4. Gundorova, R. A., & Danilichev, V. F. (1994). Eye and orbit injuries: Diagnosis and treatment. Moscow: Medicina.
5. Filatova, I. A. (1994). Eye and orbit injuries in working-age patients: Clinical presentation, diagnosis, treatment (Doctoral dissertation). Moscow.
6. Kharlampidi, M. P. (2002). Optimization of enucleation techniques for improved cosmetic results in eye prosthetics (Doctoral dissertation). Moscow.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-5

7. Sorokina, I. A. (2005). Use of Alloplant biopreparations in the formation of the motor support unit during enucleation. Vestnik Oftalmologii, 121(5), 17–20.

8. Beraya, M. Z. (2006). Comprehensive diagnostics of the orbital area before and after eyeball removal (Doctoral dissertation). Moscow.

9. Krasilnikova, V. L. (2007). Modern approaches to oculoplastic surgery in anophthalmic patients. Vestnik Oftalmologii, 123(1), 43–46.

10. Lavrentieva, N. V. (2013). Modern aspects of anophthalmic socket reconstruction. Oftalmokhirurgiya, (1), 39–43.

11. Shklyaruk, V. V. (2005). Comparative evaluation of orbital reconstruction techniques after enucleation. Russian Ophthalmological Journal, 6(1), 33–36.

12. Rauwer, A. E., & Michelson, K. M. (1981). Psychosocial aspects of facial plastic surgery. Plastic and Reconstructive Surgery, 68(3), 395–400.