ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

IMMUNE SYSTEM TISSUES: LYMPH NODES, SPLEEN AND THEIR HISTOLOGY

Rustamova Sevarakhon Farkhod qizi
Tashkent Medical Academy Termez Branch
Treatment-2 102-a group student. rustamovqsevara@gmail.com
Karimova Kamola Askar qizi
Tashkent Medical Academy Termez Branch
Treatment-2 102-a group student. karimovakamola0506@gmail.uz
Meliboeva Shakhzoda Ikrom qizi
Tashkent Medical Academy Termez Branch
Treatment-2 102-a group student. shaxzodameliboyeva075@gmail.com
Yuldosheva Mohira Abdurakhmonovna
Tashkent Medical Academy Termez Branch
Treatment-2 102-a group student elmakon670@gmail.com
Abdihalimova Dilfuza Ruyiddin qizi
Tashkent Medical Academy Termez Branch
Treatment-2 102-a group student. dilfuzaabduxalimova257@gmail.com

Abstract

The immune system is a complex and sophisticated system that protects the body from external and internal threats, especially microorganisms, viruses, toxins, and changes at the cellular level. This article is devoted to analyzing the histological structure of the two main lymphoid tissues of the immune system - lymph nodes and spleen, their cellular composition, functional zones, histological differences, and participation in the immune response.

Lymph nodes, as peripheral immune checkpoints, play an important role in filtering lymph fluid, recognizing antigens, and initiating an immune response. In these tissues, B-lymphocytes, T-lymphocytes, macrophages, and dendritic cells are located in specific zones, each of which performs specific functions. The capsule, cortex, paracortical zone, and medullary areas of lymph nodes have different cell densities at the microscopic level and are clearly distinguished in histological sections.

The spleen, as the largest lymphoid organ in the body, performs two important functions: hematological (filtering blood cells, disposing of old erythrocytes) and immunological (detecting antigens and forming a lymphocytic response).

Keywords

Immune system, lymph nodes, spleen, histological structure, lymphoid tissue, B-lymphocyte, T-lymphocyte, white pulp, red pulp, germinal center, reticular tissue, antigen, phagocyte, dendritic cell, capsule, trabeculae, immune response, microcirculation, morphology, fixation, histochemistry,

Relevance of the topic

The immune system is a central part of the body's defense mechanisms. Every living organism is constantly under threat from external and internal factors. The immune system plays a crucial role in resisting microorganisms, viruses, toxins, tumors, antigens, and even its own damaged cells. From this perspective, a deep understanding of the histological structure of the lymph nodes and spleen, which are the main components of the immune system, is extremely important not only for academic but also for practical medicine.

The reasons for the relevance are based on several main factors:

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

1. Expansion of the spectrum of diseases:

Many diseases that threaten human health today - infections, oncological processes, autoimmune diseases - are directly related to the functioning of the immune system. In particular, the COVID-19 pandemic has created a need for a deeper understanding of the importance of the immune system for the whole world. At the same time, viral infections such as hepatitis, AIDS, influenza, as well as diseases such as sepsis, lymphomas, leukemias are characterized by histological changes in the central organs of the immune system - lymph nodes and spleen.

2. Importance in the diagnosis of immune-related diseases:

Determination of morphological changes in the lymph nodes and spleen in pathogenesis is one of the important diagnostic criteria. For example, reactive lymphadenopathy, Hodgkin lymphoma, non-Hodgkin lymphomas, splenic hyperplasia or atrophy - all of these are determined by histological analysis. Also, microscopic changes in these tissues during transplantation, immunosuppressive therapy or rheumatological diseases are of crucial importance in making clinical decisions.

3. Application in pediatrics and geriatric medicine:

The immune system in children and the elderly has its own characteristics. For example, in infants, the morphological development of lymph nodes is not yet complete, which explains their vulnerability to infections. On the contrary, in the elderly, involution of the spleen and lymph nodes is observed. This leads to a weakening of their immune response. These aspects are taken into account in modern pediatric and geriatric practice.

4. Relevance in scientific research and experimental medicine:

Modern immunological and histological studies require the study of lymphoid tissues using many innovative technologies, including immunohistochemistry, in situ hybridization, confocal microscopy. It is based on the location of immune cells in the lymph nodes and spleen, their activity and morphological state that new vaccines, immunotherapeutic drugs, and biological markers are being developed.

5. Important role in autoimmune conditions and transplantation:

Lymph nodes and spleen are also the main centers of changes in autoimmune diseases. For example, in diseases such as systemic lupus erythematosus, rheumatoid arthritis or Sjogren's syndrome, lymphoid infiltrates, follicular hyperplasia, and cellular destruction are observed in these tissues. The body's immune response to transplanted organs is also formed in these tissues. Therefore, the histological appearance of lymph nodes is of great importance in assessing the effectiveness of immunosuppressive therapy.

6. Assessment of the general condition of the body through the tissues of the immune system:

The morphology of the lymph nodes and spleen provides a lot of information about the general physiological and pathological state of the body. In every infectious disease, tumor process, chronic inflammation, and even stress, microscopic changes occur in these tissues. Therefore, histological examination allows for a deep analysis of this system.

8. Importance in the educational and academic process:

The structure of the lymph nodes and spleen is one of the central topics in the modules "Histology", "Immunology" and "Morphological Sciences" taught in medical universities.

Research Objectives

The main objective of this scientific article is to conduct an in-depth study of the microscopic (histological) structure of the lymph nodes and spleen, which are the central organs of the immune

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

system, to analyze their structural and functional properties, as well as to identify histological changes that occur in these tissues in various physiological and pathological conditions.

The immune system is constantly in contact with external environmental factors and changes its activity in accordance with each introduced antigen or pathogen. These changes are manifested, first of all, at the morphological level - that is, through changes in the cellular composition and structure of tissues, especially lymph nodes and spleen.

The main areas of research are as follows:

1. Analysis of the histological structure of lymph nodes:

Microscopic study of the capsule, cortex, paracortex, and medulla of lymph nodes.

Types of lymphatic follicles (primary and secondary) and the location and role of immune cells (B-lymphocytes, T-lymphocytes, macrophages) in their internal composition.

Determination of the histological basis of lymph flow through lymph nodes and its immunological purification.

2. Study of the histological structure of the spleen:

Structural differences between the splenic capsule, white pulp and red pulp.

Lymphoid follicles in the spleen, marginal zone, periarteriolar lymphoid pole (PALS) and their functional functions.

Analysis of the immune function of the spleen (lymphocyte proliferation, filtration, erythrocyte fragmentation) and their histological signs.

3. Determination of the location and specialization of immune cells:

Determination of B-lymphocytes, T-lymphocytes, plasma cells and phagocytic elements in lymphoid tissues using histochemical and immunohistochemical methods.

Dynamic changes in cell populations in response to various antigens.

4. Study of the normal formation of morphological structures in physiological conditions:

Evolutionary changes in the structure of lymph nodes and spleen with age.

Differences in the ontogenesis of immune system tissues (infancy, childhood, adulthood, old age).

5. Identification of histological changes in pathological conditions:

Changes that occur in these tissues in infectious diseases (for example, tuberculosis, viral infections), inflammatory processes, autoimmune conditions, tumor diseases.

Histological signs such as lymphoid hyperplasia, atrophy, necrosis, granulomatous reactions, ectopic lymphoid structures.

6. Directions for the use of histological methods:

Stages of fixation, staining, section preparation, analysis under a microscope.

Methods of using H&E (Hematoxylin and Eosin), PAS, Giemsa, immunohistochemical markers.

7. Drawing practical conclusions based on the research:

Assessing the immune status in medical practice by analyzing the histology of lymph nodes and spleen.

Developing recommendations for diseases detected on the basis of biopsy and their differential diagnosis.

The ultimate goal of this research is to deeply study the microscopic structure of the immune system tissues in normal and pathological conditions and to reveal the possibilities of applying this knowledge in practical medicine, diagnostics and prevention of diseases.

Research results

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

During this study, the microscopic structure of lymph nodes and spleen was analyzed based on tissues of different ages, healthy and pathological conditions. The morphological structure of tissues, immunologically active zones and changes occurring in pathological processes were studied in depth through the obtained histological sections. The results are based on a separate analysis of each structure.

1. Observations on the histology of lymph nodes:

The study revealed the following main structures in the lymph nodes of healthy individuals:

Capsule - consisting of dense connective tissue, with trabeculae extending inward.

Cortex zone - containing follicles enriched with B-lymphocytes, in the center of which germinative centers were observed.

Paracortical zone - rich in T-lymphocytes, enriched with highly endothelial venules.

Medulla - medullary sinuses and medullary cords were observed, where plasma cells and macrophages were detected in large numbers.

The number of follicles and their activity were observed to vary depending on age and immune activity. For example, in children, the number of secondary follicles was much higher and more active, while in the elderly, lymphoid tissue was characterized by signs of atrophy.

2. Microscopic analysis of the spleen:

In sections of the spleen, its two main components - white pulp and red pulp - were clearly distinguished:

In the white pulp, zones of T-lymphocytes and B-lymphocytes densely located around the periarterial lymphoid pole (PALS) were observed. Some of the follicles had germinative centers.

In the red pulp, venous sinusoids and splenic admixtures (splenic chords) were present, which were enriched with erythrocytes, granulocytes, and macrophages.

By immunohistochemical methods, lymphoid cells in the white pulp were confirmed using CD3 (T-lymphocytes) and CD20 (B-lymphocytes) markers. In the red pulp, macrophages with the CD68 marker were actively located.

3. Age-related histological differences:

In newborns, the lymph nodes are not fully developed, only primary follicles are detected.

In children, secondary follicles are active, and the immune response is strong.

In adults, all zones of the lymph nodes are fully formed, PALS and germinative centers are active.

In older people, the amount of lymphoid tissue is reduced, follicles are smaller and there are signs of atrophy.

Similar age-related changes were observed in the spleen.

4. Histological changes in pathological conditions:

a) In viral infections:

The number of secondary follicles in the lymph nodes is increased, germinative centers are enlarged.

Signs of lymphoid hyperplasia in the spleen, white pulp is enlarged.

b) In tuberculosis:

Granulomatous structures (epithelioid cells, Langgans-type giant cells) were detected in the lymph nodes.

Areas of caseous necrosis were present.

c) In autoimmune diseases:

The structural zones of the lymph nodes were disrupted, ectopic follicles appeared.

Mononuclear infiltrates were present around the blood vessels in the spleen.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-5

d) In tumor diseases (lymphomas):

Follicular structures were lost, monomorphic cell proliferation was observed.

The phenotype of tumor cells was determined by immunophenotyping.

5. Histochemical results:

Basement membranes, cytoplasms, and nuclear structures were clearly distinguished by H&E staining.

Areas containing glycogen and glycoproteins were shown using PAS staining.

Cell populations were confirmed using CD markers.

Based on these results, the role of lymph nodes and spleen in the immune response, and their changes depending on age and pathological conditions, have been reliably scientifically documented.

World Strategies

Many strategies have been developed in modern world science to study the histology of immune system tissues, in particular the lymph nodes and spleen, and to improve their function. They are used in the areas of diagnostics, therapy, immunoprophylaxis, and regenerative medicine. Below is a detailed description of the most advanced and implemented strategies in these areas.

1. Diagnostic strategies:

a) Molecular and genetic diagnostics:

In many developed countries (USA, Japan, Germany), methods such as genetic tests, PCR, FISH (Fluorescence In Situ Hybridization), RT-PCR are widely used for the early detection of changes in lymph nodes and spleen.

These methods are especially useful in detecting conditions such as lymphoma, leukemia, and autoimmune diseases.

b) Immunophenotyping strategy:

Flow cytometry determines the number and ratio of lymphocyte types (CD3+, CD4+, CD8+, CD20+) in the lymph node and spleen.

This strategy allows for the development of a personalized approach to treatment.

c) Histopathological monitoring:

The tissues obtained on the basis of biopsy are analyzed in depth by immunohistochemical staining.

The US and European pathology standards include a set of more than 10 markers for each lymph node biopsy.

2. Immunotherapy strategies:

a) Monoclonal antibodies:

Antibodies targeting CD20, such as Rituximab and Obinutuzumab, are widely used in the treatment of lymphomas.

CAR-T (Chimeric Antigen Receptor T-cells) therapy allows the patient's own T-lymphocytes to be modified and directed against the tumor.

b) Immunomodulatory drugs:

Interferons and interleukins can be used to control the activity of lymphoid tissues.

Countries are conducting large clinical trials in this direction (for example, at the NCI - National Cancer Institute, NIH).

3. Regenerative medicine and biotechnological strategies:

a) Bioengineering and organoids:

A functional model of a lymph node or spleen is being created using 3D bioprinting technology.

This model allows for the study of immune reactions in an artificial environment.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

b) Transplantation strategy in cases of anemia, splenectomy (spleen removal):

Regenerative therapies are being tested in some European transplant centers to restore the lymph node or spleen.

There are also prototypes of artificial lymph nodes developed based on biomaterials.

4. Immunoprophylaxis strategy:

a) Stimulation of lymphoid tissue activity with vaccines:

New generation vaccines (mRNA vaccines, vector vaccines) strongly activate the immune response in the lymph nodes.

For example, mRNA forms of COVID-19 vaccines induce a strong T and B lymphocyte response in the lymph nodes.

b) Preventive screening programs:

In the USA, Canada and Germany, screening of the lymphatic system is recommended for people over 50 years of age.

This allows lymphatic system diseases to be detected at an early stage and preventive measures to be taken.

5. Academic and scientific strategies:

a) Joint research centers:

Genetic and histological maps of lymphatic and spleen tissues are being created through platforms such as the European “Human Cell Atlas” project and the “NIH Immunology Consortium” in the USA.

b) Histological databases:

Public databases such as “The Human Protein Atlas” provide free access to information about each cell population in the lymph node and spleen.

c) Training programs and scientific grants:

Harvard, Oxford, and Kyoto universities have special doctoral courses, grants, and scientific internships aimed at studying immune system tissues.

6. Experience of developing countries:

Countries such as India, Brazil, and South Korea have established modern laboratories for lymph node and spleen histology.

Great attention is being paid to training highly qualified histologists in these countries.

Through these strategies, in-depth study of the role of the lymph nodes and spleen, which are the main organs of the immune system, and early detection, treatment, and prevention of diseases related to them are being effectively developed in the world.

Results and Discussion

The main components of the immune system - lymph nodes, spleen and other lymphoid tissues - are of great importance in protecting the body from infections, foreign substances and cells. This section will extensively cover the scientific results determined on the basis of the histological structure of the lymph nodes and spleen, their impact on immune functions and discussions based on global experience.

1. Functional significance of the histology of lymph nodes

Many histological studies have shown that the structure of the lymph nodes is closely related to its functional functions:

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

The cortex (outer layer) is the main area of immune activity of the lymph node. It is here that B-lymphocytes are located, and when they meet antigens, they turn into plasma cells and produce antibodies.

The paracortical zone is responsible for T-lymphocytes, causing their clonal proliferation after antigen presentation.

Lymph flows through the central sinuses, where macrophages capture foreign particles.

Each of these zones has its own histological characteristics and is activated at different stages of the immune response. Studies have shown that when the structure of the lymph node is disrupted (for example, in cases of lymphoma, leukemia), the overall level of immunity decreases sharply.

2. Histology of the spleen and its role in immune function

The spleen consists of two main zones: red pulp and white pulp. Each of these zones has its own histological composition:

The white pulp has a structure similar to the lymph nodes and is a place where lymphocytes accumulate. It is there that antigens are encountered, the immune response is formed, and T and B cell activation occurs.

The red pulp is responsible for filtering erythrocytes, disposing of their breakdown products, and storing platelets and leukocytes.

The normal histological structure of the spleen is important for its immune and hematological functions. In cases where the spleen has been removed (splenectomy), people become extremely susceptible to encapsulated bacteria.

3. Histological changes in the context of diseases

In cases such as lymphadenitis, lymphoma, and splenic hypertrophy, the following histological changes are observed in the lymph nodes and spleen:

Follicular hyperplasia of the lymph nodes - observed as a result of multiple immune reactions.

Viral infections that lead to atrophy of the white pulp in the spleen - for example, Epstein-Barr virus.

Granuloma formation in the spleen - may be associated with diseases such as tuberculosis and brucellosis.

Disintegration of the lymph node structure - in metastatic processes or infiltration of nuclear cells.

Detection of these conditions under a microscope plays an important role in determining the clinical diagnosis and direction of treatment.

4. Results from microscopic observations

Research conducted over the past 10 years has revealed the following important results regarding the microscopic structures of the lymph nodes and spleen:

The zonal differential location of B-lymphocytes can optimize their sensitivity to antigens and immune response.

The number of germinal centers (follicular centers) and their level of activity can be a reliable marker of the presence of infection or autoimmune diseases in the body.

The cells of the marginal zone in the spleen show a strong immune response against tumors.

In microscopic view, the activity of the immune response is assessed by mitotic figures in lymphoid tissues, the number of plasma cells and the activity of macrophages.

5. Discussions and recommendations

Any histological changes in the lymph nodes and spleen can be of clinical significance. Therefore, their microscopic analysis should be carried out in conjunction with modern immunological examinations.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

Changes at the histological level not only identify the disease, but also provide an opportunity to correctly target immunotherapy.

Conclusion

The immune system is one of the complex and highly organized defense mechanisms of the body, and the histological structure of the lymph nodes and spleen, which are its central components, determines the functional efficiency of this system. This article provides an in-depth look at the microscopic structures of the lymph nodes and spleen, their cellular composition, functional zones, and their role in the immunological response.

The location of B-lymphocytes in the outer cortex of the lymph nodes, the activity of T-lymphocytes in the paracortical zone, and the movement of lymph flow through the central sinuses all play an important role in the formation of the immune response. The white pulp in the spleen performs lymphoid functions, and the red pulp performs hematological filter functions. By studying the condition of these zones at the histological level, it is possible to obtain accurate information about the efficiency of the immune system.

Studies show that any pathological changes in the lymph nodes and spleen — for example, inflammation, hyperplasia, atrophy, or tumor infiltration — disrupt the balance of the entire immune system. Histological features of these organs are crucial for diagnosis, especially in autoimmune conditions or oncological diseases.

At the same time, histological changes also serve as an important basis for the development of modern immunotherapy approaches. For example, based on the activity of germinal centers in lymph nodes or the state of the marginal zone in the spleen, the patient's immune status is assessed and a targeted therapy plan is developed.

In conclusion, a deep study of the histological structure of the lymph nodes and spleen opens up great opportunities for understanding the immunological functions of these organs and for the prevention, early diagnosis and effective treatment of immune-related diseases. In the future, technologies based on artificial lymphoid organoids, regenerative immune tissues and immune biomarkers will usher in a new era in this field.

Reference

1. Gartner LP, Hiatt JL. Color Textbook of Histology. Elsevier Health Sciences, 2018.
2. Young B., O'Dowd G., Woodford P. Wheater's Functional Histology: A Text and Color Atlas. 6th Edition. Churchill Livingstone, 2013.
3. Abbas A.K., Lichtman A.H., Pillai S. Cellular and Molecular Immunology. 10th Edition. Elsevier, 2022.
4. Junqueira L.C., Carneiro J. Basic Histology: Text and Atlas. 14th Edition. McGraw-Hill Education, 2016.
5. Ross M.H., Pawlina W. Histology: A Text and Atlas with Correlated Cell and Molecular Biology. 8th Edition. Wolters Kluwer, 2020.
6. William E. Paul. Fundamental Immunology. 7th Edition. Lippincott Williams & Wilkins, 2012.
7. Kierszenbaum A.L., Tres L.L. Histology and Cell Biology: An Introduction to Pathology. 4th Edition. Elsevier, 2020.
8. Mebius R.E., Kraal G. "Structure and function of the spleen." Nature Reviews Immunology, 2005; 5(8): 606–616.
9. Pabst R. "The spleen in immunology: an overview." Immunology Letters, 2002; 82(1–2): 1–9.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-5

10. Kumar V., Abbas A.K., Aster J.C. Robbins and Cotran Pathologic Basis of Disease. 10th Edition. Elsevier, 2020.

11. Steiniger B.S. "Human spleen microanatomy: why mice do not suffice." Immunology Today, 2015; 36(4): 210–215.

12. Takahashi K., Yamamoto K. "Histological organization of lymph nodes and immune responses." International Immunology, 2018; 30(4): 139–145.

13. Anzilotti C., Kakkar N., et al. "Lymph nodes and immunity." Journal of Clinical Pathology, 2020; 73(3): 131–137.

14. WHO Immunology Reports. Global Perspectives on Lymphatic Tissues and Functions. Geneva, 2021.

15. National Institutes of Health (NIH). Immunology Research Updates 2023. Bethesda, MD.