ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

AIR POLLUTION IN SAMARKAND CITY DURING 1995–2010 AND ITS ANALYSIS

Toshturdiyev Nurbek Nurali ugli
National University of Uzbekistan named after Mirzo Ulugbek
Faculty of Physics, Department of Hydrometeorology
3rd year student
Phone: +998 88 910 42 46
Email: nurbektoshturdiyev86@gmail.com

Abstract: This article analyzes the level of air pollution in Samarkand city during the period from 1995 to 2010. Based on monitoring data, the annual variation dynamics of concentrations of suspended particulates, sulfur dioxide, carbon monoxide, nitrogen oxides (NO and NO_2), phenol, hydrogen fluoride, and ammonia were examined. The results of the study indicate a decrease in the levels of some pollutants during the given period, while others remained relatively stable. The main sources of atmospheric pollution were identified as industrial activities, transport vehicles, and municipal waste. The article also briefly analyzes the impact of this pollution on public health.

Keywords: Samarkand city, air quality, pollution, suspended particulates, SO_2, CO, NO_2, NO, environmental monitoring, health risk.

Environmental pollution is one of the most pressing ecological problems of the modern era. In particular, the quality of atmospheric air directly affects human health and living conditions. An increase in the concentration of harmful substances in the air can lead to respiratory diseases, cardiovascular diseases, and many other adverse health effects. Therefore, regular monitoring of the state of atmospheric air, identification of existing pollutants, and study of their dynamics are of great importance. Samarkand city is one of the ancient, large, and developing centers of industry, transport, and tourism in Uzbekistan. In this region, industrial enterprises, automobile transport, and communal waste contribute to the emission of various harmful substances into the atmosphere. The population growth and rapid urbanization have further increased the ecological burden.

This article analyzes the annual changes in the main pollutants in the atmospheric air of Samarkand city (suspended particles, sulfur dioxide, carbon monoxide, nitrogen oxides, phenol, hydrogen fluoride, and ammonia) during the period 1995–2010 based on statistical data. The aim of the study is to determine how the air quality changed over these years and to highlight the main factors influencing this condition.

Pollutants in the atmosphere enter the air from both natural and anthropogenic sources. Their presence negatively impacts air quality, posing risks to human health, flora and fauna, and the overall ecological balance. According to monitoring results conducted in Samarkand, the air quality depends primarily on the following pollutants:

Suspended particles (dust) are solid particles that remain suspended in the air for a long time and are mainly produced by industrial activities, vehicle traffic, and soil lifted by the wind. They penetrate the human respiratory tract and harm health. Sulfur dioxide (SO_2) is released during the combustion of fuels, especially coal and petroleum products. It causes acid rain, which damages plants and human health. Carbon monoxide (CO) forms due to incomplete fuel combustion and mainly originates from transport vehicles. This gas binds with hemoglobin, limiting oxygen delivery to tissues. Nitrogen dioxide (NO_2) and nitrogen oxide (NO) are nitrogen oxides emitted during fuel

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

combustion, especially from vehicles and industrial enterprises. They adversely affect the human respiratory system and contribute to the formation of photochemical smog. Phenol is a harmful organic compound produced in chemical manufacturing, characterized by a sharp odor and toxic properties. Phenol can irritate the respiratory tract, skin, and eyes. Hydrogen fluoride (HF) is released in some industrial sectors, particularly metallurgy and chemical industries. It is highly toxic and causes severe harm to living organisms. Ammonia (NH_3) mainly arises from agriculture, manure, and waste decomposition. In certain concentrations, it irritates the respiratory tract.

In this study, we analyze the values of these pollutant gases and compounds in Samarkand city over the period 1995–2010.

Table 1

Air Pollution in Samarkand City Atmosphere for 1995–2010 (Average Values, mg/m³)

No	Years	Suspended Particles (mg/m³)	Sulfur Dioxide (SO_2) (mg/m³)	Carbon Monoxide (CO) (mg/m³)	Nitrogen Dioxide (NO_2) (mg/m³)	Nitric Oxide (NO) (mg/m³)	Phenol (mg/m³)	Hydrogen Fluoride (HF) (mg/m³)	Ammonia (NH_3) (mg/m³)
1	1995	0.2	0.005	2	0.03	0.01	0.002	0.007	0.04
2	1996	0.2	0.005	3	0.02	0.01	0.002	0.008	0.03
3	1997	0.2	0.007	2	0.02	0.01	0.001	0.006	0.02
4	1998	0.2	0.007	2	0.01	0.01	0.002	0.005	0.02
5	1999	0.2	0.007	1	0.01	0.01	0.001	0.003	0.02
6	2000	0.2	0.005	1	0.01	0.01	0.002	0.004	0.02
7	2001	0.2	0.004	1	0.02	0.01	0.002	0.004	0.04
8	2002	0.1	0.004	1	0.01	0.01	0.002	0.004	0.02
9	2003	0.1	0.005	1	0.01	0.01	0.002	0.003	0.03
10	2004	-	0.005	1	0.01	0.01	0.002	0.003	0.04
11	2005	-	0.005	1	0.01	0.01	0.001	0.005	0.03
12	2006	-	0.006	1	0.01	0.01	0.002	0.005	0.03
13	2007	-	0.006	1	0.01	0.01	0.002	0.003	0.02
14	2008	-	0.005	1	0.01	0.01	0.002	0.002	0.02
15	2009	-	0.005	1	0.01	0.01	0.002	0.002	0.02
16	2010	-	0.006	1	0.01	0.01	0.002	0.003	0.03

Table 2

Air Pollution in Samarkand City for 1995–2010 (Annual Amount in Thousand Tons)

No	Years	Suspended Particles (mg/m³)	Sulfur Dioxide (SO_2) (mg/m³)	Carbon Monoxide (CO) (mg/m³)	Nitrogen Dioxide (NO_2) (mg/m³)	Nitric Oxide (NO) (mg/m³)
1	1995	1.77	0.473	3.548	0.825	0.064
2	1996	1.174	0.876	3.141	0.303	0.506

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

3	1997	0.904	0.482	2.465	0.230	0.371
4	1998	1.195	0.529	2.141	0.209	0.347
5	1999	1.587	0.855	2.524	0.317	0.566
6	2000	1.895	0.714	0.243	0.281	0.425
7	2001	-	-	-	-	-
8	2002	1.818	0.636	2.133	0.261	0.386
9	2003	1.399	0.408	2.193	0.508	0.067
10	2004	-	0.434	1.75	0.458	0.098
11	2005	-	0.434	1.466	0.381	0.098
12	2006	-	0.835	1.177	0.237	0.142
13	2007	-	0.658	0.907	0.197	0.055
14	2008	-	-	-	-	-
15	2009	-	-	-	-	-
16	2010	-	-	-	-	-

Let us now focus on the changes in each of the above-mentioned pollutant gases over the years:

Suspended particles (dust). From 1995 to 2003, the concentration of suspended particles in the air remained around 0.2 mg/m³, with a noticeable decrease to 0.1 mg/m³ starting from 2002. From 2004 to 2010, there are no available data, which indicates either a suspension of monitoring activities or uncertainty in the measurements. The reduction in these particles may indicate strengthened control over dust emissions from industrial waste and vehicles.




Figure 1. Fluctuation graph of sulfur dioxide in Samarkand city during 1995–2010

Sulfur dioxide (SO_2). From 1995 to 2003, the concentration of this gas ranged between 0.004 and 0.007 mg/m³, while during 2004 to 2010, this indicator remained mostly around 0.005 to 0.006 mg/m³. This stability suggests a consistent level of pollution, which may indicate a relatively high pollution level, poor fuel quality, or insufficient filtration systems.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6




Figure 2. Fluctuation graph of carbon monoxide in Samarkand city during 1995–2010

Carbon monoxide (CO). In 1995, the concentration of this gas was 2 mg/m³, increasing to 3 mg/m³ in 1996. In the following years, it stabilized at around 1 mg/m³. The high levels in the early years could be associated with a large number of vehicles and the use of outdated engines, while the subsequent decrease may indicate improved monitoring and better vehicle maintenance practices.




Figure 3. Fluctuation graph of nitrogen dioxide in Samarkand city during 1995–2010

Nitrogen dioxide (NO_2). In 1995, the concentration of this gas was 0.03 mg/m³, while during 1996–2003 it remained within the range of 0.01–0.02 mg/m³. From 2004 to 2010, the level stabilized at 0.01 mg/m³. Since nitrogen dioxide mainly originates from vehicle emissions, the low fluctuations may be linked to relatively low traffic volumes or the renewal of the vehicle fleet.Nitric oxide (NO). Throughout all the observed years, the concentration of this compound remained stable at 0.01 mg/m³. Given the minimal variation, it can be concluded that the pollution level was constant but still present. Phenol. Between 1995 and 2010, the concentration of phenol remained around 0.001–0.002 mg/m³. Although this is not a high level, in certain years (such as 1997, 1999, and 2005), it dropped to as low as 0.001 mg/m³. This fluctuation may be related to the operation levels of chemical industry enterprises.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6




Figure 4. Fluctuation graph of hydrogen fluoride in Samarkand city during 1995–2010

Hydrogen fluoride (HF). In 1995, the concentration of this substance was 0.007 mg/m³, but it gradually decreased to a range of 0.002–0.005 mg/m³ by 2010. This indicates improved control over the use of fluorides in industrial processes.

Ammonia (NH_3). The level of this gas was 0.04 mg/m³ in 1995 and fluctuated between 0.02–0.04 mg/m³ in the following years. Ammonia primarily originates from agriculture and organic waste. The downward trend in its concentration may be associated with more environmentally conscious approaches in agricultural practices.

Based on observations and statistical data from 1995 to 2010, it is evident that air pollution levels in Samarkand city have varied over time. While the concentration of certain pollutants decreased significantly in specific years, others showed stable or only slightly variable trends. For instance, particulate matter (suspended particles) was relatively high in the early years but then declined. Similarly, carbon monoxide and nitrogen dioxide levels were initially high but later stabilized. Other substances such as sulfur dioxide, phenol, hydrogen fluoride, and ammonia displayed fluctuating trends throughout the period. This suggests that air pollution in the city is influenced by constant sources such as transportation, industrial activity, population density, and meteorological conditions.

Based on the findings above, the following recommendations are proposed:

1. Strengthen the continuous air quality monitoring system, particularly to address the data gap for suspended particles since 2004.
2. Enhance environmental control of vehicles operating in the city, and reduce the number of cars that do not meet ecological standards.
3. Introduce modern waste treatment technologies in industrial enterprises, and regularly inspect existing purification equipment.
4. Raise public awareness of environmental issues and promote knowledge about the health risks of air pollution.
5. Expand green spaces, tree planting, and maintenance efforts to support natural air purification processes.
6. Align local pollutant concentration standards with international norms, and incorporate global best practices into local implementation.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

References:

1. The Center of the Hydrometeorological Service of the Republic of Uzbekistan (Uzhydromet) (1995–2010). Annual air pollution monitoring data. Tashkent.
2. Environmental Protection Agency of the Republic of Uzbekistan. (2023). Annual air quality report. Tashkent.
3. State Committee for Ecology. (2023). Monitoring results of air pollution. Tashkent.
4. Ministry of Health. (2023). Statistics on respiratory diseases. Tashkent.
5. Qodirov, A., & Nurmatov, D. (2018). *Ekologiya va atrof-muhit muhofazasi asoslari*. Toshkent: Fan va texnologiya.
6. Ergashev, B., & To‘rayev, M. (2020). *O‘zbekiston ekologiyasi: muammolar va istiqbollar*. Samarqand: Zarafshon nashriyoti.
7. Toxirov, M. A. (2019). *Atmosfera havosining ifloslanishi va uni monitoring qilish usullari*. Toshkent: Ekosan.
8. World Health Organization (WHO). (2022). *Air pollution and child health: prescribing clean air*. Geneva: WHO Publications.
9. United Nations Environment Programme (UNEP). (2021). *Air Quality Policies and Strategies in Central Asia*. Nairobi: UNEP.