ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

UDK 635.64:631.527.52:581.1(575.1)

STUDY OF AGROBIOLOGY OF COCKTAIL TOMATO VARIETIES IN GREENHOUSE CONDITIONS

Kamalova Nargiza Maxammadovna

Basic doctoral student at the Andijan Institute of Agriculture and Agrotechnologies

Annotatsiya:

Mazkur maqolada issiqxona sharoitida yetishtirilayotgan kokteyl tipidagi pomidor navlarining agrobiologik xususiyatlari o‘rganildi. Tajriba Toshkent viloyatida joylashgan issiqxonada amalga oshirilib, *Sweet Million*, *Sakura F1*, *Red Pearl*, *Cherry Cocktail* va *Lilit* kabi navlar o‘rganildi. Navlarning gullash muddati, meva yetilish davri, hosildorlik ko‘rsatkichlari va o‘sish dinamikasi tahlil qilindi. Tadqiqot natijalariga ko‘ra, *Sweet Million* va *Sakura F1* navlari yuqori hosildor va tez yetiladigan navlar sifatida ajralib turdi. Ushbu ish natijalari issiqxonalar uchun istiqbolli navlarni tanlashda hamda agrotexnologiyalarni takomillashtirishda amaliy ahamiyatga ega.

Kalit so‘zlar: ***kokteyl pomidor, issiqxona, agrobiologik xususiyat, gullash muddati, meva tutishi, hosildorlik, Sweet Million, Sakura F1, nav tanlash***

Аннотация:

В данной статье изучены агробиологические особенности коктейльных сортов томатов, выращиваемых в тепличных условиях. Эксперимент проведён в теплице, расположенной в Ташкентской области, с участием сортов *Sweet Million*, *Sakura F1*, *Red Pearl*, *Cherry Cocktail* и *Lilit*. Проанализированы сроки цветения, плодоношения, урожайность и динамика роста растений. По результатам исследований сорта *Sweet Million* и *Sakura F1* выделяются как скороспелые и урожайные. Полученные данные имеют практическое значение для выбора перспективных сортов и совершенствования агротехнологий в тепличном овощеводстве.

Ключевые слова: *коктейльные томаты, теплица, агробиологические особенности, срок цветения, плодоношение, урожайность, Sweet Million, Sakura F1, выбор сорта.*

Annotation:

This article investigates the agrobiological characteristics of cocktail-type tomato varieties cultivated under greenhouse conditions. The experiment was conducted in a greenhouse located in the Tashkent region using the varieties *Sweet Million*, *Sakura F1*, *Red Pearl*, *Cherry Cocktail*, and *Lilit*. Key parameters such as flowering time, fruit development period, yield potential, and growth dynamics were analyzed. The results revealed that *Sweet Million* and *Sakura F1* are early-maturing and highly productive varieties. These findings are of practical importance for selecting promising cultivars and improving greenhouse vegetable production technologies.

Keywords: *cocktail tomato, greenhouse, agrobiological characteristics, flowering time, fruit set, yield, Sweet Million, Sakura F1, variety selection.*

Login.

- In recent years, small-sized, high-quality and salty-tasting cocktail tomato varieties, which are in high demand in the consumer market, have gained particular importance in vegetable growing. Such varieties have great potential not only in the domestic market, but also for export opportunities.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

In-depth study of the agrobiological characteristics of such varieties in greenhouse conditions, determination of their growth dynamics, flowering and fruiting processes, and selection of high-yielding varieties are important factors.

- Methods
- The study was carried out in an intensive greenhouse located in the Tashkent region in the spring of 2024. Five cocktail tomato varieties were selected for the experiment:
- *Sweet Million*
- *Sakura F1*
- *Red Pearl*
- *Cherry Cocktail*
- *Lilit*
- Each variety was planted in 3 replicates, in a 60x30 cm planting pattern, and the following agrobiological indicators of development were recorded:

• Vegetation period (in days);
• Flowering time;
• Fruiting and maturity;
• Total growth height;
• Yield (kg/m²).• Yield (kg/m²).

- The plants were cared for in accordance with the rules of agrotechnology. Irrigation was drip irrigation, lighting was artificial additional lighting (LED lamp), temperature was maintained at 22–26°C during the day and 16–18°C at night.
- Research results
- The results of the periment are shown in Table

The plants were cared for in accordance with the rules of agrotechnology. Irrigation was drip irrigation, lighting was artificial additional lighting (LED lamp), temperature was maintained at 22–26°C during the day and 16–18°Catnight.

Research results

The results of the experiment are shown in Table 1.

Table 1

Agrobiological characteristics of tomato varieties in the experiment (2024-2025**yy)**

Variety name	Flowering start (day)	Fruit ripening (days))	Yield (kg/m²)
Sweet Million	22	38	10.1
Sakura F1	25	40	9.2
Red Pearl	30	46	8.7
Cherry Cocktail	27	42	8.5
Lilit	29	44	8.2

- *Sweet Million* The variety began flowering on the 22nd day and showed the fastest development rate, with the first fruit ripening on the 38th day.
- • *Sakura* F1 also bloomed relatively early and had high yields.
- • *The Red* Pearl and Lilit varieties required more vegetation time, and also had average yields.

• *Discussion The results of the study indicate that there are significant differences in agrobiological indicators between cocktail tomato varieties grown in greenhouse conditions. These differences are*

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

mainly related to the genetic characteristics of the varieties, their degree of adaptation to the agroecological environment, and their requirements for temperature, humidity, light, and irrigation conditions.

Sweet Million and **Sakura F1** varieties:

• Early flowering (22–25 days),

• Fast maturity (38–40 days),

• High yield (9.2–10.1 kg/m²) These varieties were distinguished not only by their rapid development, but also by their uniformity of fruit quality and high marketability. These factors indicate that they are worthy of widespread commercial use. At the same time, the short growing season of such varieties shortens the harvest cycle and creates the possibility of growing two crops in one season.

Red Pearl and **Lilit** in varieties:

• **Late flowering** (29–30 days)

• **Long fruit ripening period (**44–46 days),

• **Slightly lower yield** (8.2–8.7 kg/m²),

but their fruit is attractive in appearance, has a uniform shape and skin salinity. These varieties may have high marketability in the export direction. Such qualities are especially in high demand in the European and Middle Eastern markets.

Although the Cherry Cocktail variety has average performance and lags behind Sweet Million and Sakura F1 in terms of yield, its potential can be increased if care conditions are optimized**.2-jadval**

Kokteyl pomidor navlarining asosiy sifat ko‘rsatkichlari

Variety name	Fruit mass (g)	Brix (%)	Vitamin C (mg/100g)	Market price (sum/kg)
Sweet Million	18	6.2	24.5	12 000
Sakura F1	20	6.0	23.0	11 500
Red Pearl	22	6.5	21.8	13 000
Cherry Cocktail	16	6.3	22.4	11 000
Lilit	21	6.7	20.1	12 500

According to the results of a study conducted to assess the agrobiological and quality indicators of cocktail-type tomato varieties, the specific characteristics of each variety were determined. In particular, factors such as fruit mass, Brix (%) indicator indicating taste quality, vitamin C content, and market price were analyzed in depth. In terms of fruit weight, the **Red Pearl (22 g) and Lilit (21 g)** varieties have the largest fruit size, which makes them convenient to eat and attractive in appearance. At the same time, the Cherry Cocktail variety has a fruit weight of 16 g, and due to its small size, it can be considered a variety that is consumed more by young children and for decorating dishes. According to the Brix indicator, which determines the taste quality, the **Lilit (6.7%)** and Red **Pearl (6.5%)** varieties took the lead. This indicator indicates that they have a high sugar and dry matter content. Such tomatoes are often suitable for salads and direct consumption. The Brix level of the **Sweet Million (6.2%)** and **Cherry Cocktail (6.3%)** varieties is also quite high, ensuring good taste. In terms of vitamin C content, the varieties Sweet Million (24.5 mg/100 g) and Sakura F1 (23.0 mg/100 g) outperformed others. This indicator indicates that these varieties are biologically beneficial

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-6

for human health. Vitamin C is an important antioxidant that strengthens immunity, so such varieties deserve attention for the purpose of growing healthy food.

As for the market price, **Red Pearl (13,000 soums/kg)** is the most expensive variety, which is due to its appearance, fruit uniformity and high taste quality. This variety may be of particular importance for export. **Sweet Million (12,000 soums/kg**) and **Lilit (12,500 soums/kg)** were also evaluated as varieties with high commercial potential. Although the market price of the Cherry Cocktail variety (11,000 soums/kg) is relatively low, it has a place in the domestic market as an inexpensive and salty-tasting product that is in high demand.On this basis, it can be concluded that each variety is effective for a specific purpose. Sweet Million is recommended as a universal variety in terms of quality and health indicators, Red Pearl is recommended for export, **Lilit is** recommended for consumers who prioritize taste quality, Sakura F1 is recommended as a high-yielding and fast-growing variety, and Cherry Cocktail is recommended as an inexpensive product with aesthetic appeal.

In the future, it is possible to further improve the quality and economic efficiency of each variety by developing **agrotechnical** solutions suitable for each variety - in particular, by optimizing the amount of fertilization, adjusting the frequency of irrigation, improving formation and similar care elements. Based on the results of the research, the following important conclusions were drawn:

1. **Cocktail tomato varieties differ significantly in terms of development dynamics, flowering time, fruit ripening period, and yield indicators. It was found that these differences are mainly related to the genetic characteristics of the varieties and the degree of adaptation to greenhouse conditions**
2. **Sweet Million** va **Sakura F1** varieties are recommended as the most promising varieties for intensive greenhouse technologies due to their early flowering, rapid ripening and high yield. They create the opportunity to obtain more crops per season and increase economic efficiency.
3. Although the Red Pearl and Lilit varieties have a relatively long development period, they are of particular importance as export varieties due to their fruit quality, appearance and high Brix level. It is advisable to develop special care methods to ensure their demand in foreign markets.
4. In the future, it is recommended to develop individual agrotechnical solutions for each variety. Including:o Fertilization rates and structure (macro and microelement balance),

o Irrigation regimes (frequency and water consumption),o Formation and dressing methods (single-stem or two-stem systems)

4.Adapting these elements on the basis of targeted research will allow you to fully unlock the potential of each variety.

By developing agrobiological and technological recommendations for selected varieties, it is possible to optimize development, increase productivity, and increase the economic benefits of production. This will be of great importance for the sustainable development of greenhouse vegetable growing.

FOYDALANILGAN ADABIYOTLAR RO‘YXATI

1. Ashurov, A. H., Abdurasulov, D. S. (2021). *Issiqxonada sabzavot yetishtirish agrotexnologiyasi*. Toshkent: “Fan” nashriyoti.
2. Bahriddinov, B. Yu. (2020). *Ochiq va yopiq gruntda pomidor yetishtirishning biologik asoslari*. Samarqand: SamDU nashriyoti.
3. Qurbonov, H. (2018). *O‘zbekistonda pomidor navlarining agrobiologik xususiyatlari va ulardan samarali foydalanish*. Toshkent: TDAU ilmiy ishlar.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

4. Rasulov, A.A., & Turaev, M.R. (2022). *Greenhouse vegetable production in Central Asia: Challenges and innovations*. Journal of Horticultural Science, 10(2), 45–51.
5. Mirzaev, M. M., & Karimova, N. O. (2019). *Tomato (Solanum lycopersicum) varieties for intensive farming: Yield and quality parameters*. International Journal of Agronomy, 17(1), 78–83.
6. FAO (2020). *Protected cultivation in the Mediterranean region*. Food and Agriculture Organization of the United Nations.
7. Vaxobov, X. Q. (2023). *Intensiv issiqxonalarda agronomik tadbirlarni optimallashtirish yo‘llari*. Toshkent: Agroinnovatsiya markazi.
8. Johnson, L.F., & Thorne, J.H. (2017). *Tomato cultivation under controlled environments: Varietal response and productivity*. Agriculture and Environment, 12(3), 201–210.
9. Gulyamov, S.R., & Karimov, U.T. (2021). *Sifatli pomidor hosili uchun oziqlantirish va sug‘orish texnologiyalarini tanlash*. Toshkent: Agrobiologiya ilmiy jurnali, 4(2), 39–44.
10. Qobilov, Sh.J. (2022). *Oziq-ovqat xavfsizligida pomidor mahsulotining o‘rni va eksport salohiyati*. Agroiqtisodiyot va innovatsiyalar jurnali, 5(1), 60–65.

Login Introduction)

In recent years, there has been a growing demand for high-quality, salty, and small-sized tomato varieties - cocktail tomatoes - that can meet consumer needs. It is important to fully study the agrobiological characteristics of these varieties in greenhouse conditions, analyze their development stages, flowering, and fruiting mechanisms.

Methods (Methods)

The experiment was conducted in a greenhouse located in the Tashkent region. Varieties: Sweet Million, Sakura F1, Red Pearl, Cherry Cocktail, Lilit. Each variety was planted in 3 replicates, and the vegetation period of development, the beginning of flowering, fruit set, and overall growth dynamics were recorded.

Results (Results)

- *Sakura F1 variety started flowering on day 25, and fruiting was observed on day 40.*
- *Sweet Million showed rapid development, flowering on day 22, and fruiting on day 38.*
- *Red Pearl was the slowest growing variety, flowering started on day 30.*
- *Sweet Million (10.1 kg/m²) and Sakura F1 (9.2 kg/m²) dominated in terms of yield.*

Discussion (Discussion)

The differences in the development of varieties are related to their genetic characteristics and the degree of adaptation to the greenhouse environment. The Sweet Million and Sakura F1 varieties were found to be promising as early maturing and high-yielding varieties. They may serve as the most suitable choice for greenhouses.