ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

PHYSIOLOGICAL PROPERTIES OF BASIL (OCIMUM BASILICUM) PLANTS UNDER HYDROPONIC CONDITIONS.

Abdumalikova Zarnigor Komil qizi
Teacher at Tashkent Vocational and Medical College

Abstract:This study investigates the physiological characteristics of basil (Ocimum basilicum) grown under hydroponic conditions. Parameters such as growth rate, leaf surface area, chlorophyll content, transpiration rate, and photosynthetic efficiency were analyzed during the experiment. The results demonstrated that basil cultivated in a hydroponic system exhibited higher vegetative growth and improved biochemical indicators compared to those grown in conventional soil conditions. The findings confirm that hydroponic technology is an effective agronomic approach for the cultivation of medicinal and aromatic plants.

Keywords:Hydroponics, basil, Ocimum basilicum, physiological characteristics, photosynthesis, transpiration, chlorophyll, growth rate, agronomic technology.

Introduction

Basil (Ocimum basilicum L.) is one of the most widely cultivated aromatic plants, valued for its culinary, medicinal, and industrial applications. Rich in essential oils, phenolic compounds, and bioactive metabolites, basil plays a significant role in the pharmaceutical, food, and cosmetic industries. Traditionally, basil has been grown in soil-based systems; however, increasing demand for high-quality produce and the need for sustainable agricultural practices have stimulated interest in soilless cultivation methods, particularly hydroponics.

Hydroponic cultivation, which involves growing plants in nutrient-rich aqueous solutions without soil, offers several advantages, including precise control of nutrient supply, efficient water use, faster growth cycles, and reduced risk of soil-borne diseases. Previous studies have shown that hydroponic systems can enhance the growth performance and secondary metabolite content of various herbs. Nevertheless, the physiological responses of basil under hydroponic conditions, especially in relation to photosynthetic activity, water balance, and chlorophyll accumulation, require further investigation. Understanding the physiological adaptations of basil in hydroponics is crucial for optimizing growth conditions, improving yield, and maximizing the quality of bioactive compounds. This study aims to evaluate the growth parameters and physiological traits of O. basilicum grown in a controlled hydroponic environment and to compare the results with conventional cultivation methods.

Methodology

Plant Material and Experimental Design

The experiment was conducted using Ocimum basilicum L. seeds of a locally adapted cultivar. Seeds were germinated in inert rockwool cubes under controlled laboratory conditions and, after the emergence of two true leaves, seedlings were transferred to a deep-water culture (DWC) hydroponic system. The study was arranged in a completely randomized design with two treatments: hydroponic cultivation and conventional soil-based cultivation (control), each with three replicates.

Hydroponic System Setup

The hydroponic units consisted of opaque nutrient tanks (20 L) fitted with aeration pumps to maintain dissolved oxygen levels above 6 mg/L. Plants were supported using net pots and grown in a nutrient

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

solution formulated according to the Hoagland and Arnon standard composition, adjusted to pH 6.0–6.2 and an electrical conductivity (EC) of 1.6–1.8 mS/cm. The nutrient solution was replaced weekly to prevent nutrient imbalance.

Environmental Conditions

Both hydroponic and control groups were maintained under identical environmental conditions in a growth chamber: air temperature 24 ± 2 °C, relative humidity 60 ± 5%, and a 16 h photoperiod with a light intensity of 200 $\mu mol \cdot m^{-2} \cdot s^{-1}$ provided by LED lamps.

Physiological Measurements

Growth parameters such as plant height, number of leaves, and leaf surface area were measured weekly. Chlorophyll content was determined using a SPAD-502 chlorophyll meter, while photosynthetic rate and transpiration rate were measured with a LI-6400 portable photosynthesis system. Leaf relative water content (RWC) was calculated following the Barrs and Weatherley method.

Data Analysis

All measurements were statistically analyzed using one-way analysis of variance (ANOVA) followed by Tukey's post hoc test at a 95% confidence level ($p < 0.05$). Statistical analyses were performed using SPSS 26.0 software.

Results

The results indicated a significant difference between hydroponically grown basil and soil-grown controls in terms of growth and physiological parameters. After 30 days of cultivation, the hydroponic group exhibited a 28.5% greater plant height (average 36.8 ± 1.2 cm) compared to the control group (28.6 ± 1.5 cm). The leaf surface area was also significantly higher in hydroponic plants (172.3 ± 4.5 cm^2) than in soil-grown plants (136.8 ± 3.9 cm^2).

Chlorophyll content, as measured by SPAD values, averaged 46.2 ± 1.1 in hydroponic plants, compared to 39.4 ± 1.3 in controls, indicating enhanced photosynthetic pigment accumulation. Photosynthetic rate in the hydroponic group was 18.6 ± 0.7 $\mu mol\ CO_2 \cdot m^{-2} \cdot s^{-1}$, which was 21.4% higher than that of the control. Transpiration rates followed a similar trend, with hydroponic plants exhibiting 4.2 ± 0.2 $mmol\ H_2O \cdot m^{-2} \cdot s^{-1}$ versus 3.3 ± 0.1 $mmol\ H_2O \cdot m^{-2} \cdot s^{-1}$ in soil-grown plants.

Statistical analysis confirmed that all measured parameters showed significant differences between the two cultivation methods ($p < 0.05$).

Discussion

The results demonstrate that hydroponic cultivation significantly enhances the vegetative growth and physiological performance of Ocimum basilicum. The observed increase in plant height and leaf area is consistent with previous findings by Dorais et al. (2016), who reported that precise nutrient control in hydroponics stimulates rapid biomass accumulation. The higher chlorophyll content and photosynthetic rate in hydroponic basil may be attributed to improved nutrient availability and root-zone aeration, which optimize metabolic processes.

Furthermore, the elevated transpiration rates suggest more efficient water transport and stomatal regulation under hydroponic conditions, potentially leading to greater CO_2 assimilation efficiency. This aligns with the findings of AlShrouf (2017), who noted that hydroponic systems can improve both water-use efficiency and photosynthetic performance in leafy herbs.

From a commercial perspective, these results indicate that hydroponics not only accelerates growth but may also enhance the quality attributes of basil, such as pigment concentration and potentially secondary metabolite levels, which are important for culinary and medicinal uses.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

Conclusion

This study confirms that hydroponic cultivation has a positive impact on the physiological traits and growth performance of Ocimum basilicum. Compared to conventional soil cultivation, hydroponics produced plants with greater height, larger leaf area, higher chlorophyll content, and improved photosynthetic activity. These findings highlight the potential of hydroponic systems as an efficient and sustainable method for producing high-quality basil.

Future research should focus on optimizing nutrient formulations for maximum secondary metabolite yield and assessing the economic feasibility of large-scale hydroponic basil production.

REFERENCES:

1. Дораис М., Пападопулос А.П., Грауэрман А. Гидропонные системы и контроль роста растений // Canadian Journal of Plant Science. – 2016. – Т. 96, №3. – С. 356-370.
2. AlShrouf A. Hydroponics, aeroponic and aquaponic as compared with conventional farming // American Scientific Research Journal for Engineering, Technology, and Sciences. – 2017. – Vol. 27, №1. – P. 247-255.
3. Hoagland D.R., Arnon D.I. The water-culture method for growing plants without soil // California Agricultural Experiment Station Circular. – 1950. – №347. – P. 1–32.
4. Бобокулов Ф.Х., Хамроқулов М.М. Гидропоника технологиясининг илмий асослари ва қишлоқ хўжалигида қўлланилиши // Агробиология журнали. – 2020. – №2. – С. 45-51.
5. Маҳмудов Ш.Қ., Юсупова Н.М. Гидропоникадаги озуқа эритмалари таркиби ва уларнинг ўсимлик ўсишига таъсири // Ўзбекистон аграр фанлар хабарномаси. – 2021. – №4. – С. 78-84.
6. Resh H.M. Hydroponic Food Production: A Definitive Guidebook. – 7th ed. – Boca Raton: CRC Press, 2013. – 560 p.