ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

EFFECTS OF THYROID DYSFUNCTION ON METABOLIC PROCESSES

Kokand University, Andijan Branch Faculty of Medicine Department of Natural and Medical Sciences Assistant: **Sh.D. Ismoilova**

Student: **Sh.I. Qochqarova** (Group 24/04, General Medicine)

ANNOTATION

The thyroid gland is one of the primary endocrine organs responsible for regulating metabolic processes through the secretion of thyroxine (T4) and triiodothyronine (T3). These hormones play a critical role in controlling cellular energy production, protein synthesis, lipid metabolism, and carbohydrate turnover. According to the World Health Organization (WHO), thyroid disorders affect approximately **5–7%** of the global population, with hypothyroidism accounting for around 3.5% and hyperthyroidism for 1–2% of cases. Regional studies in Uzbekistan have reported a prevalence of **6–8%**, highlighting the public health significance of thyroid dysfunction in this area. This article examines the metabolic alterations resulting from thyroid dysfunction, including changes in basal metabolic rate, lipid profile, glucose homeostasis, and protein turnover. A review of scientific literature demonstrates that hypothyroidism slows down metabolism, leading to weight gain, hyperlipidemia, and reduced energy expenditure, whereas hyperthyroidism accelerates metabolism, causing weight loss, muscle wasting, and thermoregulatory disturbances. The findings underscore the importance of regular monitoring of thyroid function and early diagnosis to prevent complications related to metabolic imbalance.

Keywords: Thyroid dysfunction, metabolism, hypothyroidism, hyperthyroidism, thyroid hormones, metabolic rate, lipid metabolism, carbohydrate metabolism, endocrine system, hormonal imbalance

INTRODUCTION

The thyroid gland is one of the most critical components of the endocrine system and plays a central role in maintaining metabolic homeostasis. It synthesizes and secretes two primary hormones: thyroxine (T4) and triiodothyronine (T3), both of which regulate a wide spectrum of metabolic activities at the cellular level. These hormones influence oxygen consumption, heat production, glucose utilization, lipid metabolism, protein turnover, and even affect the function of nearly every organ system, including the heart, brain, liver, and muscles.

Thyroid dysfunction occurs when the gland produces either too little hormone (hypothyroidism) or too much (hyperthyroidism), disrupting the normal physiological balance. In hypothyroidism, the slowed production of thyroid hormones leads to a decrease in basal metabolic rate (BMR), weight gain, cold intolerance, bradycardia, fatigue, constipation, and depression. Biochemically, it is associated with increased serum cholesterol levels, impaired glucose tolerance, and reduced thermogenesis.

In contrast, hyperthyroidism results in an accelerated BMR, weight loss despite normal or increased appetite, heat intolerance, tachycardia, restlessness, anxiety, and frequent bowel movements. On the biochemical level, it leads to increased gluconeogenesis, enhanced lipolysis, decreased serum cholesterol, and increased nitrogen excretion due to elevated protein catabolism.

Both conditions can have profound effects on overall health and quality of life if left untreated. Thyroid disorders are relatively common worldwide and may be caused by autoimmune diseases (such as Hashimoto’s thyroiditis or Graves’ disease), iodine deficiency, congenital defects,

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

medication side effects, or tumors. Importantly, even subclinical thyroid dysfunction—where hormone levels are only slightly abnormal—can still significantly impact metabolic parameters and cardiovascular risk.

Given the essential role of the thyroid in regulating metabolism, understanding the physiological and biochemical consequences of its dysfunction is vital for early detection, accurate diagnosis, and effective treatment. This article aims to provide a comprehensive overview of how thyroid hormone imbalances affect metabolic processes, including insights into the underlying mechanisms, clinical implications, and potential therapeutic strategies.

Literature Review

Thyroid dysfunction and its impact on metabolism have been widely studied across clinical and experimental research. Numerous studies have confirmed that thyroid hormones, particularly T3, play a direct role in regulating basal metabolic rate (BMR), mitochondrial activity, and thermogenesis. According to Silva and Rabelo (1997), T3 increases the expression of mitochondrial uncoupling proteins, leading to enhanced heat production and energy expenditure.

In individuals with **hypothyroidism**, several researchers have observed a consistent reduction in BMR and lipid clearance. A study by Duntas and Brenta (2012) highlighted that hypothyroidism is frequently associated with dyslipidemia, including elevated LDL cholesterol and triglycerides, due to reduced LDL receptor activity and decreased hepatic lipase function. Additionally, hypothyroidism impairs glucose metabolism, often contributing to insulin resistance (Brenta, 2011).

Conversely, in **hyperthyroidism**, studies have demonstrated increased lipolysis, enhanced gluconeogenesis, and a hypercatabolic state. Bianco and McAninch (2013) showed that elevated thyroid hormone levels stimulate hepatic glucose output and increase muscle protein breakdown, leading to negative nitrogen balance and muscle wasting. This aligns with clinical findings of weight loss, increased appetite, and fatigue in patients with uncontrolled hyperthyroidism.

Autoimmune thyroid diseases—such as **Graves' disease** and **Hashimoto's thyroiditis**—are also of significant interest in the literature due to their chronic impact on metabolism. Research by Antonelli et al. (2015) emphasizes the metabolic alterations seen in autoimmune thyroid patients, including increased risk of metabolic syndrome, obesity, and cardiovascular disease.

Furthermore, **subclinical thyroid dysfunction**, where TSH is abnormal but T3/T4 levels remain within reference ranges, has been shown to influence metabolic parameters subtly but significantly. Rodondi et al. (2005) demonstrated that even subclinical hypothyroidism is associated with increased cholesterol levels and a higher risk of coronary heart disease.

Recent advances in molecular biology have also deepened our understanding of thyroid-metabolism interactions. Studies on thyroid hormone receptors and intracellular signaling pathways reveal that thyroid hormones regulate gene expression in hepatocytes, adipocytes, and myocytes, directly altering energy balance and metabolic efficiency (Yen, 2001).

Overall, the body of literature underscores the multifaceted and systemic nature of thyroid hormone action on metabolism. However, despite significant progress, further research is needed to understand inter-individual variability, long-term metabolic outcomes, and optimal treatment strategies in borderline or subclinical thyroid conditions.

Methodology

This article is based on a comprehensive review of current scientific literature related to thyroid gland dysfunction and its impact on metabolic processes. A systematic search was conducted using electronic databases, including **PubMed**, **ScienceDirect**, **Google Scholar**, and **Scopus**, to identify

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

relevant peer-reviewed articles published between **2000 and 2025**. Keywords used in the search included *"thyroid dysfunction"*, *"hypothyroidism"*, *"hyperthyroidism"*, *"metabolism"*, *"thyroid hormones"*, *"basal metabolic rate"*, and *"endocrine disorders"*.

Inclusion criteria consisted of:

Original research articles, reviews, and meta-analyses in English;

Studies involving human subjects or well-established animal models;

Publications with clear data on metabolic outcomes associated with thyroid dysfunction.

Exclusion criteria included:

Studies not available in full text;

Articles focusing exclusively on pediatric or rare genetic thyroid disorders;

Case reports without broader clinical relevance

A total of approximately **60 articles** were reviewed in depth, of which **30** were selected based on their methodological quality, relevance, and contribution to understanding the relationship between thyroid dysfunction and metabolism. Information extracted included study design, population characteristics, hormonal levels, metabolic parameters (e.g., lipid profile, glucose metabolism, BMR), and clinical outcomes.

The collected data were qualitatively analyzed and categorized into thematic sections: effects of hypothyroidism, effects of hyperthyroidism, autoimmune thyroid disease and metabolism, and subclinical thyroid dysfunction. This narrative review aimed to synthesize current evidence and highlight gaps in knowledge that require further investigation.

No experimental or clinical interventions were performed as part of this study, and no ethical approval was required, as the methodology relied solely on previously published data.

Results

A thorough review of the literature revealed consistent evidence that both hypothyroidism and hyperthyroidism significantly affect metabolic processes. Key findings include:

Basal Metabolic Rate (BMR): Hypothyroidism is associated with a marked decrease in BMR, while hyperthyroidism leads to its increase.

Lipid Metabolism: In hypothyroid patients, increased levels of total cholesterol and LDL cholesterol were consistently observed. In contrast, hyperthyroid states were linked to enhanced lipid breakdown and reduced serum cholesterol levels.

Carbohydrate Metabolism: Hypothyroidism often leads to impaired glucose tolerance and insulin resistance. Hyperthyroidism, meanwhile, increases hepatic glucose production and may lead to transient glucose intolerance.

Protein Metabolism: Hypothyroid individuals may exhibit reduced protein synthesis, while hyperthyroid patients often show increased protein catabolism and muscle wasting.

Body Weight: Most studies confirm weight gain in hypothyroidism and weight loss in hyperthyroidism, independent of dietary intake.

Thermoregulation: Cold intolerance in hypothyroidism and heat intolerance in hyperthyroidism were consistently reported. Furthermore, autoimmune thyroid diseases, particularly Hashimoto's thyroiditis and Graves' disease, were linked to long-term metabolic consequences, including increased cardiovascular risk and altered energy homeostasis. Subclinical thyroid dysfunction was also shown to influence metabolism, although the effects were milder and sometimes controversial.

Discussion

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

The metabolic alterations observed in thyroid dysfunction are largely attributed to the systemic effects of thyroid hormones, especially T3, which acts directly on nuclear receptors to regulate gene expression involved in energy balance. The literature consistently demonstrates that even minor changes in thyroid hormone levels can significantly affect multiple metabolic pathways.

In hypothyroidism, the reduced availability of T3 leads to decreased mitochondrial activity and oxygen consumption, which in turn slows metabolism and promotes lipid accumulation. These mechanisms explain the clinical manifestations such as weight gain, fatigue, and dyslipidemia. Conversely, hyperthyroidism increases mitochondrial biogenesis and oxidative phosphorylation, enhancing thermogenesis and catabolic processes.

An important point is the impact of thyroid dysfunction on cardiovascular metabolism. Hypothyroidism contributes to atherosclerosis through increased LDL levels and reduced clearance, while hyperthyroidism may induce arrhythmias due to increased cardiac output and sensitivity to catecholamines.

Subclinical thyroid dysfunction, although often asymptomatic, has emerged as a significant area of research due to its subtle but clinically relevant effects on lipid metabolism, insulin sensitivity, and cardiovascular risk markers. These findings suggest the importance of early detection and individualized treatment approaches, even in borderline cases.

Despite clear trends in the literature, some inconsistencies remain—particularly in studies involving subclinical conditions and inter-individual variability in metabolic responses. Factors such as age, sex, comorbidities, and genetic background may influence outcomes, necessitating more personalized approaches in future research.

CONCLUSION

Thyroid dysfunction has profound and multifaceted effects on human metabolism. Both hypothyroidism and hyperthyroidism disrupt normal metabolic processes, affecting lipid, carbohydrate, and protein metabolism, as well as thermoregulation and energy expenditure. The degree and nature of these disturbances vary with the type and severity of thyroid disorder.

This review highlights the central role of thyroid hormones in maintaining metabolic homeostasis and underscores the importance of timely diagnosis and management. Even subclinical thyroid dysfunction can influence metabolic health and should not be overlooked in clinical practice.

Future studies should focus on better understanding the molecular mechanisms underlying these effects, improving diagnostic sensitivity for early dysfunction, and developing targeted therapies that address not only hormonal imbalance but also its metabolic consequences.

REFERENCES

1. Brent, G. A. (2012). Mechanisms of thyroid hormone action. *The Journal of Clinical Investigation*, 122(9), 3035–3043. https://doi.org/10.1172/JCI60047
2. Chaker, L., Bianco, A. C., Jonklaas, J., & Peeters, R. P. (2017). Hypothyroidism. *The Lancet*, 390(10101), 1550–1562. https://doi.org/10.1016/S0140-6736(17)30703-1
3. Smith, T. J., & Hegedüs, L. (2016). Graves' Disease. *The New England Journal of Medicine*, 375(16), 1552–1565. https://doi.org/10.1056/NEJMra1510030
4. Vanderpump, M. P. J. (2011). The epidemiology of thyroid disease. *British Medical Bulletin*, 99(1), 39–51. https://doi.org/10.1093/bmb/ldr030
5. World Health Organization (2020). *Thyroid disorders – Key facts and statistics*. WHO Press.
6. Ministry of Health of the Republic of Uzbekistan. (2022). *Statistical report on the epidemiology of endocrine diseases*. Tashkent.
7. Zimmermann, M. B., & Boelaert, K. (2015). Iodine deficiency and thyroid disorders. *The Lancet Diabetes & Endocrinology*, 3(4), 286–295. https://doi.org/10.1016/S2213-8587(14)70225-6