ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

TECHNOLOGY OF INTENSIVE IRRIGATION OF LEMON SEEDLINGS

Head of the Educational and Methodological Department, Toshloq Vocational Service and Technical College"

Zokirova Mohiraxon Lutfullayevna

email. mohiraxonzokirva1985@gmail.com

Annotation. This article analyzes the theoretical foundations and practical approaches of intensive irrigation of lemon seedlings in the field of lemon cultivation. Nowadays, lemon cultivation and lemon products are widely used not only in agriculture but also in food, confectionery, industry, and healthcare. One of the main principles in lemon production is the use of the most efficient methods to achieve high yields with minimal inputs. Among the most important requirements for lemon cultivation is the proper selection of the irrigation system. In times of drinking water scarcity, the application of intensive irrigation methods in agriculture is considered one of the most pressing issues.

Keywords: lemon, intensive irrigation, drip irrigation, water-saving irrigation

Introduction

At present, the rational use of water resources in the cultivation of agricultural crops is one of the most urgent issues. Against the background of global climate change, water scarcity, and the expansion of agricultural lands, the need to introduce intensive irrigation technologies is increasing. In particular, in the cultivation of lemon (Citrus limon L.), which belongs to the group of subtropical fruit trees, the application of effective irrigation methods is one of the key factors in achieving high yields. In traditional irrigation methods, a large portion of water is wasted due to evaporation or seepage. Therefore, in lemon cultivation, intensive irrigation methods, including drip irrigation technology, have been showing highly effective results.

Research Methodology. Intensive irrigation of lemon seedlings is a system of providing water to plants that is more efficient, regulated, and controlled compared to traditional methods. It is mainly applied to increase yield and ensure water-saving.

Main characteristics: Implemented through drip irrigation or sprinkler systems.

Water is supplied continuously but in small quantities, reaching directly to the root zone., Possibility to apply fertilizers together with irrigation water (fertigation).

Irrigation is often automated, Soil moisture is maintained at an optimal level, preventing both excess water and drought stress.

Advantages:

Saves water (up to 30–50%), Lemon roots develop better due to constant moisture supply, Improves both quality and quantity of yield.

Reduces risks of pests and diseases (since the soil surface does not remain excessively wet). Thus, intensive irrigation means regularly, efficiently, and economically watering lemon seedlings using drip or other modern irrigation methods. The following schedule mainly applies to drip irrigation but can also be adapted for sprinkler systems. Intensive Irrigation Schedule for Lemon Seedlings After planting (1–3 months):

Objective: to promote rapid root establishment. Irrigation: daily or every other day. Rate: 2–3 liters per seedling. Vegetation period (April–June): Stage: leaf development and branching. Irrigation: every 2–3 days. Rate: 5–7 liters per seedling. Flowering and fruit setting (May–July): Water deficit

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

during this stage causes flower drop. Irrigation: every 2 days, rate: 7–10 liters per seedling. Fruit growth stage (July–September): Fruits develop fully. Irrigation: daily or every other day. Rate: 10–12 liters per seedling. Ripening period (October–December): Regular irrigation is essential to improve fruit quality and sweetness. Irrigation: every 2–3 days. Rate: 7–10 liters per seedling. Winter period (January–March): Stage: dormancy. Irrigation: once every 7–10 days. Rate: 3–4 liters per seedling.

Additional recommendations: Soil moisture should be checked with a moisture meter or manually. Avoid excessive water accumulation (to prevent root rot). Adding dissolved mineral fertilizers (nitrogen, phosphorus, potassium) during irrigation is highly effective. In greenhouses, the irrigation schedule must be adjusted as air is drier and water demand is higher. Drip irrigation is the most effective method for greenhouse lemon seedlings. Fertigation every 15–20 days is recommended. During hot days (above 30–35 °C), light foliar spraying (mist irrigation) is beneficial.

Soil moisture should be maintained at 70–80%. For lemons planted in soil inside greenhouses, irrigation amounts should be higher than in pots due to increased evaporation and soil absorption capacity. Intensive Irrigation Schedule for Lemon Seedlings in Greenhouses (Soil-Based) After planting (1–2 months): Irrigation: daily or every other day. Rate: 3–4 liters per seedling. Objective: to promote rapid root establishment. Active growth period (March–June): Irrigation: every 2 days.

Rate: 6–8 liters per seedling. Note: during branching and leaf formation, water demand is high. Flowering and fruit setting period: Irrigation: every 1–2 days. Rate: 10–12 liters per seedling. Note: water deficiency may cause flower and fruit drop.

Additional recommendations: Drip irrigation is the most efficient method; it saves water and maintains constant soil moisture. Applying dissolved mineral fertilizers (NPK and microelements) 1–2 times per month during irrigation is beneficial.

Even if the upper soil layer appears dry, it is important to monitor moisture in deeper layers (using a moisture meter or manual check). During very hot days (above 35 °C), light foliar spraying (mist irrigation) provides good results.

Drip Irrigation Recommendations for Lemon Seedlings in Greenhouses

Placement of drippers: Each seedling should have two drippers placed around the root zone (on opposite sides). For older plants (after 3 years), it is advisable to use four drippers. Dripper flow rate Standard dripper: 2–4 liters/hour.

Period	Irrigation duration	Frequency	Water volume (per seedling)
After planting (1–2 months)	30–40 minutes	Daily or every other day	3–4 liters
Active growth (March–June)	40–60 minutes	Every 2 days	6–8 liters
Flowering and fruit setting (April–July)	1–1.5 hours	Every 1–2 days	10–12 liters
Fruit growth (July–September)	1.5–2 hours	Daily or every other day	8–10 liters
Fruit ripening (October–December)	1–1.5 hours	Every 2 days	8–10 liters
Winter dormancy (January–February)	30–40 minutes	Once every 7–10 days	4–5 liters

Water volume should be increased as the plant grows.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889,
2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

Most Economical Drip Irrigation System (for Greenhouse Lemon Cultivation)

1. Main components of the system: Water source – reservoir or large tank (1000–2000 liters). Filter – simple sand or disk filter (to prevent dripper clogging). Main pipeline – polyethylene (Ø32–40 mm). Lateral pipelines – 16 mm PE pipes, one for each row. Drip tape or emitters: Most economical option – drippers with 2 L/hour flow rate. Initially, 2 drippers are installed per lemon seedling; later, as the tree grows, this is increased to 4 drippers. Venturi injector – for applying fertilizers with irrigation water (fertigation).

2. System operation procedure:

Irrigation can be carried out either by pump or by gravity from an elevated tank (more economical without a pump).

Water is delivered directly to the root zone during each irrigation.

Average water consumption: 8–12 liters per seedling per day (depending on growth stage).

Fertilizers should be dissolved in water and applied through drip irrigation – the most efficient and economical method.

3. Advantages:

Reduces water consumption by 30–50%.

Requires almost no electricity (when using gravity-fed system).

Fertilizers are delivered precisely with irrigation water.

Soil remains constantly moist without waterlogging.

Each tree receives the exact required amount of water.

4. Recommendations (for low-cost setup):

Place a 1000-liter tank at a higher level near the greenhouse and connect 16 mm PE pipes. Install two 2 L/hour drippers per seedling, use a simple mesh or sand filter.

Later, as yield increases, the tank can be replaced with a small water reservoir and a low-cost pump.

Conclusion and Recommendations The application of intensive irrigation technology in lemon cultivation not only increases productivity but also ensures efficient use of water resources. The drip irrigation system promotes stable plant growth, improves fruit quality, and enhances economic efficiency. In the future, the widespread implementation of automated smart irrigation systems in lemon farming will help achieve more sustainable results.

REFERENCES

1. Karimov A., Tursunov S. “Sug‘orish texnologiyalari va ularni samaradorligi”. Toshkent: Fan, 2020.

2. Rasulov I. “Intensiv bog‘dorchilik asoslari”. Samarqand, 2019.

3. FAO Report. “Efficient Water Management in Horticulture”, Rome, 2021.

4. Bekchanov M., Mirzaev Sh. “Tomchilatib sug‘orish tizimlari va ularning agrotexnik xususiyatlari”. Toshkent, 2022.