ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

UDK: 631.41:631.445:631.8

RESULTS OF AGROCHEMICAL ANALYSIS OF TYPICAL SIEROZEM SOILS OF RAINFED AGRICULTURAL AREAS

I.SH. Mamatqulov. J.T.Naxalbaev. A.S.Muratkasimov.U.Z.Usmanov

Lalmikor dehqonchilik ilmiy tadqiqot instituti

e-mail: uzniizerno@yahoo.com

Annotation. This article describes the results of the study on determining the agrochemical properties and mechanical composition of typical sierozem soils of rainfed agricultural areas of the Gallaorol district, as well as analyzing their level of productivity.

Keywords: typical sierozem soils, siderate crops, agrochemical, agrophysical properties, water-physical properties, soil mechanical composition, soil nutrient regime.

Introduction. To improve the productivity and quality of agricultural crops, of course, the restoration and improvement of soil fertility is the most relevant. It is known that the procedure for short-term crop rotation is being established instead of long-term crop rotation, which is currently practiced in rainfed fields. In increasing soil fertility, the productive use of organic fertilizers, green manure crops is considered important.Decree of the president of the Republic of Uzbekistan dated June 24, 2024 PD-233 “on measures to create a sustainable agroecosystem in relation to climate change and increase the adaptability of agricultural products growers to the risks associated with climate change”Annex 1

Paragraph 40.1 sets out important strategic tasks to” improve the knowledge of agricultural producers to improve soil fertility by planting green manure crops ". Therefore, it is important to improve the agrophysical and agrochemical properties of degraded soils in the rainfed regions, grow green manure crops that serve to effectively use natural moisture, increase soil fertility and crop yields, wrap them in their impact on soil fertility and physical properties.

Analysis of thematic literature. One of the factors that prevent the effective use of dryland areas in the Republic today is low soil fertility. For many years, it was found that the single domination of grain crops in these areas, a sharp decrease in annual and perennial legumes and pure plowing areas, the continuous plowing of soil with ploughs to a depth of 20-22sm, the almost non-use of organic and mineral fertilizers, weeds, various diseases and pests, with an increase in the level of erosion,According to the FAO, the main problems of rich soils found in arid regions, including the rainfed lands of Uzbekistan, were discussed: salinity, abundance of carbonates, lack of organic matter. Agroecological adaptive approaches, sustainable fertilization systems, soil monitoring and alternative technologies have been recommended by the FAO to address these problems [3].A.Karimov and Sh.Juraev have conducted a comprehensive analysis of the agrophysical and agrochemical condition of typical rich soils found in the rainfed regions of Uzbekistan. The study argues that low humus content, limited nitrogen and phosphorus reserves make it difficult for crops to grow on such soils. It has also been noted the need to use organic matter to increase soil fertility [4].

A.Hidirov's scientific research focuses on increasing soil fertility through biological methods. Biostimulants, microcircuits and green fertilizing technologies have been found to have positive effects on lalmikor soils.[6].

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

The morphogenetic indicators of the soil change mainly depending on the characteristics of use. In particular, the color, mechanical composition, structurally different from these indicators of the ploughing layer of arable land.[7].The mechanical composition of the soil is associated with the composition of the mother genera, and as a result of crop rotation, qualitative processing of the soil, its general physical properties are improved[7].

Research methodology. Field experiments in research is a “methodological guide to conducting field experiments "developed by scientists from USRIC [2], soil analysis Y.V.Arinushkina's “Руководство по химическому анализу почв” [9], statistical analysis of the results obtained B.A.Dospekhov's methodological manuals “Методика полевого опыта” [8] and all agrotechnological activities were carried out on the basis of agrorecommendations developed by the Research Institute of Rainfed Agriculture in Gallaaral district .

Field – experimental research work on the effect of green manure crops on soil fertility and physical properties in Rainfed Fields was carried out in the Central Experimental Farm of the Research Institute of Rainfed Agriculture of Gallaaral district**.**

Analysis and results. Typical loamy soils are common in Central Asia and other arid climate areas. Formed mainly in arid climates, they are important for agriculture.

Total plowed agricultural land in Uzbekistan is 773.6 thousand hectares, 229.4 thousand hectares in Jizzakh region, 258.4 thousand hectares in Kashkadarya region, 30.9 thousand hectares in Navoi region, 182.1 thousand hectares in Samarkand region, 39.5 thousand hectares in Surkhandarya region, 33.3 thousand hectares in Tashkent Region [5].

Mechanical composition, which is one of the most important properties of soil, is one of the main indicators that determine its fertility, knowledge of information about the mechanical composition of dryland areas, ensures the improvement of the soil treatment system, the correct placement of crop species and an increase in the efficiency of the technique.

We used classification of mechanical elements that was recommended by N.A.Kachinsky. According to the classification of mechanical elements recommended by Kachinsky, the mechanical composition of the soil - water permeability, moisture capacity, Air is one of the important properties that determine soil fertility, being directly related to the water and heat order [10]. Soils of different mechanical composition have different levels of productivity, and the physical and mechanical properties of the soil, which determine the resistance to agricultural techniques when plowing the soil, will also depend on the mechanical composition.

The mechanical composition of the soil is associated with the composition of the mother genera, and as a result of crop rotation, qualitative processing of the soil, its umimic physical properties are improved.

Table 1 shows the mechanical composition, i.e. particle fractions (as a percentage of weight) and the mechanical composition, according to the different layers of the typical peat soil section, which is found in rainfed (non-irrigated) conditions. Based on the table, the mechanical composition of the layers is as low as below.

Layer A_1 (0-22 sm) is the layer of the soil surface, and it is on this floor that the most active agrobiological processes take place. The amount of sand fraction (especially fine sand) in it is relatively high, ensuring light soil processing, good ventilation and water permeability. The physical clay content is 34.2%, which maintains the lightweight structure of the layer.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

In general, the A_1 layer is the most important agrotechnical layer, with a high level of productivity, where plant roots are actively located. The medium sand content makes this layer a favorable environment for crops.

Despite the fact that the A_2 layer (22-37 sm) is close to the upper layer, it has a higher amount of physical clay, which makes it possible to evaluate the layer as middle sand. The presence of 36.6% of physical clay ensures that moisture is relatively retained. This layer creates favorable conditions for the good development of the roots.

The amount of sand in the B_1 layer (37-70 sm) decreased, while the proportion of clay fractions increased. This condition increases the compaction of the layer, which can negatively affect air exchange. At the same time, the amount of physical clay (37.4 %) increases the water retention property of the soil. In this layer, nutrients can accumulate a little more, which ensures a deeper development of plant roots.

The B_2 layer (70-122 sm) has a slightly higher physical clay content (38.2 %) than the b_1 layer, which brings the layer closer to a heavier state than the middle sand. The water storage property is strong, but the possibility of air exchange is reduced. The spread of plant roots in this layer can be difficult. However, as a moisture-retaining layer, it plays a beneficial role in arid conditions.

The C_1 layer (122-160 sm) is the mother genus, which represents the final stages of soil formation. The amount of physical clay in it (42.9%) is higher, which is the density of the layer and holds water

Table 1

Mechanical composition of typical sierozem soils of rainfed agricultural areas

Profile number and name	Layer	Layer thickness, cm	Mechanical composition (fraction weight),%								Nomenclature according to mechanical composition
			>0,25	0,25-0,1	0,1-0,05	0,05-0,01	0,01-0,005	0,005-0,001	<0,001	Physical clay. <0.01	
1	A_1	0-22	2,0	2,4	20,1	41,3	9,5	19,9	4,8	34,2	Medium sand
	A_2	22-37	1,7	3,5	24,0	34,2	11,9	19,1	5,6	36,6	Medium sand
	B_1	37-70	2,8	3,6	22,1	34,2	12,7	18,3	6,4	37,4	Medium sand
	B_2	70-122	1,9	3,7	22,9	33,4	12,7	17,5	8,0	38,2	Medium sand
	C_1	122-160	1,9	1,4	17,2	36,6	16,7	18,3	8,0	42,9	Heavy sand

In order to study the initial state of the soils of the experimental research area, the main soil profile was obtained. Humus content and total nutrient elements (nitrogen, phosphorus and potassium) as well as soil environment indicators were found to contain in these typical sierozem soils of rainfed agricultural areas .

The amount of humus in typical sierozem soils of rainfed agricultural areas under research is 1,549-1,181% in the plow layer and sub-plow layers, with the humus indicator averaging, while in

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-8

the lower layers its amount is 1,101-0,896%, making it a low indicator. These soils have a C: N ratio of 8.0-10.2, with an average nitrogen supply of humus (Table 2).

The total nitrogen content is 0.051 - 0.091 %, the phosphorus content is 0.120-0.350% in layers, reduced from above along the soil profile to the lower layers, is in the upper pointer. The total potassium content is 0.542-1.320% by profile and corresponds to a low indicator. These soils have been found to be neutral and weakly alkaline environments with an environment indicator range of 7.0-7.4.

Table 2

Agrochemical composition of typical sierozem soils of rainfed agricultural areas , %

№ Profile	Layer thickness, cm	Humus	N	C:N	P	K	pH
1	0-22 (A_1)	1,549	0,091	9,9	0,350	1,320	7,4
	22-40 (A_2)	1,181	0,080	8,6	0,230	1,007	7,0
	40-75 (B_1)	1,102	0,080	8,0	0,170	0,817	7,2
	75-140 (B_2)	0,937	0,059	9,2	0,120	0,621	7,1
	140-200 (C_1)	0,896	0,051	10,2	0,120	0,542	7,0

In the resulting cross section, it was observed that humus, ammonium state nitrogen, phosphorus in general, potassium in general and nitrogen in the ammonia state decrease downward from the top layer**.**

Conclusions and suggestions. As a result of the analysis carried out, it was found that there are significant differences in the mechanical composition of the layers of the typical sierozem soils of rainfed agricultural areas section at different depths. The upper layers of the soil were assessed as having a middle sand mechanical composition. Deeper layers are also of medium sand content, in which there is a gradual increase in the amount of physical clay. The lowest layer of the cross-section is a heavy sand formation, with a physical clay (<0.001 mm) content of 42.9, indicating a high soil density and water retention capacity. This situation is important in land processing, when choosing the types of crops. In particular, the predominance of the middle sand layers creates the ease of water permeability of the soil, respiration and agrotechnical activities.

Agrochemical analysis of typical sierozem soils of rainfed agricultural areas has shown that they are inadequate in terms of organic matter and food elements.

In general, these soils are poor in humus, nitrogen and phosphorus, but rich in potassium, and a neutral or slightly alkaline pH makes them conditions for the cultivation of various crops. However, agrotechnical measures are a prerequisite for obtaining a high yield.

Bibliography.

1. A.S.Muratkasimov .The current state of typical sierozem soils of rainfed agricultural areas and ways of their effective use (on the example of soils of Gallaorol district): his dissertation, written to obtain the degree of Doctor of Philosophy (PhD)in Agricultural Sciences.Tashkent, 2019. - S. 57-58 page.

2. Methods of conducting field experiments// USRIC, Tashkent, 20.07, page. 1-146.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-8

3. FAO Soils Bulletin No. 79. (2001). Soil Fertility Management in Support of Food Security in Sub-Saharan Africa. – Rome: FAO. – 132 b.

4. Karimov A., Zhuraev Sh. (2020). Soils of Uzbekistan and their fertility. - Tashkent: publishing house" Fan". – 156 p.

5.R.A.Turayev, R.N.Sharopov. Rainfed field monitoring / / monograph. - Tashkent "Lesson Press" LLC, 2023. 15 p.

6.Khidirov A. (2019). The effect of biostimulants on the soil of rainfed fields. - Journal "Agriculture and innovation", №4. – 60-66.

7.Yusupov X., Muratkasimov A., Usmanov U. Description of the morphological structure and mechanical composition of typical sierozem soils. Eurasian Journal of Medical and Natural Sciences, 2023, 3(2 Part 2), 70-74 PP.

8.Dospekhov B.A. Methodology of Field Experiments (with the Basics of Statistical Processing of Research Results) Moscow. Agropromizdat, 1985. -351 p.

9.E.V. Arinushkina. Manual on Chemical Analysis of Soils, Moscow University Press, 1970. 137-218 p.

10. Methods of agrochemical, agrophysical, and microbiological research in agriculture of the cotton region // Soyuz NIKSI, Tashkent. 1963. Pp. 270-274.