ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

ENERGY VARIATIONS OF 4s-3d ORBITALS AND EXCEPTIONS IN ELECTRON CONFIGURATIONS OF Cl AND Fe ELEMENTS

Eshchanov Ruzumboy Abdullaevich, Prof., Chirchik State Pedagogical University, Chirchik
ORCID: https://orcid.org/0000-0002-8594-2907
ruzimboy@gmail.com

Abstract. This article analyzes the variability of the energy levels of 4s and 3d orbitals, as well as the peculiarities of electron filling sequences in the elements Cl and Fe. It is shown that the various valence states of Cl (1, 3, 5, and 7) and Fe (2, 3, and 6) depend on the electron configuration and orbital energies. Furthermore, it is noted that the classical interpretation of Klechkovskiy's rule does not always apply to real atomic systems, as electrons are distributed between outer and inner shells depending on their interactions and energy balance. The examples of Cl and Fe demonstrate that the complex distribution of electrons between outer and inner shells creates difficulties for students and in teaching theoretical chemistry.

Keywords: 4s orbital, 3d orbital, electron configuration, valence states, Cl element, Fe element, orbital energy, Klechkovskiy rule, outer shell, inner shell, electron distribution, valence exceptions, energy balance, teaching difficulties.

The arrangement of electrons in orbitals is one of the most important factors determining the properties of chemical elements. In modern quantum chemistry, this arrangement is based on Klechkovskiy's rule (or the (n + l) rule) [1]. According to this rule, electrons first occupy orbitals with the smallest (n + l) value; if two orbitals have the same (n + l) value, the orbital with the smaller n is filled first [2]. Following this principle, the 4s orbital, having a smaller (n + l) value than the 3d orbital, should be filled before the 3d orbital.

However, experimental observations and quantum mechanical calculations show that this rule does not always work precisely. In particular, for elements such as chlorine (Cl) and iron (Fe), the actual electron distribution deviates from Klechkovskiy's rule. Such deviations are especially pronounced in excited states or during ionization processes.

For example, in the excited state of a chlorine (Cl) atom, an electron can move from the 3p orbital to the 3d orbital, even though the energy of the 4s orbital is theoretically considered lower. This phenomenon is explained by the complexity of electron–electron interactions, the close energy levels of the orbitals, and spin–orbital coupling. Similarly, in an iron (Fe) atom, the 4s orbital is filled before the 3d orbital, but during ionization, electrons are removed first from the 4s orbital. This demonstrates that the energy levels of the 4s and 3d orbitals are variable and depend on the specific conditions.

Thus, Klechkovskiy's rule should be regarded only as an empirical guideline applicable to ideal cases. Although it determines the general order of electron filling, in real atomic systems, as seen in Cl and Fe, orbital energies are not constant but vary depending on electron density, shielding effects, and the charge of the atomic nucleus. Therefore, Klechkovskiy's rule alone is insufficient to explain the energy balance between the 4s and 3d orbitals.

The arrangement of electrons in orbitals and the valence states of atoms in Cl and Fe are associated with complex phenomena and represent exceptions to the simple interpretation of

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

Klechkovskiy's rule. In particular, the various valence states of chlorine (Cl) — 1, 3, 5, and 7 — are directly related to its electron configuration and orbital energies [3].

The 1-valence state of a chlorine (Cl) atom is observed in its ground (normal) state. In this state, the electrons in the outer shell are sufficient, no additional energy is required, and the atom remains relatively stable. However, in the 3-valence state of a Cl atom, the electrons occupy the configuration $3s^2\ 3p^4\ 3d^1$ rather than $3s^2\ 3p^4\ 4s^1$. This occurs because forming the configuration by removing or adding electrons in the 4s orbital would result in relatively higher energy. Therefore, the 3-valence state of Cl requires the electron to move into the 3d orbital, which is explained by the energetic balance between orbitals and electron–electron interactions. This phenomenon cannot be fully explained by Klechkovskiy's rule, as its empirical nature does not account for energy variations in real atomic systems.

Similarly, in an iron (Fe) atom, the 4s orbital is filled before the 3d orbital. However, the various valence states of Fe, specifically 2, 3, and 6, create complexities in electron distribution. For example, in the 2-valence state of Fe, in its active form, electrons occupy the $4s^1$ and $4p^1$ orbitals. In the 3-valence state, although the $4s^1$ and $4p^1$ electrons remain in the outer shell, the atom also utilizes electrons from the inner $3d^6$ orbitals. Likewise, in the 6-valence state, electrons are not limited to the outer shell but also interact with the inner 3d orbitals.

Consequently, the distribution of electrons between the outer and inner orbitals in Fe causes confusion for many students and learners. This situation also presents challenges in theoretical chemistry and teaching, as classical rules and principles cannot fully explain the actual electron distribution in the valence states of the Fe atom.

Thus, the transition of electrons between outer and inner orbitals observed in the examples of Cl and Fe is explained by the condition-dependent variation in the energy levels between the 4s and 3d orbitals. These phenomena cannot be fully explained by Klechkovskiy's rule and require a more detailed analysis to determine the actual electron configuration and valence states of the atom.

In this article, the variability of the energy levels of 4s and 3d orbitals and the exceptions in the electron filling order in Cl and Fe elements are analyzed. In the case of Cl, it was shown that the different valence states — 1, 3, 5, and 7 — depend on the electron configuration and the variation in orbital energies. In particular, in the 3-valence state of a Cl atom, electrons occupy the configuration $3s^2\ 3p^4\ 3d^1$ rather than $4s^1\ 3p^4$, which is explained by achieving a more energetically stable state. In this way, the electron distribution of the Cl atom manifests as an exception to classical rules and principles, including Klechkovskiy's rule.

In the case of the iron (Fe) element, the 4s orbital is filled before the 3d orbital; however, when valence states are formed, the electrons are not limited to the outer 4s and 4p orbitals but also involve the inner 3d orbitals. For example, in the 2-valence state of Fe, electrons occupy the $4s^1$ and $4p^1$ orbitals, while in the 3-valence and 6-valence states, in addition to the outer shell electrons, electrons from the inner 3d orbitals are also utilized. Consequently, the balance of electrons between the outer and inner orbitals in Fe causes confusion for students and cannot be fully explained by classical rules and principles.

In conclusion, the exceptions observed in the electron configurations and valence states of Cl and Fe create complexities in understanding the true behavior of these atoms and also demonstrate the difficulty of explaining these topics in a sufficiently simple manner in teaching and theoretical chemistry. Therefore, when explaining the distribution of electrons between outer and inner orbitals,

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

the condition-dependent energy of orbitals, and exceptions in valence states, it is necessary not to rely solely on empirical rules but to develop new theoretical approaches to understand such exceptions.

References

1. OpenStax. Chemistry 2e: Section 6.4 “Electronic Structure of Atoms (Electron Configurations)”. OpenStax, 2019.
2. LibreTexts. “Electron Configuration” (Introductory module). UCalgary & LibreTexts, 2023.
3. University of Calgary. “Electron Configurations of Atoms: Introductory Chemistry Textbook Material.”
4.