ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

CLINICAL CHARACTERISTICS OF VIRAL DIARRHEA IN HIV-INFECTED CHILDREN

Otajanov Shamsiddin Zarifbayevich

Urgench State Medical Institute
Assistant Professor, Department of Infectious Diseases, Epidemiology and Phthisiology, PhD.

Annotation. Viral intestinal infections play an important role in infectious morbidity in children of different ages.Case histories of 135 children were analyzed, the examination algorithm of which included polymerase chain reaction using primers manufactured by AmpliSense® Rotavirus/Norovirus/Astrovirus-FL or AmpliSense®, AmpliSense® Enterovirus-FL » manufactured by Interlabservis. It was found that when using the polymerase chain reaction, the diagnosis was etiologically deciphered in 80 (59%), and the proportion of viral diarrhea among the deciphered acute intestinal infections was 77.35%. It has been shown that children under 3 years of age (45%) suffer predominantly from viral diarrhea. The prevalence of rotavirus infection (40.44%) was revealed, and the proportion of adenoviral gastroenteritis was also high (21.18%). It was established that rotavirus is the first in importance in the age group of children under 6 years of age (56%), in school-age patients adenovirus was in first place (45.2%). The incidence of rotavirus infection remained high throughout the study period (44.6–58.9%), the incidence of adenovirus gastroenteritis increased from 24% in 2018 to 32% in 2019. In most cases (82%), viral gastroenteritis occurred in moderate form, the main complication was dehydration syndrome of 1–2 degrees.

It has been proven that using modern diagnostic methods, at least 60% of cases of infectious diarrhea in HIV-infected people are deciphered, and the main etiological agents are rotaviruses and adenoviruses. Our results are fully consistent with the data of foreign sources on the epidemiology of AII in children before the start of mass vaccination of rotavirus infection.

Key words: intestinal infections, children, viral diarrhea, HIV-infection, PCR.

Actuality. The urgency of the problem of acute intestinal infections (AII) is due to the fact that this group of infectious diseases steadily retains one of the leading places in the structure of childhood morbidity and mortality. Every year, AII causes the death of more than 1 million children worldwide [1]. Mortality is especially high in children of the first years of life. Currently, rotaviruses are considered as the main etiological factor of AII, especially in children under 5 years of age. According to the World Health Organization (WHO), about 138 million ases of rotavirus infection are cregistered annually in the world [3].

In the US and other developed countries, according to U.D. Parashar, J.S. Bresee, prior to the introduction of specific prophylaxis, approximately 2.3 million children under 5 years of age were infected with rotavirus annually, resulting in an economic cost of more than $275 million per 500,000 medical consultations and 50,000 hospitalizations [3].

The damage from rotavirus infection is incomparably higher in the developing countries of South Asia, India, Latin America, where from 20 to 70% of hospitalizations are associated with this etiological agent. Mortality from rotavirus infection ranges from 454 to 705 thousand cases per year and averages 611 thousand cases [4]

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

It is believed that rotaviruses infect almost every child before they reach the age of 3–5 years [5]. In developing countries, the median age at first rotavirus infection ranges from 6 to 9 months (80% of cases occur in children under 1 year of age), while in developed countries, the first episode of infection may appear at 2–5 years of age, although most cases do observed among infants (65% of cases are recorded in children under the age of 1 year) [5].

At the same time, after the introduction of routine vaccination against rotavirus gastroenteritis, norovirus infection caused more than 50% of cases among AII outbreaks recorded in the United States according to data for 2013 [6, 7].

With the expansion of diagnostic capabilities, the deciphering of other viral intestinal infections in HIV-infected children has also increased.

According to most literature sources, adenoviruses of the serogroup (the so-called "intestinal" adenoviruses) are the third most important etiological agent of viral AII. Adenovirus affects all age groups, has no distinct seasonality, it is characterized by a contact route of infection. Often there are reports of outbreaks of adenovirus infection. Species D adenovirus, in particular HAdV-67, was isolated from faeces during a major AII outbreak in Bangladesh, and serogroup F adenoviruses have been reported in India in children with acute diarrhea in the under-5 year age group [10–12]. In the world literature, a relatively small number of publications are devoted to astrovirus infection. The study of the role of astroviruses in the development of acute non-bacterial gastroenteritis began in 1975, when using the method of electron microscopy, they were first detected in the feces of children with diarrhea. When studying the prevalence of astrovirus infection among children, it was possible to establish that up to 71-75% of children aged 3 to 10 years have antibodies to astroviruses, although they had no signs of the disease in their anamnesis. It is known that astroviral gastroenteritis is often mild in HIV-infected children. Virus-neutralizing antibodies to serotype 1 were detected in 92% of cases, serotype 3 - in 69%, serotype 4 - in 50%, serotype 5 - in 36% and serotype 2 - in 31% of the examined children. Outbreaks of diarrhea associated with astroviruses occur in preschool institutions, schools, nursing homes. Most often, outbreaks of astrovirus infections are caused by the first serotype of the virus [13]. Interesting data were obtained in the course of a study in which 17 adults took part. It was found that astroviruses have low pathogenicity, because, despite infection, clinical signs of the disease were registered in only one person, while the antibody response was detected in most subjects [13].

According to Australian scientists, in 2009 an astrovirus caused an outbreak of AII among schoolchildren [14].

In Japan, after the introduction of mass vaccination against rotavirus in developing countries, the frequency of detection of astrovirus infection increased from 5 to 20% [14].

The purpose of the study is to study the etiological structure and features of the course of viral diarrhea in HIV-infected children in 2018–2021.

Materials and methods.

Under supervision in 2018–2021 There were 135 children hospitalized at the DIB clinic, of which 80 (59%) were with verified AII, 61 (45.18%) were with confirmed viral AII in HIV-infected children. The proportion of diarrhea of viral origin among the deciphered AII was 77.35%.

Assessment of the clinical course of infectious diarrhea was carried out in accordance with the principle of determining the type, severity, nature of the course of the infectious process [15]. When assessing the severity of the disease, the severity of symptoms of intoxication, temperature reaction and local gastrointestinal syndrome were taken into account. Particular attention was paid to

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

the course of the infectious process, namely, the non-smooth nature of the course: the presence of complications, exacerbations or relapses. An assessment was made of the dehydration syndrome and forms of severity in 77.35% of the examined children with viral diarrhea of various etiologies.

All patients underwent a standard laboratory examination, which included: clinical blood and urine tests, biochemical blood tests, coprocytogram; etiological interpretation of the diagnosis included, in addition to the microbiological method, bacteriological seeding of feces on solid nutrient media to isolate a pure culture of the pathogen, PCR of feces for bacterial and viral agents (detection of RNA / DNA of pathogens). Testing of the collected fecal samples was performed using the reagents "AmpliSense® Rotavirus/Norovirus/Astrovirus-FL" or "AmpliSense® OKI screen-FL", "AmpliSens® Enterovirus-FL" by polymerase chain reaction (PCR) with hybridization-fluorescence detection of production " Interlabservis.

Results and discussion.

In the course of the study, it was possible to identify the following features of viral diarrhea in children.

In 2018–2021 when using routine methods for verifying infectious diarrhea (microbiological method and faecal ELISA), the frequency of deciphered intestinal infections was 48%, of which 12.5% were salmonellosis, 10.3% were campylobacteriosis, 7% each were shigellosis and AII conditionally pathogenic etiology, 3.5% for yersinia infection, 5.4% for escherichiosis, and 52.3% for viral gastroenteritis [16].

When analyzing the structure of diarrhea decoded using PCR in 2018–2021, the proportion of AII decoded in HIV-infected children was 59%.

According to world statistics, in recent years there has also been an increase in the frequency of detection of rotaviruses in the structure of acute intestinal infections among hospitalized children [3].

We compared the etiological structure of viral diarrhea in different periods. According to the Department of Intestinal Infections, in 2004-2009. the proportion of rotavirus infection was 56% and occupied a dominant position in the etiological structure of viral AII, 25% was assigned to the share of calicivirus gastroenteritis [16].

The incidence of rotavirus infection in HIV-infected children remained high throughout the study period (44.6–58.9%), the incidence of adenovirus gastroenteritis increased from 24% in 2018 to 32% in 2021, the proportion of viral-viral associations increased from 8.4% in 2018 to 14.4% in 2021.

It has been proven that the age structure of children with viral diarrhea is as follows: 45.31% are children from 1 year to 3 years old, in second place are children from 3 to 6 years old - 20.42%, the proportion of children under 1 year old and children with 6 to 14 years is 19.82% and 12.68% respectively.

Conclusion.

Thus, it was found that using the polymerase chain reaction method, the frequency of etiological interpretation of AII in HIV-infected children increases (from 48% in 2018 to 60% in 2021).

The predominance of diarrhea of viral origin among the deciphered AII was shown - 77.35%. A high proportion of rotavirus infection in the overall structure of viral diarrhea was revealed - up to 48±1.6%; – 2.2±0.3% in the structure of morbidity.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

The age characteristics of viral monoinfections and mixed viral gastroenteritis have been established: up to 45% occur in children under 3 years of age, the second most common detection group is a group of children from 3 to 6 years old - 20.4%. In children older than 6 years, the proportion of adenovirus infection was 45.2%, viral-viral associations were more often detected in children of the first year of life - 13.1%. It has been proven that the frequency of development and the degree of dehydration depends on the etiological agent; with rotavirus infection and mixed viral diarrhea, up to 60% of cases occur with the development of dehydration of 1–2 degrees.

References

1. Kramarev SA, Zakordonets LV. Zdorovye rebenka. 2011; 1 (28): 53-6.
2. Annual Report of reference center for monitoring pathogens of acute intestinal infections (RTSKI) for 2012. Available from: http://www.epid-oki.ru/cont/analit_pism/Godovoj_otchet_referens centra_po_monitoringu_vozbuditelej_ostryh_kishechnyh_infekcij.pdf
3. Parashar UD, Gibson CJ, Bresee JS, et al. Rotavirus and severe childhood diarrhea. Emer. Infect. Diseases. 2006; 12(2): 304-6.
4. Parashar UD, Gentsch JR, Glass RI. RotavirusEmerg. Infect. Dis. 1998; 4 (4):561-570.
5. Ezhenedelnyy epidemiologicheskiy byulleten. 2013; 88: 49-64. Available from: http://www.who.int/wer.
6. Li Y, et al. An outbreak of norovirus gastroenteritis associated with a secondary water supply system in a factory in south China. BMC Public Health. 2013; 13:283.
7. Siddiqui U, Bini EJ, Chandarana K, et al. Prevalence and impact of diarrhea on health-related quality of life in HIV-infected patients in the era of highly active antiretroviral therapy. J Clin Gastroenterol. 2007;41:484–490.
8. MacArthur RD, DuPont HL. Etiology and pharmacologic management of noninfectious diarrhea in HIV-infected individuals in the highly active antiretroviral therapy era. Clin Infect Dis. 2012;55:860–867.
9. Gupta R, Ordonez RM, Koenig S. Global impact of antiretroviral therapy-associated diarrhea. AIDS Patient Care STDS. 2012;26: 711–713.
10. daCosta DiBonaventura M, Gupta S, Cho M, Mrus J. The association of HIV/AIDS treatment side effects with health status, work productivity, and resource use. AIDS Care. 2012;24:744–755.
11. Аскарова, Р. И., & Отажонов, Ш. З. (2020). Туберкулез на фоне сопутствующей патологии. *European science*, (3 (52)), 106-108.
12. Отажанов, Ш. З., Худайцулова, Г. К., & Муминова, М. Т. (2019). THE COURSE OF DIARRHEA OF VIRAL ETIOLOGY IN HIV INFECTED CHILDREN. *Новый день в медицине*, (4), 252-255.
13. Аскарова, Р. И., Атажанов, Ш. З., & Маткурбанов, Х. И. (2019). Анализ первичных форм туберкулеза у детей, проживающих в Хорезмской области. *Наука, техника и образование*, (9 (62)), 64-68.
14.Артиков, И. А., Отажанов, Ш. З., & Садуллаев, С. Э. (2025). ПАТОМОРФОЛОГИЧЕСКИЕ АСПЕКТЫ ДЕТСКОГО ЛИМФОБЛАСТНОГО ЛЕЙКОЗА. *Multidisciplinary Journal of Science and Technology*, *5*(6), 257-264.
15.Kadamovna, A. D., Rustombekovich, N. R., Zarifboyevich, O. S., & Ernazarovich, S. S. (2025). COMBINATIONS OF HEPATITIS B WITH PULMONARY TUBERCULOSIS. *Multidisciplinary Journal of Science and Technology*, *5*(5), 60-65.

16.Rustombekovich, N. R., Zarifboyevich, O. S., Kadamovna, A. D., & Ernazarovich, S. S. (2025). THE STATE OF THE ANTIOXIDANT DEFENSE SYSTEM IN CHRONIC HEPATITIS C. *Multidisciplinary Journal of Science and Technology*, *5*(5), 79-84.

17.Ernazarovich, S. S., Zarifboyevich, O. S., Rustombekovich, N. R., & Kadamovna, A. D. (2025). DYNAMICS OF THE COVID-19 PANDEMIC AND ITS CLINICAL CONSEQUENCES. *Multidisciplinary Journal of Science and Technology*, *5*(5), 85-90.

18.Худайкулова, Г. К., Муминова, М. Т., & Отажанов, Ш. З. (2021). АНАЛИЗ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ ВИРУСНЫХ ДИАРЕЙ У ВИЧ-ИНФИЦИРОВАННЫХ ДЕТЕЙ. *Journal of cardiorespiratory research*, *2*(3), 31-34.

19.Аскарова, Р. И., & Отажанов, Ш. З. (2020). Туберкулез на фоне сопутствующей инфекции. *European Science*, (3), 52.

20.Ibraximova, H. R., Artiqov, I. A., & Otajanov Sh, Z. (2025). HELMINTH INFECTIONS: CLASSIFICATION, DEVELOPMENTAL CONDITIONS, AND CLINICAL MANIFESTATIONS. *Multidisciplinary Journal of Science and Technology*, *5*(6), 1241-1245.

21.Юсупов, Ш. Р., Отажонов, Ш. З., Артиков, И. А., & Ибодуллаева, С. С. (2025). МИКРОБИОЛОГИЧЕСКАЯ СТРУКТУРА ОСТРЫХ ДИАРЕЙНЫХ ЗАБОЛЕВАНИЙ В ЮЖНОМ ПРИАРАЛЬЕ: РЕЗУЛЬТАТЫ ЭПИДЕМИОЛОГИЧЕСКОГО МОНИТОРИНГА. *Multidisciplinary Journal of Science and Technology*, *5*(6), 237-242.

22.Urinovna, M. O., Kadamovna, A. D., Zarifboyevich, O. S., & Axmedjonovich, A. I. (2025). PUBLIC HEALTH AND EPIDEMIOLOGICAL INSIGHTS INTO THE COVID-19 PANDEMIC. *Multidisciplinary Journal of Science and Technology*, *5*(6), 243-248.

23.ARTIQOV, I., & ABDULLAYEVA, D. (2025). THE SPREAD OF DIARRHEAL DISEASES IN THE KHOREZM REGION. *Multidisciplinary Journal of Science and Technology*, *5*(3), 672-677.

24.Otajanov, S. Z. (2025, January). CYTOKINE LEVELS IN HIV-INFECTED CHILDREN WITH VIRAL DIARRHEA. In *CONFERENCE ON THE ROLE AND IMPORTANCE OF SCIENCE IN THE MODERN WORLD* (Vol. 2, No. 1, pp. 148-148).

25.Худайкулова Гулнара Каримовна, Муминова Махбуба Тешаевна, & Отажанов Шамсиддин Зарифбаевич (2021). АНАЛИЗ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ ВИРУСНЫХ ДИАРЕЙ У ВИЧ-ИНФИЦИРОВАННЫХ ДЕТЕЙ. Journal of cardiorespiratory research, 2 (3), 31-34. doi: 10.26739/2181