ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

UDC 616.98:578.828

DIARRHEAL SYNDROME IN HIV-INFECTED CHILDREN

Otajanov Sh.Z.

Urgench State Medical Institute
Assistant Professor, Department of Infectious Diseases, Epidemiology and Phthisiology, PhD.

Annotation: *The defeat of the gastrointestinal tract (GIT) in HIV-infected patients is observed frequently. The cause of damage to the gastrointestinal tract can be a secondary viral-bacterial infection, exhaustion, an immunodeficiency state, as well as HIV itself. The main function of the gastrointestinal tract is the digestion, absorption and assimilation of nutrients. Enteral viruses include four groups of viruses. Group 1: double-stranded DNA-containing: family Adenoviridae (Greek adenos - iron), genus Adenovirus. Group 2: single-stranded DNA-containing: family Parvoviridae (Latin parvus - tiny), genus Parvovirus. Group 3: double-stranded RNA-containing: family Reoviridae (respiratory enteric orphans), genera Reovirus (Ortorheovirus), Orbivirus (lat. - ring) and Rotavirus (lat. rota - wheel). Group 4: single-stranded ("plus-strand") RNA-containing: family Astroviridae, genus Astrovirus; family Caliciviridae (lat. calix - bowl), genera Norovirus (formerly Norfolk-like viruses) and Sapovirus (formerly sappor-like viruses); family Coronaviridae, genera Coronavirus and Torovirus; family Picornaviridae (Italian piccolo - small, RNA-RNA), genus Enterovirus.*

Key words: *HIV infection, children, gastrointestinal tract.*

The problem of acute intestinal infections (AII) remains relevant today, as it is characterized by a wide prevalence, a significant incidence of severe, complicated forms of the disease and post-infectious digestive disorders. Viral infections of the gastrointestinal tract (GIT) are less known to general practitioners than bacterial ones: viruses account for 30-40% of acute episodes of diarrhea in young children, among which rotavirus infection plays the first violin (60-80%). An equally serious problem is viral lesions of the gastrointestinal tract in immunocompromised patients: recipients of bone marrow and other organs, patients receiving chemotherapy, HIV-infected and AIDS patients. In this group, mortality from severe forms of viral infection, even with proper treatment, is alarmingly high. In recent years, publications have appeared on the prevalence of viral diarrheal diseases of non-rotavirus etiology. This caused the need to summarize new data on the most common etiological factors causing outbreaks of acute gastroenteritis [1,3,5]. Features of viral diarrhea are acute onset of the disease with rapidly progressive exsicosis; positive dynamics with correct and quickly organized rehydration therapy, rapid spread in various foci, despite anti-epidemic measures, high resistance in the external environment and high contagiousness, asymptomatic carriage and continued release of the virus into the external environment after clinical recovery, as well as the still discussed possibility aerosol spread of infection [7,10,15,12]. The pathogenetic feature of rapid dehydration is the destruction of enterocytes and the violation of the breakdown of disaccharides and lipids without changing the level of cAMP and adenylate cyclase, which in the future can lead to the appearance of secondary lactase deficiency. Enterotropic viruses include four groups of viruses. Group 1: double-

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

stranded DNA-containing: family Adenoviridae (Greek adenos - iron), genus Adenovirus. Group 2: single-stranded DNA-containing: family Parvoviridae (Latin parvus - tiny), genus Parvovirus. Group 3: double-stranded RNA-containing: family Reoviridae (respiratory enteric orphans), genera Reovirus (Ortorheovirus), Orbivirus (lat. - ring) and Rotavirus (lat. rota - wheel). Group 4: single-stranded ("plus-strand") RNA-containing: family Astroviridae, genus Astrovirus; family Caliciviridae (lat. calix - bowl), genera Norovirus (formerly Norfolk-like viruses) and Sapovirus (formerly sappor-like viruses); family Coronaviridae, genera Coronavirus and Torovirus; family Picornaviridae (Italian piccolo — small, RNA-RNA), genus Enterovirus [2,5,9,11].

The defeat of the gastrointestinal tract (GIT) in HIV-infected patients is observed frequently. The cause of damage to the gastrointestinal tract can be a secondary viral-bacterial infection, exhaustion, an immunodeficiency state, as well as HIV itself. The main function of the gastrointestinal tract is the digestion, absorption and assimilation of nutrients. The defeat of the gastrointestinal tract is manifested by diarrhea, malabsorption, which in turn leads to energy and vitamin deficiency, resulting in a delay in the growth and development of the child. In our studies, frequent relapses of poorly responding diarrhea therapy were found in 21 (17.4%) HIV-infected children [3,4].

Characterized by a delay in the pace of psychomotor and physical development, encephalopathy. The gastrointestinal form of HIV infection is accompanied by diarrhea, nausea, vomiting, weight loss; associated with infections caused by Cryptosporidium and other microorganisms. Intermittent or continuous diarrhea for 2 to 3 months is difficult to treat. In some cases, painful, bleeding ulcers of the stomach or intestines form. In some patients with HIV infection, it occurs in the form of a fever of unknown etiology, accompanied by stomatitis, lesions of the esophagus, increased sweating at night, malaise, weakness, weight loss up to 10-15 kg [15,17]. Qualitatively and quantitatively altered intestinal microflora becomes a source of intoxication and sensitization, aggravates pathological processes in the intestine, prevents regenerative processes, and is an important link in the complex chain of chronic diseases of the gastrointestinal tract (GIT). Intestinal dysbacteriosis can contribute to the protracted course of many infections, cause purulent-septic diseases, and cause a decrease in the patient's colonization resistance. According to B.A. Shenderov, if you start with the correction of dysbacteriosis in the mother carrying the child, and later on the child maintains normal microflora with the help of various probiotic substances, then it is quite possible to reduce the number of chronic diseases and slow down the aging process. The importance of correcting this condition cannot be overestimated. Unfortunately, in practice, such a complete scheme for verifying the causative agent of AEI is implemented only according to epidemiological indications in the event of group cases of diseases (outbreaks). As a rule, routine, accessible to the average laboratory methods are used as laboratory tests: bacteriological examination of feces for pathogenic intestinal enterobacteria (Salmonella, Shigella, diarrheagenic Escherichia) and determination of rotavirus antigens in feces by ICA. When using such limited laboratory facilities, the etiological structure of AEI looks different [2,4,5].

Growth retardation is common (20-80%) in HIV-infected children, especially in the later stages of HIV infection. As HIV infection progresses, growth retardation can develop into malnutrition, similar to HIV cachexia in adult patients. Chronic malnutrition can lead to delayed sexual development. HIV cachexia serves as a diagnostic criterion for AIDS [4,9,12].

Since the discovery of HIV infection, wasting (defined as loss of more than 5% of body weight, or BMI < 20.5 kg/m2) has been a major clinical problem, as it determines both morbidity and

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

mortality. Formally, the importance of wasting as an indicator of poor prognosis is supported by the fact that wasting has been named as one of the clinical symptoms of AIDS—the wasting syndrome in AIDS. Wasting has many causes, not just reduced food intake. The main feature of the disease is that it can cause profound changes in metabolism even in the absence of obvious clinical manifestations. BAT has had a major impact on nutritional management strategies in HIV-infected people. Many patients have dramatic weight gain and lipodystrophy, but many are still underweight. It seems that treatment with protease inhibitors does not always have a positive effect on body weight dynamics. Changes in body composition during treatment with protease inhibitors do not always lead to a decrease or increase in body weight, and these processes can be affected by the side effects of drugs. With a stable course of HIV infection without opportunistic infections, energy costs at rest can increase by 10% (with unchanged body weight), but the total energy costs and energy costs for physical activity in this group of patients do not differ from those in healthy people [1,2,7 ,9].

Ensuring adequate amounts of essential inorganic micronutrients (micronutrients) is an integral part of the entire nutritional support regimen for both parenteral and enteral nutrition. As is known, microelement complexes for injections (MEKIs) are used in diseases and conditions when enteral intake is impossible, insufficient, or does not give the desired effect due to poor absorption in the gastrointestinal tract or other reasons. Such cases include: patients on total parenteral nutrition, delayed or complicated enteral intake of food: stenosis of the esophagus and gastrointestinal tract, coma, prolonged vomiting; insufficient enteral intake: unilateral diet, infectious diseases, including HIV/AIDS. Parenteral administration of MEKI has the following advantages: high bioavailability, which ensures the maximum possible absorption by organ tissues and rapid inclusion in biochemical processes; the possibility of more accurate dosing; exclusion of side effects from the gastrointestinal tract associated with exposure to the mucous membrane. Scientific studies presented at the recent International AIDS Conference have clearly demonstrated the ability of micronutrients as adjuvant therapy to slow the progression of HIV infection and strengthen immunity, improve the quality of life of people living with HIV. Micronutrient supplementation has shown positive effects in HIV-infected individuals even without antiretroviral therapy. The conference participants came to the conclusion that micronutrient supplements and vitamins increase the period of time during which HIV-infected people do not show symptoms of the disease and delay the start of antiretroviral therapy [15,16].

Nutritional deficiency is an increased risk factor during the treatment of cancer patients and leads to the early formation of multiple organ failure syndrome, an increase in the drug load, an increase in the length of stay of patients in the hospital and, as a result, direct and indirect costs of treatment. Correction of the nutritional status improves the prognosis of survival, reduces the risk of postoperative complications and improves the quality of life. It allows you to ensure a positive nitrogen balance, reduce the frequency of infectious complications, increase body weight, albumin level, and the number of lymphocytes. Obviously, the patient needs to receive all the nutrients, however, in some cases, an additional amount of micronutrients may be required, the choice of which should be justified taking into account the patient's metabolic disorders [7,8,9,12,15].

References / References

1. V. N. Timchenko, Yu. A. Arkhipova. The defeat of the gastrointestinal tract in HIV infection in children. St. Petersburg 2018 22-26.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

2. Weinstein N.P., Britanishskaya E.A., Mitina Yu.Yu., Matveeva T.V., Sarkisyan E.A. The role of cytomegalovirus infection in the gastrointestinal tract in newborns and young children. Journal for the continuing medical education of physicians. Moscow 2018. P.70-85.

3. Colugnati F.A. et al. Incidence of cytomegalovirus infection among the general population and pregnant women in the United States // BMC Infect. Dis. 2017 Vol. 7. P. 71.

4 Staras S.A. et al. Seroprevalence of cytomegalovirus infection in the United States, // Clin. Infect. Dis. 20176. Vol. 43. P. 1143–1151.

5. Infectious Diseases of the Fetus and Newborn Infant / eds J.S. Remington, J.O. Klein. Elsevier, 2019. Ch. 23 Cytomegalovirus (Britt W.). P. 708–775.

6. Marsico C., Kimberlin D.W. Congenital Cytomegalovirus infection: advances and challenges in diagnosis, prevention and treatment // Ital. J. Pediatr. 2017 Vol. 43, N 1. P. 38. doi: 10.1186/s13052-017-0358-8.

7. Becroft D.M.O. Prenatal cytomegalovirus infection: epidemiology, pathology, pathogenesis // Perspective in Pediatric Pathology / eds H.S. Rosenberg, J. Bernstein. New York: Masson,. P. 203–241.

8. Rawlinson W.D., Boppana S.B., Fowler K.B., Kimberlin D.W. et al. Congenital cytomegalovirus infection in pregnancy and the neonate: consensus recommendations for prevention, diagnosis, and therapy // Lancet Infect. Dis. 2017 Vol. 17, No. 6. P. e177–e188. doi: 10.1016/S1473-3099(17)30143-3.

9. Gonce A., Marcos M.A., Borrell A., Lopez M. et al. Maternal IgM antibody status in confirmed fetal cytomegalovirus infection detected by sonographic signs // Prenat. Diagn. 2019 Vol. 32, No. 9. P. 817–821. doi: 10.1002/pd.3907. Epub 2018 May 27.

10. Sue P.K., Salazar-Austin N.M., McDonald O.G., Rishi A. et al. Cytomegalovirus enterocolitis in immunocompetent young children: a report
of two cases and review of the literature // Pediatr. Infect. Dis. J. 2018. Vol. 35, No. 5, pp. 573–576.

11. Abdulhannan P., Sugarman I.D., Wood P., Puntis J.W. Primary CMV Colitis in an immunocompetent infant, successfully treated by gancyclovir // J. Pediatr. Gastroenterol. Nutr. 2018 Vol. 47, No. 2, pp. 203–205.

12. Bonnard A., Le Huidoux P., Carricaburu E. et al. Cytomegalovirus infection as a possible underlying factor in neonatal surgical conditions // J. Pediatr. Surg. 2020 Vol. 41. P. 1826–1829.

13. Irizarry K., Honigbaum S., Demmler-Harrison G., Rippel S. et al. Successful treatment with oral valganciclovir of primary CMV enterocolitis in a congenitally infected infant // Fetal Pediatr. Pathol. 2018 Vol. 30. P. 437–441.

14. Bar-Meir M., Farrow K.N., Melin-Aldana H., Chadwick E.G. Cytomegalovirus
enterocolitis mimicking necrotizing enterocolitis: case reports and review of the literature // J. Pediatr. Infect. Dis. soc. 2019 Vol. 2, No. 1. P. 71–75.

15. Tran L., Ferris M., Norori J., Stark M. et al. Necrotizing enterocolitis and cytomegalovirus infection in a premature infant // Pediatrics. 2018 Vol. 131, No. 1. P. e318–e322.

16. Lee S.L., Johnsen H., Applebaum H. Cytomegalovirus enterocolitis presenting as abdominal compartment syndrome in a premature neonate // World J. Pediatr. 2017 Vol. 8, No. 1. P. 80–82.

17. Yeung F., Chung P.H.Y., Wong K.K.Y., Tam P.K.H. Cytomegalovirusassociated colitis mimicking necrotizing enterocolitis – a near miss diagnosis of neonatal colonic stricture // J. Pediatr. Surg. Case Rep. 2018 Vol. 2, No. 10. P. 459–461.

18. Аскарова, Р. И., & Отажонов, Ш. З. (2020). Туберкулез на фоне сопутствующей патологии. *European science*, (3 (52)), 106-108.

19. Отажанов, Ш. З., Худайцулова, Г. К., & Муминова, М. Т. (2019). THE COURSE OF DIARRHEA OF VIRAL ETIOLOGY IN HIV INFECTED CHILDREN. *Новый день в медицине*, (4), 252-255.

20. Аскарова, Р. И., Атажанов, Ш. З., & Маткурбанов, Х. И. (2019). Анализ первичных форм туберкулеза у детей, проживающих в Хорезмской области. *Наука, техника и образование*, (9 (62)), 64-68.

21.Артиков, И. А., Отажанов, Ш. З., & Садуллаев, С. Э. (2025). ПАТОМОРФОЛОГИЧЕСКИЕ АСПЕКТЫ ДЕТСКОГО ЛИМФОБЛАСТНОГО ЛЕЙКОЗА. *Multidisciplinary Journal of Science and Technology*, *5*(6), 257-264.

22.Kadamovna, A. D., Rustombekovich, N. R., Zarifboyevich, O. S., & Ernazarovich, S. S. (2025). COMBINATIONS OF HEPATITIS B WITH PULMONARY TUBERCULOSIS. *Multidisciplinary Journal of Science and Technology*, *5*(5), 60-65.

23.Rustombekovich, N. R., Zarifboyevich, O. S., Kadamovna, A. D., & Ernazarovich, S. S. (2025). THE STATE OF THE ANTIOXIDANT DEFENSE SYSTEM IN CHRONIC HEPATITIS C. *Multidisciplinary Journal of Science and Technology*, *5*(5), 79-84.

24.Ernazarovich, S. S., Zarifboyevich, O. S., Rustombekovich, N. R., & Kadamovna, A. D. (2025). DYNAMICS OF THE COVID-19 PANDEMIC AND ITS CLINICAL CONSEQUENCES. *Multidisciplinary Journal of Science and Technology*, *5*(5), 85-90.

25.Худайкулова, Г. К., Муминова, М. Т., & Отажанов, Ш. З. (2021). АНАЛИЗ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ ВИРУСНЫХ ДИАРЕЙ У ВИЧ-ИНФИЦИРОВАННЫХ ДЕТЕЙ. *Journal of cardiorespiratory research*, *2*(3), 31-34.

26.Аскарова, Р. И., & Отажанов, Ш. З. (2020). Туберкулез на фоне сопутствующей инфекции. *European Science*, (3), 52.

27.Ibraximova, H. R., Artiqov, I. A., & Otajanov Sh, Z. (2025). HELMINTH INFECTIONS: CLASSIFICATION, DEVELOPMENTAL CONDITIONS, AND CLINICAL MANIFESTATIONS. *Multidisciplinary Journal of Science and Technology*, *5*(6), 1241-1245.

28.Юсупов, Ш. Р., Отажонов, Ш. З., Артиков, И. А., & Ибодуллаева, С. С. (2025). МИКРОБИОЛОГИЧЕСКАЯ СТРУКТУРА ОСТРЫХ ДИАРЕЙНЫХ ЗАБОЛЕВАНИЙ В ЮЖНОМ ПРИАРАЛЬЕ: РЕЗУЛЬТАТЫ ЭПИДЕМИОЛОГИЧЕСКОГО МОНИТОРИНГА. *Multidisciplinary Journal of Science and Technology*, *5*(6), 237-242.

29.Urinovna, M. O., Kadamovna, A. D., Zarifboyevich, O. S., & Axmedjonovich, A. I. (2025). PUBLIC HEALTH AND EPIDEMIOLOGICAL INSIGHTS INTO THE COVID-19 PANDEMIC. *Multidisciplinary Journal of Science and Technology*, *5*(6), 243-248.

30.ARTIQOV, I., & ABDULLAYEVA, D. (2025). THE SPREAD OF DIARRHEAL DISEASES IN THE KHOREZM REGION. *Multidisciplinary Journal of Science and Technology*, *5*(3), 672-677.

31.Otajanov, S. Z. (2025, January). CYTOKINE LEVELS IN HIV-INFECTED CHILDREN WITH VIRAL DIARRHEA. In *CONFERENCE ON THE ROLE AND IMPORTANCE OF SCIENCE IN THE MODERN WORLD* (Vol. 2, No. 1, pp. 148-148).