ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

UDC: 633.511;631.542.4.

INFLUENCE OF DEFOLIANTS ON THE OPENING OF BOLLS OF COTTON VARIETIES

*__Убайдуллаев Мадаминжон Муминжонович__, доктор философии по сельскохозяйственных наук, доцент. E-mail: mubaydullaev6554@gmail.com
тел: 90 274-65-54
**__Мамасолиев Зафарбек Кимсанбоевич__, доктор философии по сельскохозяйственных наук
**__Эргашев Нодирбек Юлдашалиевич__, доктор философии по сельскохозяйственных наук
***
*Ферганский государственный технический университет
**Ферганской научно-опытной станции Научно-исследовательского института селекции, семеноводства и агротехнологии выращивания хлопка
***
*Ubaydullaev Madaminjon Mominjonovich, doctor of Philosophy in Agriculture associate professor
**Mamasoliev Zafarbek Kimsanboyevich, doctor of Philosophy in Agriculture
**Ergashev Nodirbek Yuldashaliyevich, doctor of Philosophy in Agriculture
***
*Fergana State Technical University.
* Fergana Scientific and Experimental Station of the Cotton Breeding, Seed Production, and Agrotechnology Research Institute

Аннотация: В вариантах 16-18 с применением смеси Хим-экстра + Этифон в норме 0,100+1,5 л/га, 0,150+1,0 л/га и 0,200+0,5 л/га относительно высокие показатели раскрытых коробочек составили 97,5%, полураскрытых 0,5%, скорость раскрытия 42,4% и разница от контроля 25,7%.

Ключевые слова: состояние почвы, слой, рост, ветвление, разветвление, точка роста.

Abstract: In variants 16-18 with the application of mixtures of Chem-extra + Etifon at a rate of 0.100+1.5 l/ha, 0.150+1.0 l/ha, and 0.200+0.5 l/ha, the relatively high number of opened bolls was 97.5%, half-opened bolls - 0.5%, the opening rate - 42.4%, and the difference from the control was 25.7%, which was obtained when using Chem-extra at a rate of 0.100 l/ha + Etifon at a rate of 1.5 l/ha.

Key words: soil condition, layer, growth, branching, branching, growth point.

Introduction. It can be said that the opening of cotton bolls depends on a number of external factors, as well as on the agrotechnical measures carried out, as well as the timing and optimal rates of defoliation [2].

For defoliation, chemical substances - defoliants - are used, which cause processes similar to aging or natural shedding of plant leaves, as a result of which the opening of cotton bolls is accelerated by 15-20% [8].

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

Research methodology. Based on the above urgent tasks, during 2018-2022, our research on the topic was conducted in the conditions of meadow-sodic, mechanically heavy loamy, slightly saline soils of the scientific experimental station of the Research Institute of Cotton Breeding, Seed Production and Growing Agrotechnologies, located in the Kuva district of the Fergana region, with a groundwater depth of 1.6-1.8 meters [5].

Research results. Experiments 1-2, conducted in the studies, were also carried out in 2018-2020, and under the conditions of 2020, under the control of variants (1-6), where the application of defoliants was prescribed for the C-8290 cotton variety with 30-40% boll opening, initially 33.8% of the bolls opened, after 14 days this indicator was 65.7%, the half-opened bolls were 18%, and the opening rate was 31.9%. This last indicator differs by 0.6% from the parallel variant of the C-8290 variety in the 1st experiment, since all cotton varieties were grown in the same climatic conditions of 2020, moreover, on the same plot. Liquid XMD was applied at a rate of 8.0 l/ha. In the 2nd variant, the above indicators were 78.9; 1.7; 44.4%, which is 12.5% higher than the control.

In the 3rd variant with the use of Kim-extra as a standard at a rate of 0.200 l/ha, the initial opening of 35.1% of bolls was observed, and after 14 days of defoliation it was 81.4%, while the half-opened bolls were 1.6%, the opening rate was 45.5%, and the difference from the control was 13.4%. The effect of this standard was 2.5; 0.9 and 0.9%, respectively. Consequently, good results are also obtained with the application of the chemical extract itself, however, with the mixed application of the chemical extract at a rate of 0.100 l/ha + Etifon at a rate of 1.5 l/ha, the above indicators were respectively 93.2; 1.5; 58.7 and 26.8%, respectively, the number of opened bolls was 11.8%, the opening rate was 13.4%, and compared to the control variant, the number of opened bolls was 27.5%, and the opening rate was 26.8% higher compared to the application of only Khim-extra at a rate of 0.200 l/ha.



Figure 2. Difference of defoliated area from control, day 7

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11

When using the chemical preparation "Chim-extra" at a rate of 0.150 l/ha + Etifon 1.0 l/ha (5 variants) with 30-40% boll opening, the number of opened bolls was 94.5%, the number of half-opened bolls was 1.7%, the opening rate was 60.7%, and the difference from the control was 28.8%, moreover, under the influence of the standard chemical preparation "Chim-extra" at a rate of 0.200 l/ha, these indicators increased by 13.1 and 15.4%, respectively. With the application of Xim-extra at a rate of 0.200 l/ha + Etifon at a rate of 0.5 l/ha, the above indicators decreased slightly, the number of opened bolls decreased by 6.1% compared to the optimal variant, and the opening rate decreased by 6.4%. In the control variants (13-18), where the application of defoliants was prescribed with 50-60% boll opening of this cotton variety, 55.1% of bolls opened initially, and after 14 days this indicator naturally reached 71.8%, and the opening rate was 16.7%.

Whereas in the parallel variant (17) of the 1st experiment, these indicators were 56.2; 73.1 and 16.9%, respectively; 1.3 and 0.2% respectively. The correlation between leaf fall and boll opening was high and was equal to r=0.905.




Figure 2. Influence of defoliants on the opening of bolls of the C-8290 cotton variety, 2020, Experiment 2

In the 2nd experiment, defoliation of the Namangan-77 variety was carried out with 30-40% boll opening; in the control variants (7-12), the number of bolls opened naturally initially was 34.1%, after 14 days it was 65.1%, the number of half-opened bolls was 2.1%, the opening rate was 31.0%, and compared to the parallel variant (1), the number of opened bolls of the C-8290 variety was 0.6%. When using liquid XMD at a rate of 8.0 l/ha for this cotton variety, the number of opened bolls was 0.5%, the number of half-opened bolls was 0.3%, and the opening rate was 0.2%, which differed from the control by 1.6%.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-11




Figure 3. Influence of defoliants on the opening of bolls of the Namangan-77 cotton variety,2020, Experiment 2

With the application of chemical extra at a rate of 0.200 l/ha and as a standard (variant 9), the number of opened bolls was 80.1%, half-opened bolls 2.3%, the opening rate was 45.0%, and the difference from the control was 14.0%. In the experiment, compared to the tested norms of Chimextra and the standard at a rate of 0.100+1.5 l/ha, 0.150+1.0 l/ha, and 0.200+0.5 l/ha, relatively high indicators were obtained at a rate of 0.150+1.0 l/ha, and the above indicators were respectively 94.5; 1.7; 57.4 and 26.4%, respectively.

Conclusion

In conclusion, it can be said that among the mixtures of defoliants Khim-extra+Ethifon, relatively high indicators were obtained in both varieties at a boll opening rate of 50-60% at a rate of 0.100+1.5 l/ha, and the number of opened bolls was 97.5-98.4%, respectively.

Thus, it was established that the optimal effect of the defoliants used in the conducted experiments 1-2 was relatively high at 30-40% boll opening and lower at 50-60% boll opening.

References

1. Madaminjon Ubaydullaev,Sultanbek Allanazarov, Jasurbek Komilov. Effect of Weeding on Cotton Weight Per Boll and Cotton Yield Depending on Seedling Thickness of Medium Fiber Cotton Varieties BIO Web of Conferences 78, 03012 (2023)

https://www.bioconferences.org/articles/bioconf/abs/2023/23/bioconf_mtsitvw2023_03012/bioconf_mtsitvw2023_03012.html

2. Ubaydullayev, M., & Kurbanova, U. (2023). THE INFLUENCE OF DEFOLIANTS ON THE TECHNOLOGICAL QUALITY INDICATORS AND CHEMICAL COMPOSITION OF SEED. Science and innovation, 2(D4), 26-30.

3. Убайдуллаев, М. М., & Саетбековна, Қ. У. (2023). ДЕФОЛИАЦИЯ ЎТКАЗИШ УЧУН ДАЛАЛАРНИ ТАНЛАШ ВА ТАЙЁРЛАШ. Journal of Science-Innovative Research in Uzbekistan, 1(9), 322-328.

4. Mo‘minjonovich, U. M., & Saetbekovna, Q. U. (2023). EFFECT OF DEFOLIANTS ON COTTON WEIGHT. Journal of Science-Innovative Research in Uzbekistan, 1(9), 316-321.

5. Ubaydullayev, M. M. (2022). YANGI DEFOLIANTLAR HOSILDORLIK GAROVI. Архив научных исследований, 2(1).

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

6.Mo'minjonovich, U. M. (2022). Effectiveness Of Defoliants. Eurasian Research Bulletin, 8, 9-

7.Mominjonovich, U. M., & Ogli, M. I. V. (2022). STUDY AND ANALYSIS OF TECHNOLOGICAL PROCESSES OF COTTON DRYING IN A CLUSTER SYSTEM. International Journal of Advance Scientific Research, 2(11), 6-10.

8.Ubaydullaev, M. M., & UT, T. (2022). DETERMINATION OF APPROPRIATE NORMS AND TERMS OF DEFOLIANTS. American Journal Of Applied Science And Technology, 2(05), 18-22.

9.Ubaydullaev, M. M., & Makhmudova, G. O. (2022). MEDIUM FIBER S-8290 AND S-6775 COTTON AGROTECHNICS OF SOWING VARIETIES. European International Journal of Multidisciplinary Research and Management Studies, 2(05), 49-54.

10.Ubaydullaev, M. M., & Komilov, J. N. (2022). EFFECT OF DEFOLIANTS FOR MEDIUM FIBER COTTON. International Journal of Advance Scientific Research, 2(05), 1-5.

11.Ubaydullaev, M. M., & Mahmutaliyev, I. V. (2022). EFFECTIVENESS OF FOREIGN AND LOCAL DEFOLIANTS ON THE OPENING OF CUPS. International Journal of Advance Scientific Research, 2(05), 6-12.

12.Ubaydullaev, M. M., & Sultonov, S. T. (2022). DEFOLIATION IS AN IMPORTANT MEASURE. European International Journal of Multidisciplinary Research and Management Studies, 2(05), 44-48.

13.Ubaydullaev, M. M., & Nishonov, I. A. (2022). The Benefits of Defoliation. Eurasian Journal of Engineering and Technology, 6, 102-105.

14.Ubaydullayev Madaminjon Mo'minjon o'g'li, & Ma'rufjonov Abdurahmon Mo'sinjon o'g'li. (2022). BIOLOGICAL EFFICIENCY OF FOREIGN AND LOCAL DEFOLIANTS. "science and innovation" international scientific journal, 1(2). https://doi.org/10.5281/zenodo.6569808

15.Ubaydullayev, M. M. (2021). G'o'zada defoliatsiya o'tkazishning maqbul me'yor va muddatlari. Monografiya.-Corresponding standards and terms of defliation of cotton. Monograph. Соответствующие нормы и сроки дефолиации хлопка. Монография. Zenodo.