ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

LABORATORY PREDICTORS OF SEVERE COVID-19: A SCIENTIFIC AND ANALYTICAL REVIEW OF RETROSPECTIVE STUDY DATA

Alimova Umidakhon Omonjon qizi,
Tashkent State Medical University, Uzbekistan

Abstract: During the COVID-19 pandemic, laboratory biomarkers have become essential tools for evaluating disease severity, predicting complications, and determining the need for intensive therapy. This article presents a scientific and analytical synthesis of retrospective data from 350 hospitalized patients with PCR-confirmed SARS-CoV-2 infection.

Key laboratory indicators with high prognostic significance were identified: lymphopenia, neutrophil-to-lymphocyte ratio (N/L), C-reactive protein (CRP), lactate dehydrogenase (LDH), fibrinogen, D-dimer, and platelet parameters. The study established that combined immuno-inflammatory and coagulopathic abnormalities shape an integral pathogenetic profile of severe COVID-19. Particular attention is given to the U-shaped relationship between fibrinogen levels (<2.0 g/L or >9.0 g/L) and mortality risk, as well as the strong association of elevated D-dimer with thrombotic complications and multi-organ failure.

These findings underscore the necessity of continuous laboratory monitoring and highlight the importance of biomarker-based personalized patient management.

Keywords: COVID-19; laboratory biomarkers; fibrinogen; D-dimer; lymphopenia; N/L ratio; CRP; LDH; cytokine storm; coagulopathy; immunothrombosis; severity prediction.

Introduction

The COVID-19 pandemic has created an unprecedented need for laboratory indicators that can reliably assess disease severity and predict clinical outcomes. Data from the dissertation research *(Alimova, 2022)* clearly demonstrate that COVID-19 is a multisystem disorder characterized by immune dysregulation, lung tissue injury, and profound thrombo-inflammatory disturbances. International studies have confirmed that laboratory biomarkers may surpass imaging modalities in diagnostic sensitivity, particularly in the early stages of disease *(Guan et al., 2020; Huang et al., 2020)*.

The analyzed cohort included 350 patients with PCR-confirmed SARS-CoV-2 infection. Laboratory markers of inflammation, immune response, and hemostasis were examined: lymphocyte count, neutrophil-to-lymphocyte ratio (N/L), CRP, LDH, ferritin, fibrinogen, D-dimer, and platelet indices. This integrated approach allowed the identification of dynamic patterns and laboratory criteria associated with unfavorable outcomes.

Immune Dysfunction: Lymphocytes, N/L Ratio, CRP, and LDH as Composite Markers

One of the most significant findings in COVID-19 is **lymphopenia**. More than 80% of patients demonstrated reduced lymphocyte counts, with severe cases showing values as low as 0.3–0.5×10⁹/L, indicating marked suppression of adaptive immunity *(Alimova, 2022)*.

The **neutrophil-to-lymphocyte ratio (N/L)** serves as an even more precise marker of severity. Values exceeding 6 were strongly predictive of cytokine storm progression. The correlation between N/L and CT-verified lung involvement reached r=0.823, demonstrating its high prognostic accuracy.

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

C-reactive protein (CRP), a classical acute-phase reactant, also exhibited strong prognostic relevance. Levels exceeding 80–100 mg/L were closely associated with rapid development of respiratory failure.

Lactate dehydrogenase (LDH), reflecting cellular injury, emerged as a sensitive predictor of lung tissue damage. Values above 500 U/L consistently correlated with unfavorable outcomes *(Wang et al., 2020)*.

The combined pattern of lymphopenia, elevated N/L, CRP, and LDH constitutes an "immuno-inflammatory profile" capable of predicting disease deterioration prior to clinical decompensation.

Fibrinogen and D-dimer as Key Indicators of COVID-19-Associated Coagulopathy

COVID-19 induces significant hemostatic impairment characterized by endothelial injury, microvascular thrombosis, and activation of fibrinolysis. The dissertation revealed a unique and clinically important phenomenon — a **U-shaped association between fibrinogen levels and mortality risk**.

Fibrinogen as a Bidirectional Predictor

• **Low fibrinogen (<2.0 g/L)** reflects consumption coagulopathy and the onset of disseminated intravascular coagulation (DIC), with a 3.2-fold increase in mortality risk.

• **High fibrinogen (>9.0 g/L)** indicates severe hypercoagulability and is associated with a 2.6-fold increase in fatal outcomes *(Alimova, 2022)*.

These findings illustrate that fibrinogen is not merely an inflammatory marker but a dual-phase indicator of hemostatic imbalance.

D-dimer as the Most Universal Predictor of Poor Outcome

Elevated D-dimer (>1–2 mg/L) was strongly associated with the development of ARDS, pulmonary embolism, and mortality *(Tang et al., 2020)*. In severe cases, rising D-dimer values were the earliest and most reliable indicators of "immunothrombosis" and impending organ failure.

The combination of **fibrinogen and D-dimer** provides a powerful framework for forecasting:

• risk of thrombotic complications,

• likelihood of DIC,

• need for escalation of anticoagulant therapy.

Platelet Dynamics and Integrative Coagulation Models

Platelet counts demonstrated a paradoxical pattern: while thrombocytopenia (110–150×10⁹/L) was common in severe cases, hypercoagulability persisted. This is explained by increased platelet consumption during microthrombus formation and possible bone marrow suppression.

The platelet-to-lymphocyte ratio (PLR) increased significantly in severe disease, reflecting simultaneous activation of inflammatory and thrombotic pathways.

Thus, platelet indices extend beyond classic hemostasis and may serve as additional markers of systemic inflammatory burden.

An Integrative Pathogenetic Model of Severe COVID-19

Based on the analytical synthesis of laboratory findings, three key pathogenetic circuits can be identified:

1. Immune Dysregulation

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

Marked by lymphopenia, elevated N/L ratio, and increased CRP. Leads to cytokine storm and immune collapse.

2. Tissue Injury

Reflected by elevated LDH and ferritin. Associated with necrotizing lung damage and systemic hypoxia.

3. Thrombo-inflammatory Coagulopathy

Characterized by abnormal fibrinogen levels, increased D-dimer, and reduced platelet counts. Drives microthrombosis, pulmonary embolism, DIC, and multiorgan failure.These circuits reinforce one another, explaining the rapid deterioration observed in a subset of patients.

Conclusion

Laboratory biomarkers represent a cornerstone in the prognostic evaluation of COVID-19. Data from the retrospective study confirm that:

- lymphopenia and elevated N/L ratio are early predictors of severe disease;
- CRP and LDH are reliable indicators of systemic inflammation and tissue injury;
- fibrinogen and D-dimer are the most critical markers of thrombo-inflammatory processes and mortality risk.

The integration of these laboratory parameters supports more accurate risk stratification and enhances personalized management strategies for patients with COVID-19.

References

1. Alimova, U. O. O. (2022). *COVID-19 bilan kasallangan bemorlarda laborator ko‘rsatkichlar dinamikasining retrospektiv tahlili* [Master’s thesis, Tashkent State Medical Academy].

2. Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., … Zhang, L. (2020). Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study. *The Lancet, 395*(10223), 507–513. https://doi.org/10.1016/S0140-6736(20)30211-7

3. Guan, W., Ni, Z., Hu, Y., Liang, W., Ou, C., He, J., … Zhong, N. (2020). Clinical characteristics of coronavirus disease 2019 in China. *The New England Journal of Medicine, 382*(18), 1708–1720. https://doi.org/10.1056/NEJMoa2002032

4. Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., … Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet, 395*(10223), 497–506. https://doi.org/10.1016/S0140-6736(20)30183-5

5. Tang, N., Bai, H., Chen, X., Gong, J., Li, D., & Sun, Z. (2020). Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *Journal of Thrombosis and Haemostasis, 18*(4), 844–847. https://doi.org/10.1111/jth.14768

6. Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., … Peng, Z. (2020). Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. *JAMA, 323*(11), 1061–1069. https://doi.org/10.1001/jama.2020.1585

7. Zhang, J., Wang, X., Jia, X., Li, J., Hu, K., Chen, G., … Dong, W. (2020). Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clinical Microbiology and Infection, 26*(6), 767–772. https://doi.org/10.1016/

8. Dekhkambayeva, Z. ., Rahimova, M. ., & Panabaeva , N. . (2025). EFFECTIVNES OF INTENSIFICATION AND ACCELERATION TECHNOLOGIES IN TEACHING CHRONIC

ISSN: 2582-4686 SJIF 2021-3.261, 2022-2.889, 2023-5.384, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-12

HEART FAILURE TO STUDENTS. International Journal of Medical Sciences, 1(4), 28+32. Retrieved from https://inlibrary.uz/index.php/ijms/article/view/97134

9. Dekhkambayeva , Z., Nishanova, Y., & Toshboeva , N. (2025). IMPROVING EDUCATIONAL EFFECTIVENESS IN TEACHING RADIONUCLIDE-NEGATIVE BREAST CANCER THROUGH COMPREHENSIVE RADIODIAGNOSIS. International Journal of Medical Sciences, 1(4), 108–112. Retrieved from https://inlibrary.uz/index.php/ijms/article/view/97163

10. Xoltajiyev Xurshid Otabek o‘g‘li, & Dexkambayeva Z.A.,. (2025). GIPERTONIYA KASALLIGINI BOSHQARISHDA TIBBIY PEDAGOGIKANING AHAMIYATI: PEDAGOGIK MAHORATNING ROLI VA SAMARADORLIGI . Medicine, Pedagogy and Technology: Theory and Practice, 3(4), 6–11. Retrieved from https://inlibrary.uz/index.php/mpttp/article/view/84157

11. Декхкамбаева, З. (2024). Zamonaviy ta'limni jismoniy tarbiya va sport vositasida takomillashtirish. Наука и инновации, 1(1), 70-7

12. Zulfiya, D. (2022). SOG ‘LOM TURMUSH TARZINI SHAKLLANTIRISHDA SOG ‘LOMLASHTIRUVCI TEXNOLOGIYALAR. Новости образования: исследование в XXI веке, 1(5), 697-700.

13. 24. Dekhkambaeva Zulfia (20025) ANALYSIS OF CONCEPTS OF DEVELOPING THE CREATIVE ABILITY OF STUDENTS IN HIGHER EDUCATIONAL INSTITUTIONS. (2025). EduVision: Journal of Innovations in Pedagogy and Educational Advancements, 1(2), 292-296.

14. 21. Нарметова, Ю., & Дехкамбаева, З. (2024). ВЛИЯНИЕ НЕРЕШЁННЫХ ПСИХОЛОГИЧЕСКИХ КОНФЛИКТОВ И АЛЕКСИТИМИИ НА СОМАТИЧЕСКОЕ ЗДОРОВЬЕ.Multidisciplinary Journal of Science and Technology,4(12), 97-99.

15. NEUROSIS AND NEUROTIC DISORDERS. (2024). *Western European Journal of Medicine and Medical Science*, *2*(12), 28-30. https://westerneuropeanstudies.com/index.php/3/article/view/1833