ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

DETERMINATION OF SELECTED HEAVY METAL IONS IN FOOD PRODUCTS USING ADVANCED VOLTAMMETRIC ANALYSIS

Authors: Muhammad Rajabov[1], Sergey Aronbayev[1], Hasan Nasimov[1]
[1] Institute of Biochemistry, Samarkand State University, Samarkand, Uzbekistan

Abstract
Heavy metal contamination in food products poses a significant threat to public health due to their persistence, bioaccumulation, and toxicity. Industrial emissions, pesticide residues, and inadequate agricultural practices contribute to the presence of metals such as Pb^{2+}, Cd^{2+}, Cu^{2+}, and Zn^{2+} in commonly consumed foods. This study investigates the quantitative determination of heavy metal ions in wheat flour, rice flour, dried potatoes, and dried carrots using voltammetric analytical methods. Samples were digested using nitric acid and hydrogen peroxide, followed by precise electrochemical measurement. Analytical calibration, interference studies, and reproducibility assessments were conducted for method validation. Results demonstrated slightly elevated concentrations of Pb^{2+} and Cd^{2+} in selected samples, while Cu^{2+} and Zn^{2+} remained within acceptable regulatory limits. The findings highlight the necessity of routine monitoring and the suitability of voltammetric techniques for rapid, sensitive, and cost-effective heavy metal determination in food matrices.
Keywords: heavy metals, food safety, voltammetric analysis, Pb^{2+}, Cd^{2+}, Cu^{2+}, Zn^{2+}, contamination monitoring

1. Introduction

Food safety remains one of the most critical global challenges, especially in regions where industrialization and agricultural expansion have intensified environmental contamination. Heavy metals—particularly lead (Pb), cadmium (Cd), copper (Cu), and zinc (Zn)—are frequently detected in soil, water, and food products. These metals exhibit non-biodegradable behavior and accumulate in living organisms, resulting in severe health implications including neurotoxicity, kidney damage, metabolic disorders, carcinogenic effects, and impaired growth.

Lead exposure, even at trace levels, affects children's cognitive development and causes chronic cardiovascular damage, whereas cadmium is associated with renal dysfunction and bone fragility. Copper and zinc, although essential micronutrients, become toxic at elevated concentrations, disrupting enzymatic activity and metabolic pathways.

Electrochemical methods, particularly voltammetric techniques, have become dominant analytical tools for heavy metal detection due to their high sensitivity, selectivity, low detection limits, and minimal sample consumption. Compared with spectroscopic techniques (AAS, ICP-OES), voltammetry offers rapid analysis, cost efficiency, and portability, making it suitable for both laboratory and field applications.

This study aims to quantify Pb^{2+}, Cd^{2+}, Cu^{2+}, and Zn^{2+} ions in selected food products using advanced voltammetric analysis and evaluate their compliance with international food safety standards.

2. Materials and Methods

2.1 Sample Collection

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

Food samples were collected from local markets in Samarkand, Uzbekistan. The following materials were analyzed:

- Wheat flour
- Rice flour
- Dried potato samples
- Dried carrot samples

Each sample was homogenized and stored in airtight containers prior to digestion.

2.2 Wet Digestion Procedure

Approximately 1.0 g of each sample was transferred into Teflon digestion vessels. A mixture of:

- **8 mL HNO_3 (65%)**
- **2 mL H_2O_2 (30%)**

was added to each vessel.

Digestion was performed using a controlled temperature program until complete mineralization was achieved. The resulting clear solutions were diluted to 50 mL using deionized water.

2.3 Voltammetric Analysis

Electrochemical measurements were performed using a multi-mode voltammetric workstation equipped with:

- Working electrode: modified carbon electrode
- Reference electrode: Ag/AgCl
- Counter electrode: platinum wire

Calibration curves were constructed using standard solutions of Pb^{2+}, Cd^{2+}, Cu^{2+}, and Zn^{2+} at known concentrations. Analytical validation included:

- Limit of detection (LOD)
- Limit of quantification (LOQ)
- Linearity ($R^2 > 0.995$)
- Repeatability (RSD < 5%)
- Interference studies via standard addition

2.4 Statistical Analysis

All measurements were performed in triplicate. Data were analyzed using OriginPro and GraphPad Prism to generate statistical plots, bar charts, and concentration curves.

---

3. Results

3.1 Heavy Metal Concentrations

The concentrations of Pb^{2+}, Cd^{2+}, Cu^{2+}, and Zn^{2+} in the analyzed samples are shown below.

Sample	Pb^{2+} (mg/kg)	Cd^{2+} (mg/kg)	Cu^{2+} (mg/kg)	Zn^{2+} (mg/kg)
Wheat flour	0.15	0.12	0.05	0.20
Rice flour	0.12	0.09	0.06	0.18
Dried potatoes	0.18	0.14	0.07	0.22
Dried carrots	0.13	0.10	0.05	0.19

3.2 Comparison with Permissible Limits

Regulatory limits (WHO/FAO):

- Pb^{2+}: 0.10 mg/kg

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

- Cd^{2+}: 0.10 mg/kg
- Cu^{2+}: 0.20 mg/kg
- Zn^{2+}: 0.50 mg/kg

Pb^{2+} and Cd^{2+} slightly exceeded permissible limits in wheat flour and dried potatoes.

3.3 Graphical Analysis




Figure 1. Bar chart of Heavy Metal Concentration in Food Samples

Concentration levels of Pb^{2+}, Cd^{2+}, Cu^{2+}, and Zn^{2+} in wheat flour, rice flour, dried potatoes, and dried carrots.

The bar chart illustrates the measured concentrations of the four heavy metals across all food samples. Pb^{2+} and Cd^{2+} show noticeably higher levels in wheat flour and dried potatoes, while Cu^{2+} and Zn^{2+} remain relatively stable across samples.




Figure 2. Metal concentration curves vs. permissible levels

This graph presents the measured values plotted against international regulatory thresholds. Pb^{2+} and Cd^{2+} exceed the maximum allowable concentration (0.10 mg/kg) in wheat flour and dried

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

potatoes, whereas Cu^{2+} and Zn^{2+} remain below their respective limits (0.20 mg/kg and 0.50 mg/kg). The graphical comparison clearly visualizes the margin of exceedance and compliance.

Graphical Construction Details:

Both figures are constructed using the mean of triplicate measurements, with error bars representing standard deviations. Statistical processing and visualization were performed using OriginPro and GraphPad Prism, ensuring high-accuracy numerical rendering. All axes were calibrated with uniform scaling for comparison clarity.

4. Discussion

The presence of heavy metals, especially Pb^{2+} and Cd^{2+}, beyond regulatory limits raises substantial concerns regarding food safety. Wheat flour and dried potatoes showed the highest contamination, possibly due to soil composition, irrigation water, or environmental exposure during cultivation and storage.

Electrochemical analysis proved highly effective due to:

- Rapid detection
- Low sample volume
- High precision
- Minimal chemical consumption

Comparative analysis with international studies confirms similar contamination patterns in regions with agricultural-industrial overlap. The technique's reproducibility and sensitivity highlight its suitability for routine monitoring in developing countries where advanced spectrometric equipment may not be readily available.

5. Conclusion

1. Heavy metals were successfully quantified using voltammetric analysis.
2. Pb^{2+} and Cd^{2+} levels exceeded international safety limits in selected food samples.
3. Cu^{2+} and Zn^{2+} remained within acceptable ranges.
4. The voltammetric method proved reliable, sensitive, and practical for food safety monitoring.
5. These results emphasize the need for continuous surveillance and stricter regulatory control over food production chains.

References

1. Abdiganiev, D. *Ecological Chemistry*. Tashkent: Fan Publishing, 2019.
2. Karimov, Sh., Rakhmatova, M., & Mukhtorov, A. *Chemical Analysis Methods for Food Products*. Samarkand: SamSU Press, 2021.
3. Holmuradov, A. *Analytical Chemistry: Laboratory Manual*. Tashkent: University Press, 2018.
4. Sharipov, B. “Development and Monitoring of Heavy Metals in Various Environments.” *Uzbekistan Journal of Chemistry*, no. 4, 2022, pp. 45–52.
5. Wang, J. *Analytical Electrochemistry*. 4th ed. New Jersey: Wiley, 2020.
6. Skoog, D.A., Holler, F.J., & Crouch, S.R. *Principles of Instrumental Analysis*. 7th ed. Boston: Cengage Learning, 2018.
7. Jarup, L. “Hazards of Heavy Metal Contamination.” *British Medical Bulletin*, vol. 68, no. 1, 2003, pp. 167–182.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-12

8. WHO/FAO. *Joint Expert Committee on Food Additives: Heavy Metals in Food*. Rome: WHO Press, 2019.
9. Gumpu, M.B., Sethuraman, S., Krishnan, U.M., & Rayappan, J.B.B. "A Review on Electrochemical Sensors for the Detection of Heavy Metals." *Sensors and Actuators B: Chemical*, vol. 213, 2015, pp. 515–533.
10. Šestáková, J., & Mikelová, R. "Voltammetric Determination of Trace Metals in Food Matrices." *Food Chemistry*, vol. 135, 2012, pp. 1718–1725.
11. Ahmed, M. "Environmental Exposure to Heavy Metals in Edible Plants." *Environmental Monitoring and Assessment*, 2022.
12. Park, Y., & Lee, J. "Electrochemical Quantification of Pb^{2+} and Cd^{2+} in Agricultural Products." *Analytical Letters*, 2020, pp. 1920–1932.
13. Ismoilov, A. "A Voltammetric Approach for Determining Lead and Cadmium in Flour Samples." *International Conference on Modern Analytical Chemistry*, Tashkent, 2022.
14. Lee, H. "Nanostructured Electrodes for Heavy Metal Detection." *IEEE Sensors Conference Proceedings*, 2020.