ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-12

PERIODONTITIS AS A MANIFESTATION OF SYSTEMIC DISEASE

Gulmuxamedov P.B.
Tashkent State Medical University

Abstract

Chronic periodontitis is a multifactorial inflammatory disease characterized by progressive destruction of periodontal tissues and alveolar bone. While microbial biofilm initiates the inflammatory process, individual variability in disease progression suggests a significant contribution of host-related factors, particularly genetic determinants of immune response. Pro-inflammatory cytokines play a key role in regulating periodontal inflammation, and genetic polymorphisms affecting cytokine expression may influence disease severity and clinical outcomes.

The aim of this study was to evaluate molecular-genetic markers associated with the progression of chronic periodontitis and to assess their potential value for personalized periodontal risk prediction.

A total of 141 patients with chronic periodontitis and 138 periodontally healthy controls were examined. Patients were stratified according to disease severity into mild, moderate, and severe forms. Clinical periodontal parameters, radiographic bone loss, and microbiological characteristics were assessed. Genetic analysis focused on polymorphisms of pro-inflammatory cytokine genes TNF-α (rs1800629) and IL-1β (rs1143634, rs16944, rs1143627), identified using polymerase chain reaction techniques. Statistical associations between genetic variants and disease severity were analyzed using odds ratios and confidence intervals.

Specific polymorphic variants of TNF-α and IL-1β genes were significantly associated with increased risk of disease progression and severe clinical manifestations. Combined unfavorable genotypes demonstrated a cumulative effect, correlating with deeper periodontal pockets and more extensive alveolar bone resorption.

These findings support the role of molecular-genetic markers as predictors of chronic periodontitis progression and highlight their relevance for the development of personalized preventive and therapeutic strategies in periodontal care.

Keywords

chronic periodontitis; molecular genetics; cytokine polymorphism; TNF-α; IL-1β; disease progression; personalized dentistry

1. Introduction

Chronic periodontitis remains a major public health concern due to its high prevalence, progressive nature, and impact on oral and systemic health. The disease is initiated by pathogenic dental biofilm; however, the extent of periodontal tissue destruction varies widely among individuals with similar microbial exposure. This clinical heterogeneity cannot be explained solely by environmental factors and indicates the involvement of host-related biological mechanisms.

Inflammatory and immune responses play a central role in periodontal tissue breakdown. Pro-inflammatory cytokines, particularly tumor necrosis factor-alpha (TNF-α) and interleukin-1 beta (IL-1β), are key mediators that regulate leukocyte recruitment, connective tissue degradation, and osteoclast activation. Elevated levels of these cytokines have been consistently associated with active periodontal inflammation and bone resorption.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-12

Recent advances in molecular genetics have demonstrated that single nucleotide polymorphisms (SNPs) in cytokine genes may affect gene transcription and cytokine production. As a result, genetically determined differences in immune response intensity may predispose certain individuals to more aggressive forms of periodontitis.

The concept of personalized dentistry emphasizes the identification of individual risk factors, including genetic susceptibility, to optimize prevention, diagnosis, and treatment. However, the clinical application of molecular-genetic markers in periodontal practice remains limited, particularly in Central Asian populations.

The present study aimed to identify molecular-genetic predictors of chronic periodontitis progression by analyzing polymorphic variants of TNF-α and IL-1β genes and to evaluate their potential role in personalized periodontal care.

2. Materials and Methods

2.1. Study Population

The study included 141 patients diagnosed with chronic periodontitis and 138 periodontally healthy individuals serving as controls. All participants were ethnic Uzbeks to minimize population stratification effects. The study protocol complied with the principles of the Declaration of Helsinki, and informed consent was obtained from all participants.

2.2. Clinical and Radiographic Assessment

Comprehensive periodontal examination was performed, including evaluation of gingival inflammation, bleeding on probing, probing pocket depth, clinical attachment loss, and tooth mobility. Radiographic analysis was conducted to assess alveolar bone resorption. Based on clinical and radiographic findings, patients were categorized into mild, moderate, and severe chronic periodontitis groups.

2.3. Microbiological Evaluation

Subgingival samples were collected from periodontal pockets for microbiological analysis. The presence of aerobic and anaerobic pathogenic microorganisms was assessed, and their distribution was analyzed in relation to disease severity.

2.4. Molecular-Genetic Analysis

Genomic DNA was extracted from venous blood and saliva samples. Genotyping of TNF-α (rs1800629) and IL-1β (rs1143634, rs16944, rs1143627) gene polymorphisms was performed using polymerase chain reaction (PCR) techniques. Genotype distributions were evaluated for Hardy–Weinberg equilibrium.

2.5. Statistical Analysis

Statistical analysis included comparison of allele and genotype frequencies between groups using chi-square tests. Associations between genetic variants and disease progression were evaluated by calculating odds ratios (ORs) with 95% confidence intervals (CIs). A p-value < 0.05 was considered statistically significant.

3. Results

Analysis revealed that specific polymorphic variants of pro-inflammatory cytokine genes were significantly associated with disease progression. The TNF-α rs1800629 A allele was more frequently observed in patients with moderate and severe chronic periodontitis compared to those with mild disease and controls. Carriage of this allele was associated with increased probing pocket depth and greater radiographic bone loss.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

VOLUME-5, ISSUE-12

The IL-1β rs16944 T allele demonstrated a significant correlation with disease severity. Patients carrying unfavorable IL-1β genotypes were more likely to exhibit extensive periodontal tissue destruction and advanced clinical attachment loss.

Combined analysis of TNF-α and IL-1β polymorphisms showed that individuals with multiple unfavorable genotypes had the highest risk of rapid disease progression. These patients presented with deeper periodontal pockets, increased tooth mobility, and more pronounced alveolar bone resorption.

4. Discussion

The findings of this study support the concept that molecular-genetic factors play a crucial role in determining the progression of chronic periodontitis. Polymorphisms in genes encoding pro-inflammatory cytokines appear to modulate the host immune response, influencing both the intensity and duration of periodontal inflammation.

The cumulative effect of unfavorable TNF-α and IL-1β genotypes suggests that genetic risk assessment based on multiple markers may provide greater predictive value than single-gene analysis. This approach aligns with the principles of personalized dentistry and precision medicine.

Incorporating molecular-genetic screening into periodontal diagnostics may allow early identification of high-risk individuals and facilitate tailored preventive and therapeutic interventions. However, further studies with larger sample sizes and longitudinal designs are required to validate these findings and determine their clinical applicability.

5. Conclusions

Molecular-genetic polymorphisms of TNF-α and IL-1β genes are significantly associated with the progression and severity of chronic periodontitis. These genetic markers demonstrate potential value as predictors of disease progression and may serve as a foundation for personalized periodontal care strategies.

References

1. Budtz-Jørgensen, E.; Lombardi, T. Denture stomatitis revisited. *Periodontology 2000* **2018**, *78*, 254–261.
2. Felton, D.A. Complete edentulism and comorbid diseases: An update. *Journal of Prosthodontics* **2016**, *25*, 5–20.
3. Grigoriadou, M.E.; Koutayas, S.O.; Madianos, P.N.; Strub, J.R. Interleukin-1 as a genetic marker for periodontal disease. *Journal of Clinical Periodontology* **2010**, *37*, 517–523.
4. Napimoga, M.H.; Nunes, L.H.; Maciel, A.A.; et al. Possible involvement of IL-1β polymorphism in chronic periodontitis. *Journal of Oral Science* **2014**, *56*, 13–19.
5. Tian, Y.; Feng, P.; Wang, Y.; et al. Association between TNF-α gene polymorphism and chronic periodontitis: A meta-analysis. *Journal of Periodontal Research* **2013**, *48*, 298–305.
6. Ayazi, G.; Pirayesh, M. Genetic polymorphisms of cytokines and susceptibility to periodontal disease. *Journal of Dental Research, Dental Clinics, Dental Prospects* **2013**, *7*, 8–15.
7. Feng, P.; Tian, Y.; Chen, Z.; et al. Association of TNF-α polymorphisms with periodontal disease risk: Evidence from a population-based study. *Inflammation Research* **2014**, *63*, 901–908.
8. Casado, P.L.; Villas-Boas, R.; de Mello, W.; et al. Genetic susceptibility in chronic periodontitis: A systematic review. *Journal of Periodontology* **2015**, *86*, 99–111.
9. Petrin, A.N.; Tsarev, V.N. Cytokine gene polymorphisms in inflammatory diseases: Clinical significance. *Infection and Immunity* **2011**, *79*, 3381–3389.

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 9.948 / 2024

10. Gulmukhamedov, P.B. Clinical and molecular-genetic aspects of periodontitis pathogenesis. *PhD Thesis Abstract*, Tashkent State Dental Institute, Tashkent, Uzbekistan, **2019**.
11. Hasanov, L.E.; Abdullaeva, N.R.; Mirkhoshimova, M.F. Genetic markers of periodontal inflammation in Central Asian populations. *Stomatologiya* **2017**, *1*, 15–19.
12. Safarov, M.T.; Dadabaeva, M.U.; Asemova, S.A.; et al. Modern aspects of mathematic modeling in dental implantation. *Science and Education: Problems and Prospects* **2020**, 354–359.
13. Xabilov, N.L.; Xabilov, B.N.; Sharipov, S.S.; et al. Plant-based complexes in dentistry. *Journal of New Century Innovations* **2024**, *66*, 203–209.
14. World Health Organization. Oral health fact sheet. WHO, Geneva, Switzerland, **2012**. Available online: https://www.who.int (accessed on Day Month Year).