ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 10.25 / 2025

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-6, ISSUE-2

IRON DEFICIENCY ANEMIA IN CHILDREN AND EFFECTIVE DIAGNOSTIC AND TREATMENT PATHWAYS

Makhpieva Guldonakhon Kabiljanovna,
PhD, Associate Professor of the Department of pediatrics,
Faculty of Medicine, ASMI

Abstract. Iron deficiency anemia is the most common nutritional anemia in childhood and a frequent reason for pediatric visits, laboratory testing, and supplementation. Its causes are usually predictable—insufficient dietary iron intake, rapid growth, prematurity, and excessive cow's milk—yet it is often missed until hemoglobin falls substantially or developmental and behavioral concerns emerge. This article summarizes effective diagnostic and treatment pathways for iron deficiency anemia in children, with a practical emphasis on outpatient primary care.

Keywords: iron deficiency anemia; pediatrics; ferritin; hemoglobin; oral iron therapy; screening; cow's milk; thalassemia.

INTRODUCTION

Iron deficiency anemia (IDA) in children is not merely a low hemoglobin value—it is a signal that oxygen delivery, growth demands, and micronutrient intake have fallen out of balance. The clinical importance is heightened in infancy and early childhood because iron supports neurodevelopment, myelination, and energy metabolism during periods of rapid brain and body growth. While severe IDA is classically associated with pallor, fatigue, and poor appetite, many children present with subtle findings such as irritability, reduced attention, pica, restless sleep, or recurrent infections that are easily attributed to "normal childhood." For this reason, pediatric organizations have long emphasized prevention and early detection through nutrition guidance and screening at defined ages, particularly around 12 months when fetal iron stores are depleted and complementary feeding patterns vary widely [1].

At the same time, clinicians must avoid two common errors: over-testing and under-treating. Over-testing occurs when broad laboratory panels are ordered without a structured pathway, increasing cost and confusion while not improving care. Under-treating occurs when low-dose or short-duration iron is prescribed without absorption counseling or follow-up, leading to "treatment failure" that is actually a systems problem—poor adherence, incorrect dosing, continued high cow's milk intake, or unrecognized blood loss.

MATERIALS AND METHODS

A practical diagnostic pathway begins with risk identification and case-finding, because the pre-test probability of iron deficiency is not equal for all children. Highest-risk groups include infants born preterm or low birth weight, infants with early umbilical cord clamping (in some contexts), children with diets low in iron-rich foods, toddlers with high intake of non–iron-fortified cow's milk, children with chronic inflammatory conditions, and those with intestinal malabsorption or recurrent gastrointestinal blood loss. Nutrition history is not optional—it is often the diagnostic clue. Clinicians should quantify cow's milk intake (often a major driver when >16–24 oz/day), assess iron sources (meat, legumes, fortified cereals), and review feeding transitions (exclusive breastfeeding without supplementation or iron-rich complementary foods). Pediatric guidance on iron deficiency prevention

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 10.25 / 2025

highlights the role of appropriate iron intake in infancy and toddlerhood and recommends screening strategies that combine hemoglobin testing with risk assessment [1].

Screening and initial testing typically start with hemoglobin or hematocrit at around 12 months, with additional testing in children with risk factors at any age. The CDC also communicates to families that anemia screening commonly occurs around 12 months and links this to practical feeding advice, reinforcing the clinical reality that the one-year visit is a key checkpoint [2]. In settings that follow older or more targeted policies, additional screening may be done for high-risk children at 9–12 months, repeated later, and continued periodically in high-risk preschoolers. Importantly, the existence of different national recommendations does not invalidate screening; it highlights the need for clinicians to align practice with local prevalence and resource realities while maintaining a coherent plan. The USPSTF has historically noted insufficient evidence to recommend for or against routine screening in asymptomatic young children, but it still summarizes that high-risk screening strategies are used by major organizations and that treatment dosing is well established [4].

RESULTS AND DISCUSSION

When a child is anemic or borderline, the next step is pattern recognition on the complete blood count. Classic iron deficiency shows microcytosis (low MCV), hypochromia (low MCH), and often an elevated RDW reflecting variable red cell sizes as iron supply fluctuates. However, microcytosis is not specific: thalassemia trait can look similar but typically has a relatively normal RDW and a higher RBC count for the same hemoglobin level. A simple clinic rule is “microcytosis plus high RDW plus diet risk strongly suggests iron deficiency,” but clinicians should confirm with iron studies when possible or use a monitored therapeutic trial when the picture is strongly suggestive and the child is clinically stable. Reticulocyte hemoglobin content, when available, can provide early evidence of iron-restricted erythropoiesis and can respond quickly to therapy, but access varies by setting.

Ferritin is the key confirmatory marker, with an important caveat: ferritin rises in inflammation, infection, and liver disease, so a “normal” ferritin does not always exclude deficiency when the child is acutely ill. Many clinical pathways therefore recommend interpreting ferritin alongside markers of inflammation (such as CRP) or repeating tests when the child is well, especially if the clinical suspicion is high and the diet history fits. Practical institutional guidelines commonly note that low ferritin is diagnostic while also emphasizing its acute-phase behavior and the need for context-based interpretation [3]. Transferrin saturation and serum iron can support the diagnosis but are more variable with recent intake; they are best used as part of a package rather than as stand-alone decision points. The differential diagnosis should also include anemia of chronic inflammation (often normal/high ferritin with low serum iron and low transferrin saturation), lead exposure (which may coexist with iron deficiency), and less common etiologies such as hemolysis or bone marrow suppression when clinical features suggest them.

A highly effective outpatient approach is to use a structured “three-step” treatment pathway for uncomplicated suspected IDA: (1) start therapeutic oral iron at an evidence-based dose, (2) provide absorption and adherence counseling immediately, and (3) schedule an early response check. Modern pediatric reviews describe therapeutic elemental iron dosing commonly in the range of 3–6 mg/kg/day for IDA (higher end for more severe anemia), using ferrous salts as first-line because of bioavailability and cost [5]. Many children tolerate once-daily dosing better than multiple daily doses, and adherence often improves when the regimen is simple; the best dose is the one the family can consistently give. Counseling should be concrete: give iron away from large milk feeds, reduce excessive cow’s milk

ISSN: 2582-4686 SJIF 2021-3.261,SJIF 2022-2.889, 2024-6.875 ResearchBib IF: 10.25 / 2025

intake, consider giving with vitamin C–containing fluids to improve absorption, and warn about expected side effects such as dark stools, constipation, or mild abdominal discomfort. Families should also be told about tooth staining risk with liquid iron and taught simple prevention (use a dropper toward the back of the mouth, rinse/brush afterward).

Monitoring response is the "truth serum" of the pathway. In a child with true IDA who is taking iron, clinicians should expect an early reticulocyte response within about 7–10 days and a hemoglobin rise over the next weeks. A pragmatic outpatient checkpoint is 2–4 weeks: if hemoglobin has not meaningfully improved, clinicians should not reflexively "change to a different iron brand" without troubleshooting.

CONCLUSION

Iron deficiency anemia in children is common, clinically meaningful, and highly treatable when managed through structured diagnostic and treatment pathways. Effective care starts with recognizing risk patterns and using age-appropriate screening anchored around the 12-month visit, then confirming the diagnosis through CBC pattern analysis and iron studies—especially ferritin interpreted in clinical context. Treatment succeeds when clinicians prescribe an adequate elemental iron dose, simplify dosing to improve adherence, counsel families on absorption barriers such as excessive milk intake, and verify response early rather than waiting months to discover failure.

REFERENCES

1. Baker R. D., Greer F. R.; Committee on Nutrition, American Academy of Pediatrics. Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0–3 years of age). – *Pediatrics*. – 2010. – Vol. 126(5). – P. 1040–1050. – URL: https://publications.aap.org/pediatrics/article/126/5/1040/65343/ (accessed: 04.02.2026).

2. Centers for Disease Control and Prevention. Iron | Infant and Toddler Nutrition. – 2024. – URL: https://www.cdc.gov/infant-toddler-nutrition/vitamins-minerals/iron.html (accessed: 04.02.2026).

3. World Health Organization. *Guideline: Daily iron supplementation in infants and children*. – Geneva: WHO, 2016 (digital edition). – URL: https://www.ncbi.nlm.nih.gov/books/NBK362027/ (accessed: 04.02.2026).

4. U.S. Preventive Services Task Force. Iron Deficiency Anemia in Young Children: Screening. – 2015. – URL: https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/iron-deficiency-anemia-in-young-children-screening (accessed: 04.02.2026).

5. Leone G., et al. Iron therapy in pediatric iron deficiency anemia: practical dosing and formulations. – *[PMC article]*. – 2026. – URL: https://pmc.ncbi.nlm.nih.gov/articles/PMC12821433/ (accessed: 04.02.2026).